



The INSTITUTE for BIRD POPULATIONS

MAPS MANUAL

2025 PROTOCOL

INSTRUCTIONS FOR THE ESTABLISHMENT AND OPERATION
OF CONSTANT-EFFORT BIRD-BANDING STATIONS AS PART OF THE
MONITORING AVIAN PRODUCTIVITY AND SURVIVORSHIP (MAPS) PROGRAM

David F. DeSante, Kenneth M. Burton, Danielle Kaschube,
Pilar Velez, Dan Froehlich, Steven Albert

The Institute for Bird Populations
PO Box 518
Petaluma, CA 94953
(707) 789-3224
www.birdpop.org

ACKNOWLEDGEMENTS


We thank MAPS biologists Emma Cox, Eric Feuss, Lauren Helton, Denise Jones, Sara Martin, Nicole Michel, Peter Pyle, Eric (Zed) Ruhlen, Hillary Smith, Brett Walker, Kerry Wilcox, and Mellissa Winfield and research scientists Phil Nott and Jim Saracco for their help in the development of this manual and for comments on earlier versions and revisions. Feedback from the many MAPS station operators also has been invaluable in refining both the manual and the MAPS Program itself.

Funding for the development, establishment, and operation of the MAPS Program has come from many sources. Among the most important are the U.S.D.A. Forest Service, U.S. Department of Defense/Legacy Resource Management Program, U.S.D.I. Fish and Wildlife Service/Office of Migratory Bird Management and National Park Service, U.S. Geological Survey/Biological Resources Division, and National Fish and Wildlife Foundation. We thank these and many other funding sources and private individuals for their support.

Finally, we thank the many hundreds of station operators and all their assistants, interns, and volunteers who, year after year, contribute their data to the MAPS Program. Without their dedication and perseverance, MAPS could not exist.

This is Contribution Number 127 of The Institute for Bird Populations.

Copyright © 2025



Recommended citation: DeSante, D.F., K.M. Burton, D.R. Kaschube, P. Velez, D. Froehlich, and S. Albert. 2025. MAPS Manual: 2025 Protocol. The Institute for Bird Populations, Petaluma, CA.

Contents

INTRODUCTION	1
An Invitation	1
Bird Safety	2
Proper Permitting.....	2
Background and Rationale	3
Design and Objectives of the MAPS Program	4
Analysis of MAPS Data.....	7
ESTABLISHMENT AND GENERAL OPERATION OF MAPS STATIONS	12
Siting a MAPS station.....	12
Establishing a MAPS station.....	14
MAPS terminology	14
General configuration.....	15
Mist nets (number)	15
Mist nets (density).....	15
Mist nets (placement)	16
Mist nets (size, type, and mesh size).....	17
Operating a MAPS station.....	17
Station registration	17
Banding - dates of operation.....	17
Effort data	19
Breeding Status data	19
Habitat Structure Assessment (HSA) data	20
Instructions and data forms.....	20
Recording data and making corrections	20
The use of MAPSNet	20
MAPS data-use policy.....	21
STATION REGISTRATION.....	23
Joining the MAPS Program.....	23
Instructions for completing the MAPS Station Registration Form	23
MIST-NETTING EFFORT	27
Operation of nets	27
Frequency of mist-netting effort.....	28
MAPS Periods and Intended Periods	28
Making up missed effort	29
Minimum allowable effort	30
STANDARD NET OPENING AND CLOSING TIMES.....	31
Instructions for completing the Standard Net Opening and Closing Times form	31
SUMMARY OF EFFORT	33
Instructions for completing the Summary of Mist-Netting Effort form	33
COLLECTION AND RECORDING OF BANDING DATA	36

General procedures for recording banding data.....	36
Primary MAPS data	36
Supplemental data	36
Optional data	37
Codes, scales, and forms.....	37
Multiple-station locations	37
Non-MAPS data	37
Page headings and other notations	38
New bands	38
Lost and destroyed bands	38
Recaptures.....	38
Changed bands.....	39
Added bands.....	39
Unbanded birds	39
Mortalities.....	39
Banding-data fields	40
BANDER'S INITIALS.....	40
CODE	40
BAND NUMBER.....	40
SPECIES NAME.....	41
SPECIES ALPHA CODE.....	41
AGE	42
WRP.....	45
HOW AGED.....	49
SEX	50
HOW SEXED.....	51
SKULL	51
CL. PROT.....	53
BR. PATCH.....	54
FAT	55
BODY MLT.....	57
FF MOLT	57
FF WEAR	58
JUV. BDY PL.	59
MOLT LIMITS & PLUMAGE	60
WING	64
BODY MASS.....	64
STATUS.....	64
DATE (MO/DAY)	64
CAPTURE TIME.....	64
STATION	64
NET	65
DISP	65

NOTE NUMBER	65
FTHR. PULL	65
NOTE	66
A strategy for ageing and sexing birds.....	66
SUMMARY OF MIST-NETTING RESULTS	68
Instructions for completing the Summary of Mist-Netting Results form	68
BREEDING STATUS LIST	70
PERIOD STATUS	72
YEAR STATUS	73
OBSERVER'S INITIALS.....	77
HABITAT STRUCTURE ASSESSMENT (HSA)	78
HABITAT TYPES	78
A STEP-BY-STEP APPROACH:	79
Habitat Structure Assessment form H1	82
DATE	82
HABITAT DESIGNATION	82
HABITAT NAME	82
PERCENTAGE OF STATION COMPRISED OF THIS HABITAT TYPE	82
NATIONAL VEGETATION CLASSIFICATION STANDARD DIVISION.....	82
NATIONAL VEGETATION CLASSIFICATION STANDARD ALLIANCE.....	85
GENERAL DESCRIPTION OF HABITAT TYPE INCLUDING HABITAT AGE	85
VEGETATIVE LAYERS PERCENT COVER.....	85
AVERAGE HEIGHT OF TREE CANOPY	86
AVERAGE HEIGHT OF SHRUBS	86
AVERAGE HEIGHT OF HERBACEOUS VEGETATION	86
NUMBER OF SNAGS	86
GEOGRAPHIC FEATURES.....	86
MANAGEMENT/DISTURBANCE HISTORY	87
DATA SUBMISSION	88
What data to submit	88
How to submit MAPS data	88
Submitting data through MAPSNet	88
Submitting paper copies of banding data	90
Due date	91
LITERATURE CITED.....	93

Figures

Figure 1. Map of the continental U.S. and Canada showing the eight MAPS Regions.	5
Figure 2. Diagram of an idealized MAPS station.	16
Figure 3. Recommended starting periods for MAPS stations.	18
Figure 4. Completed MAPS Station Registration Form.	26
Figure 5. Completed MAPS Standard Net Opening and Closing Times form.	32
Figure 6. Completed MAPS Summary of Mist-Netting Effort (page 1).	35
Figure 7. Completed MAPS banding-data sheet for Band Size '0' (front).	43
Figure 8. Note section for example banding page.	44
Figure 9. The Life Cycle of a Typical Migratory Songbird, Including Age and Common WRP Codes.	48
Figure 10. Completed MAPS Summary of Mist-Netting Results form.	69
Figure 11. Completed MAPS Breeding Status List losing their BPs and CPs.	74
Figure 12. Completed Station Map (truncated)	80
Figure 13. Completed MAPS HSA Form H1	84

Tables

Table 1. Adult and young MAPS superperiods.	30
Table 2. Species alpha codes for gallinaceous birds.	41
Table 3. Hierarchical categories of Period Breeding Status	72
Table 4. MAPS Banding-data file structure for 2025 data	91

INTRODUCTION

An Invitation

Welcome to the Monitoring Avian Productivity and Survivorship (MAPS) Program! MAPS is a cooperative effort among public agencies, private organizations, and the bird banders of the continental United States, Canada, and Mexico to provide critical, long-term data on population and demographic parameters for over 150 target landbird species at multiple spatial scales. As part of the MAPS family, you team with hundreds of private individuals, and workers from federal and state agencies and non-governmental organizations to gather important data for the conservation of birds and their habitat.

The MAPS Program utilizes standardized, constant-effort mist netting and banding during the breeding season at an extensive network of stations. The MAPS methodology provides annual indices of adult population size and post-fledging productivity from data on the numbers and proportions of young and adult birds captured; and annual estimates of adult survivorship, adult population size, proportion of resident individuals in the adult population, recruitment into the adult population, and population growth rate (λ) from mark-recapture data on adult birds. IBP and our collaborators use this data to study the causes of population changes in North America's landbirds. This manual (and all the forms associated with operating a MAPS station) is available for download on the MAPS web page <http://www.birdpop.org/pages/maps.php>.

Any public agency, private organization, or independent bird bander currently operating or able to establish one or more banding stations operated regularly through the breeding season is encouraged to participate in the MAPS Program. All that is required is the standardized operation of a series of about ten nets at permanent sites on only one day during each of six to ten consecutive ten-day periods between May and August.

Operating a MAPS station is relatively simple, but also a substantial commitment. Standardization from year to year and continuation of the study **for at least five consecutive years** at each station are necessary to provide reliable productivity indices and survivorship estimates ("vital rates"). Continuation of the study **for ten to twenty consecutive years** at most stations will likely be necessary to obtain reliable trend information on these critical vital rates. This manual is designed to guide you through all the steps involved in operating a MAPS station and address any questions. Everything contained herein is important; take the manual with you on every visit to your station and, **please, read and use it.**

The Institute for Bird Populations is excited about the possibility of working with you to monitor the productivity, survivorship, and population trends of North American landbirds. We cordially invite you to join the MAPS Program. The methodology outlined below may seem formidable at first glance. It is, however, relatively simple: standardized mist netting and banding during the breeding season, documentation of the apparent breeding status of the birds present at the station, and preparing a simple habitat map and habitat structure assessment. In addition, IBP provides technical assistance and guidance year-round to answer your questions and guide you through the process.

2 - 2025 MAPS Manual

Furthermore, the MAPS Program requires the operation of nets on only six to ten days during the breeding season. Thus, the effort required to gather these extremely valuable data on the vital rates (productivity and survivorship) of landbirds is quite manageable. We invite you to become an important part of this exciting, cooperative endeavor by establishing one or more MAPS stations in your area.

Bird Safety

The protocols and objectives outlined in the following pages are designed to collect valuable data for bird conservation. However, **protocols should never be followed at the expense of bird or human safety**. As a responsible bander, if safety is ever a concern, you should suspend protocols until the problems are addressed. Please ensure that all of the banders at your station know and practice safe banding techniques.

We encourage you to review ethical banding and safety procedures with your crews in depth at least once a year and always strive towards a safer banding experience. We recommend reviewing materials such as those provided by the North American Banding Council (NABC; www.nabanding.net), The Ornithological Council (*Guidelines to the Use of Wild Birds in Research*; https://birdnet.org/wp-content/uploads/2022/09/guidelines_august2010.pdf), *The Mist Netter's Bird Safety Handbook* available from IBP and the [2013](https://www.birdpop.org/pages/mapsChatArchive.php) and [2018](https://www.birdpop.org/pages/mapsChatArchive.php) MAPS Chats (<https://www.birdpop.org/pages/mapsChatArchive.php>) which provide bird safety suggestions for keeping your station as bird safe and smooth running as possible.

While bird safety is important, we must keep our banding crews safe. While most bird diseases are not regularly transmitted to humans, it is good to know that diseases such as Avian Flu H5N1 exist in the wild bird population. These diseases seem most likely to infect those who work with larger flocks of domestic birds, such as chickens, ducks, and geese. However, as passerine banders, there is a minimal risk that we could handle birds carrying the virus. At the time of the release of this manual, there are no official recommendations for protective gear while banding passerines, but the bird banding offices of the United States or Canada recommend that banders regularly wash their hands with soap and water or use hand sanitizer. And avoid touching their faces, especially their noses and eyes, before washing their hands.

Proper Permitting

All banders applying to operate or currently operating MAPS stations must adhere to all federal and state permitting requirements. Check that you have addressed these issues before beginning banding each season, and that relevant permits are current. **Note:** A special addendum is required on your federal banding permit to allow you to pull feathers. If you plan to participate in the cooperative Bird Genoscape feather sampling study (birdgenoscape.org) or similar studies, please ensure that your permits include these special permissions. See <http://www.birdpop.org/pages/mapsDataForms.php> for the latest information on feather sampling.

Background and Rationale

Earth's biosphere and its landbird populations are facing a growing number of environmental threats of ever-increasing severity, many of which, such as climate change, habitat loss, invasive species, and toxic pollution, are global in scale (Brown 1991). It is not surprising, therefore, that a number of large-scale, long-term monitoring programs for landbirds were already in place on this continent before the MAPS Program started. They include the Breeding Bird Survey (BBS), the Breeding Bird Census and Winter Bird Population Study, and the Christmas Bird Count. All of these efforts provide annual information on landbird populations, and many of the resulting trends indicate serious population declines in many species, including forest- and scrub-inhabiting Nearctic-Neotropical migrant species (Robbins *et al.* 1989, Terborgh 1989) and grassland species (Knopf 1994). These population declines prompted the establishment of the Neotropical Migratory Bird Conservation Initiative, "Partners in Flight" (PIF), to which most federal agencies and many state and private organizations have become signatories

The above-mentioned monitoring efforts, however, all fail to provide data on the primary demographic parameters or vital rates (productivity and survivorship) of landbirds. Without these critical data, it is difficult if not impossible to test competing hypotheses to account for observed population changes, or to determine the stage(s) in the life cycle at which these population changes are taking place; that is, whether the changes are being driven by causal agents that affect birth rates or death rates or both (DeSante 1992). Efforts that monitor only avian population trends have generally been unable to determine to what extent habitat destruction and degradation (e.g., deforestation and forest fragmentation) on the temperate breeding grounds, versus that on the tropical wintering grounds, are causes for declining populations of neotropical migratory landbirds (Wilcove 1985, Holmes and Sherry 1988, Hutto 1988, Morton and Greenberg 1989, Peterjohn *et al.* 1995).

An integrated approach to monitoring primary demographic parameters and secondary population trends of landbirds is critical for determining causes of population changes and identifying management actions and conservation strategies to reverse population declines (Baillie 1990). Perhaps even more importantly, this approach aids in evaluating the effectiveness of the management actions and conservation strategies actually implemented (DeSante 1995). This is because environmental stressors and management actions affect primary demographic parameters directly and usually without the buffering or time lags that often occur with secondary population trends (Temple and Wiens 1989). Monitoring the vital rates of landbirds also allows models to be constructed regarding the viability of their populations. Habitat- and landscape-specific data on vital rates provide a clear index of habitat and landscape quality, and allow identification of habitat and landscape conditions that provide source populations and that influence population sinks (DeSante and Rosenberg 1998). An increase in demographic monitoring has been called for by the Monitoring Working Group of PIF since 1992 (Butcher and Droege 1992), and an argument for basing avian management on vital rates has been provided by DeSante *et al.* (2005).

4 - 2025 MAPS Manual

In 1989, The Institute for Bird Populations (IBP) anticipated these monitoring needs and created the Monitoring Avian Productivity and Survivorship (MAPS) Program, a cooperative effort to operate a continent-wide network of constant-effort mist-netting stations to capture and band landbirds during the breeding season (DeSante 1992, DeSante *et al.* 1993, 1995). The MAPS Program was patterned on the British Constant Effort Sites (CES) Scheme which since 1981 has been one of the cornerstones of the British Trust for Ornithology's Integrated Population Monitoring Programme (Baillie *et al.* 1986, Baillie 1990, Peach *et al.* 1996) and has inspired at least 15 other European CES efforts (Robinson *et al.* 2009). The first three years of MAPS was an IBP-sponsored feasibility study, during which time the program grew from 16 stations in 1989 to 66 stations in 1991 and the MAPS protocol became standardized. MAPS was endorsed in 1992 by the Monitoring Working Group of PIF and a four-year pilot study was sponsored by the Migratory Bird Management Office of the U.S. Fish and Wildlife Service (sponsorship later transferred to the Biological Resources Division of the U.S.G.S.).

The number of stations grew dramatically in subsequent years to nearly 500, primarily through the involvement of the Department of Defense (Legacy Resource Management Program) and U.S.D.A. Forest Service.

In 1996, the MAPS methodology underwent an extensive peer review. Some of the conclusions of this review were that "MAPS is technically sound and based on the best available biological and statistical methods" and that "MAPS complements other land bird monitoring programs such as the BBS by providing useful information on land bird demographics that is not available elsewhere" (Geissler 1997). A summary and analysis of the MAPS methodology were provided by DeSante *et al.* (2004a) and Burton and DeSante (2004). MAPS has continued to expand since 1995 to some 300-400 stations operated yearly during 2001-2025.

Design and Objectives of the MAPS Program

MAPS is organized around several monitoring, research, and management objectives: to provide (a) annual **estimates** of adult survival rate, adult population size, proportion of residents in the adult population, recruitment into the adult population, and population growth rate (λ); and (b) annual **indices** of adult population size and post-fledging productivity. MAPS provides these population and demographic indices and estimates for over 150 landbird species that are well-distributed among various migration-strategy, foraging-strategy, nest-location, and habitat-preference guilds. In addition, MAPS works at multiple spatial scales, from program-wide (essentially the entire continent north of Mexico), MAPS Regions (Fig. 1), Bird Conservation Regions (BCRs), or BBS Physiographic Strata, to small scales, such as clusters of stations, on a single national forest, park, or military installation, or local landscapes surrounding single stations (e.g., four-km radius areas).

MAPS Regions have been defined by dividing the continent into eight major areas: Alaska, Boreal & Arctic Canada, Northwest, North-central, Northeast, Southwest, South-central, and Southeast (Fig. 1). These delineations generally follow the boundaries of BBS Physiographic Strata and are based on both biogeographic and meteorological




Figure 1. Map of the continental U.S. and Canada showing the eight MAPS Regions.

considerations, including the apparent periodicity of the Jet Stream. Seasonal weather tends to be similar at many locations within a given region, but often varies considerably among regions. Population and demographic parameters will likely vary in a similar manner at many banding stations within a region, at least to the extent to which weather conditions influence them.

The achievement of the **monitoring objectives** of MAPS has been well documented in MAPS reports (DeSante *et al.* 1996, 1998, DeSante and O'Grady 2000, DeSante and Kaschube 2006, 2007, 2009) and on the <https://ibp-maps-data-exploration-tool.org/app/maps> web application, which was introduced in 2023. The website updates results provided by [VitalRatesofNorthAmericanLandbirds.org](https://vitalratesofnorthamericanlandbirds.org) (DeSante *et al.* 2015) by incorporating additional years of data (now 27 years of data [1992-2018] compared to 15 years in DeSante *et al.* 2015) and applying more recently developed Bayesian hierarchical models that provide region (Bird Conservation Regions) × year specific estimates of demographic parameters, as well as region-specific estimates of trends and annual variation in these parameters. We present results for five demographic parameters: 1)

6 - 2025 MAPS Manual

adult abundance, 2) juvenile abundance, and 3) productivity indices from generalized linear mixed models of capture data; and 4) adult apparent survival and 5) residency probabilities from Cormack-Jolly-Seber (CJS) models of capture-recapture data.

The **research objectives** of MAPS are to identify and describe: (a) temporal and spatial patterns in the demographic indices and estimates provided by MAPS (The Institute for Bird Populations 2015, DeSante *et al.* 2015) and (b) relationships between these temporal and spatial patterns and (1) ecological characteristics of the target species (e.g., migration strategy, nest location), (2) population trends of the target species (e.g., areas or locations with increasing or decreasing trends) (The Institute for Bird Populations 2015, DeSante *et al.* 2015), (3) station-specific and landscape-level habitat characteristics (e.g., total forest cover, mean forest patch size), and (4) spatially-explicit weather data (e.g., mean, min, and max temperature or precipitation, extreme events).

MAPS allows these objectives to be met for multiple species at multiple spatial scales. Achievement of these research objectives is providing empirical information regarding life-history strategies and other topics of interest to avian ecology (DeSante *et al.* 1999, DeSante 2000), including the effects of global climate cycles on avian productivity (Saracco *et al.* 2019, Nott *et al.* 2002). A future important research effort will be to integrate extensive count data (for example, from E-bird and the Avian Knowledge Network) with Cormack-Jolly-Seber capture-mark-recapture (CMR) data from MAPS and extensive remote-sensed environmental data in an effort to improve the predictive accuracy of demographic rates and abundance through space and time (Saracco *et al.* 2009a). This type of analysis has already been completed, applying integrated population models (IPMs) to MAPS and North American Breeding Bird Survey (BBS) data to better assess the driving forces behind a declining warbler species (Saracco and Rubenstein 2020).

In addition, however, by analyzing the relationships between spatial variation in population trends (using for example, BBS data) and the vital rates that drive those trends, we have been able to make inferences regarding the proximate demographic cause(s) of population decline, that is, to suggest whether the decline is caused by low productivity or low survivorship (DeSante *et al.* 2001). This, in fact, is the **first management objective** of MAPS, an objective that can be achieved by no other North American avian monitoring program (Saracco *et al.* 2008).

The **second management objective** of MAPS is to determine the ultimate (environmental) causes of population trends and to identify and formulate landscape-scale management actions and conservation strategies to reverse population declines and maintain stable or increasing populations. We do this by modeling vital rates (productivity indices and survival and recruitment estimates) as functions of landscape-scale and site-specific habitat characteristics and spatially-explicit weather and climate variables to identify habitat characteristics and weather variables that exert strong effects on the vital rates of landbird species, especially species of conservation concern. Management prescriptions developed this way for species for which productivity is critical for maintaining stable populations, involve modifying habitat characteristics from those associated with low productivity to those associated with higher productivity. Examples of these types of analyses of MAPS data, that are allowing us to achieve our

second major management objective, are the focus of the management guidelines and conservation strategies that we have developed for reversing declines of landbirds of conservation concern on DOD installations in southeastern United States (Nott 2000, Nott *et al.* 2003a) and on national forests in the Pacific Northwest (Nott *et al.* 2005). We are developing decision support tools for managers, whereby they can determine the effects of their proposed management actions on the vital rates of species of concern. One way to increase populations is to increase productivity, and MAPS data has been used to determine anthropogenic noise effects on productivity, which will be important to determine which areas are important to conserve (Ng *et al.* 2020)

The **third management objective** of MAPS is to evaluate the effectiveness of the management actions and conservation strategies through the adaptive management process. If the goal is to manage for increased productivity (as is the implicit goal of most breeding-ground management), it is imperative to monitor productivity. With wetter springs in recent years in the grasslands of Nebraska, longer grasses have caused declines in Grasshopper Sparrow, but MAPS data has shown that managing these grasslands with prescribed fire can help mitigate lowered productivity (Glass *et al.* 2020). Measuring the age structure of the population, as provided in MAPS data, has also given important clues to habitat quality, with more yearling birds being present in higher numbers in reclaimed areas compared to undisturbed areas (Pyle *et al.* 2020).

MAPS' three management objectives can be achieved for multiple species at appropriate spatial scales.

Analysis of MAPS Data

IBP researchers have also completed a number of analyses and evaluations of the program in various geographical areas and landholdings. This effort began with a general analysis of the results of the first ten years (1992-2001) of the MAPS program in Alaska and adjacent Canada (DeSante *et al.* 2003a), followed by an analysis of the statistical power to detect temporal trends and spatial differences in survival of landbirds breeding in Alaska and adjacent Canada (DeSante *et al.* 2003b). These were followed by an evaluation of the data collected at MAPS stations operated on National Wildlife Refuges in the USFWS Pacific Region (DeSante *et al.* 2004b), which led to an expanded evaluation of the MAPS Program on all stations in the Pacific Northwest (DeSante *et al.* 2005). Broadening our scope, we then conducted general analyses of the statistical power to detect temporal trends and spatial differences in survival from CMR models (DeSante *et al.* 2009), and applied those models to MAPS data in each MAPS region and at the continental scale to develop a vision for expanding MAPS and integrating it into Coordinated Bird Monitoring all across North America (Saracco *et al.* 2006, DeSante and Saracco 2009). Most recently, we built upon that continental vision and provided detailed recommendations for integrating MAPS into Coordinated Bird Monitoring in the Northeast, i.e., USFWS Region 5 (DeSante *et al.* 2008). We are currently seeking resources to allow us to continue these latter analyses and provide analogous detailed recommendations for the remainder of the United States and Canada. This has become critically important as state and federal agencies and non-governmental organizations seek to develop bird conservation plans to deal with the huge threats

8 - 2025 MAPS Manual

posed by climate change. The assessment and monitoring of avian vital rates using MAPS may provide one of the optimal resources to model and predict the effects of climate change on landbird populations, guide adaptation and conservation efforts to mitigate those effects, and evaluate the effectiveness of those efforts. Annual **indices** of adult population size and post-fledging productivity (i.e., reproductive index, the ratio of young to adults in the catch) are calculated for each species from the numbers of young and adult birds captured, which are pooled over all the stations. The statistical significance is inferred from confidence intervals calculated from the standard errors of the mean percentage changes in these indices for species captured at the stations. Long-term changes in numbers of young and adult birds are assessed through application of a generalized linear ("log-linear") model (GLM) appropriate for Poisson-distributed response variables, which was used to estimate adult population size and productivity trends. Poisson models allow for a trend in variance to match the trend in mean numbers. The number of adults modeled with net-hours incorporated as an offset in a (Poisson-distributed) GLM of trend over time. Similarly, the trend in productivity is modeled as the total number of young captured, with the number of breeding adults used as an offset in the (Poisson-distributed) GLM.

Annual **estimates** of adult survival rate, proportion of residents in the adult population, recruitment into the adult population, and population growth rate (λ) are obtained for each species from modified CMR analyses. Major advances have been made in both the theory and application of data from CMR experiments (Pollock *et al.* 1990, Lebreton *et al.* 1992). These advances provide for increased precision in the resulting estimates and also allow spatial, age, and/or time dependence in the estimates of survival and recapture rates to be assessed, permit some parameters to be set equal to fixed *a priori* values, and allow any of the parameters to be related to external variables (Clobert *et al.* 1987). This approach was initially applied to mark-recapture data from both Great Tits and Black-headed Gulls in Europe (Clobert *et al.* 1987), and from Sedge and Reed Warblers in Britain (Peach *et al.* 1990, 1991) and has become a major analytical tool. By using reverse-time analysis of CMR data, models have been developed to permit estimation of recruitment and population growth rates (Pradel 1998). In addition, models have been developed to account for the negative bias of transient (non-resident) individuals on survival-rate estimates and to estimate the proportion of resident individuals among newly captured adult birds (Pradel *et al.* 1997, Nott and DeSante 2002, Hines *et al.* 2003). These transient models are incorporated into the capture-mark-recapture analyses of MAPS data (e.g., Rosenberg *et al.* 1999) and can be utilized through the computer programs SURVIV (White 1983), TMSURVIV (Hines *et al.* 2003), and MARK (White and Burnham 1999).

In order for the above mentioned analyses to be run, there need to be sufficient numbers of captures. To be able to attain minimally precise estimates of survivorship, we need six years of data and approximately four captures, and one recapture, per year of a particular species. To attain more precise estimates, six years of data and approximately 15 captures, and one recapture, per year of a particular species (Kaschube *et al.* 2022). This is fairly easy to accomplish when pooling data between stations, but can be difficult when only looking at one station.

More recently, using both reverse time (Pradel 1998) and transient (Pradel *et al.* 1997, Nott and DeSante 2002, Hines *et al.* 2003) CMR models, we can examine relationships between spatial variation in MAPS trend estimates (λ) and spatial variation in critical vital rates including adult apparent survival, recruitment, productivity, and first-year survival. As a pilot analysis, we assessed the demographic contributions of adult survival and recruitment rates to BCR-scale variation in MAPS population trends for 27 Nearctic-Neotropical migratory species using 12 years (1992-2003) of MAPS data (Saracco and DeSante 2008). We found that recruitment tended to be about three times as important as adult survival in driving spatial variation in population trends, but that first-year survival tended to be more important than productivity in driving recruitment. By looking at the overall (program-wide) MAPS population trends for these 27 species, however, we found that: first-year survival was the most important driver of spatial variation in trend for species with significant population declines, both first-year and adult survival were the most important drivers of spatial variation in trend for species with significant population increases, and productivity was most important driver of spatial variation in trend for species with non-significant (relatively stable) population trends.

Using these same methods, we examined both annual and BCR-scale spatial variation in the vital rates of 158 species of North American landbirds using 15 years (1992-2006) of MAPS data, and presented the results on the Vital Rates of North American Landbirds website (DeSante *et al.* 2015). For each of these 158 species, the site presents 1) estimates of population change (λ), adult apparent survival, recruitment, and residency, along with indices of productivity, post-breeding effects, and adult population density; 2) graphs showing annual variation and maps showing spatial variation in these vital rates; and 3) the results of temporal and spatial pairwise correlations among these vital rates. The website also includes species account narratives for these 158 species that provide hypotheses regarding the proximate demographic drivers of the observed temporal and spatial variation in their population changes, as well as suggestions as to research and management efforts to reverse population declines and maintain stable or increasing populations. These results also suggest that enhancing the survival of both adult and, especially, first-year birds must be a very important conservation strategy for slowing population declines and achieving stable populations. Because both first-year and adult survival of migratory species may be driven primarily by processes acting on the wintering ranges and migration routes of these species, identifying relationships between these vital rates and both habitat characteristics and weather on the non-breeding grounds may well be critical for successful conservation of migratory landbirds. MAPS data, used in conjunction with data from the overwintering period provided by the MoSI (Monitoreo de Sobrevivencia Invernal - Monitoring Overwintering Survival) and additional information on migratory connectivity can provide insights regarding the mechanisms whereby survival throughout the year can drive populations trends of migratory birds (Saracco *et al.* 2009b).

In cooperation with researchers at the USGS Patuxent Wildlife Research Center, we have incorporated Bayesian hierarchical spatial autoregressive models to better describe spatial variation in adult apparent survival rates and residency probabilities. Results from

10 - 2025 MAPS Manual

these models are presented on the MAPS Data Exploration tool: <https://ibp-maps-data-exploration-tool.org/app/maps>. The first iterations of these analyses were completed for American Robin, Wood Thrush (Saracco et al. 2010), and Common Yellowthroat (Saracco et al. 2011). Now, the exploration tool presents data analysis for 50 of the most commonly captured species in North America using 27 years of data (1992-2018) (IBP 2023). These spatial models represent a significant advance over approaches to investigating spatial patterns in vital rates that aggregate data at coarse spatial scales (such as the BCRs described above) and do not explicitly incorporate spatial information in the models. They tend to overcome, to some extent, difficulties caused by geographical areas (or BCRs) with sparse data and by the non-random distribution of stations, can easily accommodate missing data within the modeling framework, and permit MAPS data and results to be included in models and analyses based on any previously or subsequently established grid system. These important papers, which have appeared in *Ecology* (Saracco et al. 2010) and the *Journal of Ornithology* (Saracco et al. 2012), indicate that residency probability is often more spatially heterogeneous than survival and not positively spatially correlated with survival. They thus illustrate the importance of understanding the role of transients in local populations.

MAPS data has also helped determine how to truly define what a "population" is for a bird species (Rushing et al. 2016). Many conservation plans and management strategies work at the level of the population, so this delineation is important, but should the population be defined geographically or biologically? In cooperation with researchers at the Smithsonian Institution, an approach was developed for using Breeding Bird Survey data to quantify geographic structure in trend and abundance, and identify distinct natural populations for eight species of passerines. The researchers then used vital rates derived from MAPS data to independently validate their method of population delineation.

The MAPS program has also worked in cooperation with other programs to utilize the continent-wide range of stations to further other avian research. Using feather samples collected by MAPS cooperators, scientists from the Bird Genoscape Project isolated DNA and stable isotopes to map the connectivity of the breeding grounds to the wintering grounds. At the time of the release of this version of the MAPS manual, 16 species have completely mapped genoscapes, with the mapping of an additional 16 species in progress. (<https://www.birdgenoscape.org/species-list/#completedgenoscapes>). The first map was done on Wilson's Warbler (Ruegg et al. 2014). Using this genetic data, six distinct populations of Wilson's Warbler were delineated, allowing scientists to concentrate conservation efforts in areas where declining populations breed, migrate, and winter (Ruegg et al. 2020). Colorado State University scientists utilized MAPS data to determine the impact of West Nile virus on survival rates of 49 species, determining that for some landbird species the effect is short-lived for some, but is persistent in others (George et al. 2015).

Exciting results have also come from a cooperative effort with UCLA. Dr. Casey Youngflesh and colleagues analyzed MAPS data and found that temperature and elevation affect the size and shape of birds (Youngflesh et al. 2022). They found that within species, smaller body size was associated with warmer temperatures over space

and time. Individuals breeding at lower latitudes are smaller, and as the climate warms, individuals at a given latitude are getting smaller over time. The researchers also found that within species, wing length increased in relation to body size as breeding elevation increased. And, using the same dataset along with weather data it was determined that spring is arriving earlier and birds are responding by breeding earlier, but not at the same rate (Youngflesh et al. 2022). This phenological mismatch can cause reduced production of young, which in turn will cause declines in bird populations.

Even though MAPS is a breeding season program, IBP scientists have also used MAPS data to explore migration strategies of landbirds and to better understand molt migration (Pyle et al. 2018).

We hope we have conveyed to you the importance of demographic monitoring and the value of the MAPS Program. Each year, as more data accumulate from established stations and as additional stations are established in new areas and new landscapes, the power of the data for revealing spatial and temporal patterns in landbird demographic parameters, and thus their usefulness for avian conservation, increases dramatically. We also hope that we have conveyed some of the excitement and intense commitment we feel regarding the role of MAPS in North American landbird conservation. Again, we invite you to participate in this growing cooperative effort. But remember, MAPS may not be for everyone. Yet, if your heart is thrilled by holding in your hands the life of a precious warbler, thrush, or bunting, and knowing that you are providing data that will aid the survival of its species, then maybe MAPS is for you!

ESTABLISHMENT AND GENERAL OPERATION OF MAPS STATIONS

The following guidelines for the establishment and operation of MAPS stations will optimize the usefulness of data obtained from MAPS stations. These guidelines conform to those recommended for constant-effort mist netting in Ralph *et al.* (1993), and were discussed in DeSante *et al.* (2004a). Because a major objective of the program is to generate estimates of temporal variation in productivity and survivorship, standardization in station operations from year-to-year and station continuity over a number of years are critical. Continuity is also important for minimizing population-parameter fluctuations that may result from year-to-year changes in the geographic distribution of stations. We realize that, because of vagaries of weather and other uncontrollable factors, no station will be able to achieve perfect standardization. Nevertheless, every attempt should be made to follow these guidelines as closely as possible.

Although standardization and continuity are critical components of the MAPS Program, the first year of operation at a MAPS station should be considered a pilot year; nevertheless, all data from the first year should be submitted. Station boundaries and net sites may be shifted during or after the first field season if problems arise or net sites prove to be unproductive. Any such changes must be documented and reported, and if possible, no further changes ought to be made after the start of the second field season. If net sites are changed, the new net designations must differ from those of the discontinued sites.

Siting a MAPS station

It is important to keep in mind that the productivity indices generated at a MAPS station provide a landscape-level, rather than site-specific, measure of productivity. This is because the young birds captured by the MAPS protocol include many dispersing individuals from the surrounding landscape, as well as a few individuals that may have fledged from nests within the boundaries of the 20-ha MAPS study area. Data on the dispersal characteristics of young and adult birds after the breeding season but before fall migration are just now being obtained for a very few species from radio-telemetry studies (Anders *et al.* 1997, Vega Rivera *et al.* 1998). These studies suggest that the landscape from which the dispersing young originate may be on the order of several thousand hectares (perhaps about 10,000-12,000 acres). Although management actions occur on a site-specific basis, their effects on bird populations become pronounced only when the specific management actions occur over substantial portions of the landscape. The ability of MAPS to provide landscape-level information on productivity is one of the unique strengths of the program. Thus, when siting MAPS stations to investigate the effects of a particular habitat type or management action, it is important to consider the habitat type or management characteristics of the overall landscape, that is, of the area within perhaps a four-km (2.5-mile) radius of the station.

It is also important to consider site-specific habitat characteristics when siting MAPS stations, as these can influence the extent to which dispersing young and adult birds concentrate there. Recent work on forest-breeding species suggests that many individuals, both young and adult, desert forest interior locations immediately after the young attain independence from their parents and disperse to edge locations to molt and stage before initiating fall migration (Anders *et al.* 1997, Vega Rivera *et al.* 1998). These edge locations are generally characterized in mid to late summer by dense cover and an abundance of food resources, often fruit. Indeed, we have found that mid-to-late-summer capture rates of both adults and young are much lower at forest-interior MAPS stations than at stations that contain forest-edge or scrub habitat (DeSante 1996). Because productivity indices are calculated from the proportion of young in the mid-to-late-summer catch, the precision of the resulting indices will tend to be lower at forest-interior stations than at stations containing edge and scrub habitats. On the other hand, because late spring and early summer capture rates of breeding adults are often high at forest-interior stations, especially for forest-interior species, such stations can provide important data for estimating adult survival rates.

As mentioned above, the goals of MAPS include identifying and describing spatial and temporal patterns in demographic parameters; relating these to species-specific population trends and life history strategies, habitat characteristics, and weather variables; and using the resulting relationships to formulate management strategies for reversing population declines. As such, MAPS stations are often sited under some hypothesis-driven sampling strategy. Although we appreciate that MAPS stations can only be sited where long-term standardized mist netting is practical and permissible, the value of the data for testing hypotheses can often be enhanced if some elements of a probability-based sampling strategy can be incorporated into the siting of stations. For example, assume you are able to establish three stations in a nearby state park or forest and are particularly interested in upland oak-hickory habitat. A promising strategy might be to lay a suitably scaled grid over a GIS layer showing the distribution of upland oak-hickory habitat within the park or forest, and randomly select 10-12 grid points that lie within landscapes comprised primarily of that habitat. Then, examine the immediate landscape around each grid point and try to identify a suitable MAPS-station site within about one km of the point. If none is available around the first point, go on to each successive point until three suitable sites have been identified. A suitable site would include an area of about 20 ha (50 acres) within which long-term mist netting is both practical and permissible and that lies at least partially in upland oak-hickory habitat, but with some edge or scrub habitat as well. Remember, most of the dispersing young and adult birds that will contribute to productivity indices will originate from the surrounding landscape rather than from within the station itself. In contrast, the breeding adults that will contribute to survivorship estimates will originate from the station itself.

With these concepts in mind, we offer the following guidelines for siting stations:

- (1) If possible, try to use some elements of a probabilistic sampling strategy to site stations within the selected landscape.

- (2) Within the selected landscape, stations should be established at sites that are expected to remain accessible and free of major anthropogenic disturbance for at least

five (preferably ten) consecutive years. Note that there can be disturbance, even heavy disturbance, in the surrounding landscape. If there is disturbance at the station (or in the landscape), it should be described through the Habitat Structure Assessment (see pg. 24).

(3) Stations should be sited where substantial numbers of individuals of many of the common species breeding in the area, or of a particular target species, can be captured.

(4) In order to capture large numbers of dispersing young and adult birds, stations should contain some edge habitat, such as a forest edge, riparian corridor, montane meadow, or power-line right-of-way. Stations can be sited entirely in forest-interior situations, but capture rates at such stations will likely be low.

(5) The habitat types at the station should be fairly representative of those present in the surrounding landscape. Stations not representative of the landscape or at which large numbers of transient or migrating birds concentrate (such as narrow points of land jutting into large bodies of water, or isolated oases in desert or grassland habitats) should be avoided.

(6) Because the derived population and demographic parameters are likely to be highly sensitive to successional changes in the habitats sampled, stations generally should be sited in relatively mature habitats or where the habitat is held in a lower successional stage by active management. The latter type of station is particularly desirable for the long-term monitoring of scrub- and/or second-growth-inhabiting species. Stations sited in highly-successional habitats must be indicated as such so that habitat change can be factored into analyses.

(7) In order to ensure standardization, MAPS stations may not incorporate any artificial food or water sources such as feeders, compost piles, dumps, birdbaths, fountains, and livestock pens. Audio playback calls should also not be used.

If you have the resources available to establish and operate more than one station, it might be advantageous to select two (or more) sites of similar habitat within a few kilometers (but further than one kilometer) from each other. Such an arrangement would give greater precision to the population-parameter estimates for that habitat in that region and might allow us to examine the extent of local dispersal and site fidelity.

Establishing a MAPS station

MAPS terminology: A MAPS “**station**” is a discrete study area consisting of a number of net sites (“**nets**,” the exact places at which nets are located). Each station is given a name and a four-character code (e.g., Copper Creek = COPP or COCR). Upon receipt of a station’s first data submission, we will also assign the station a unique, five-digit station number. Each station is part of a “**location**” that may contain other stations in the same general area (e.g., on the same national forest, national park, military installation, or nature reserve) operated by the same individual or organization. Each location is identified by a four-character code (e.g., Fremont National Forest = FREM). If the location

contains only a single station and is likely to remain that way, the location and station codes generally are the same.

General configuration: An idealized MAPS station is roughly square or circular in shape and encompasses an area of about 20 hectares (50 acres, about 450 meters on a side or circular with a 250-meter radius; Fig. 2). Ten 12-meter mist nets are distributed more or less uniformly but opportunistically (where birds will be caught) within a core area of about eight hectares (20 acres, about 280 meters on a side or circular with a 160-meter radius). The station includes everything within 100 meters of any net. If nets are separated by more than 200 meters, the area between the nets (at least a 25-meter-wide corridor) should also be considered as part of the station. Note that the nets are numbered in a roughly circular arrangement; this reduces the likelihood of net numbers being recorded incorrectly as the nets are checked consecutively. We realize that many MAPS stations will not be situated in a study area that permits a square or circular configuration. Some study areas may be quite irregular in shape and others, perhaps in riparian habitats, may be long and narrow. In these cases, nets should be established as uniformly and systematically as possible in order to cover the entire core area, maintaining the recommended net density (see below) but still allowing each net run to be completed in a reasonable time. Even in these cases, the station is considered to include everything within 100 m of any net.

Mist nets (number): The number of nets utilized at a station should be the maximum number (at the appropriate density) that can be operated safely and efficiently given the personnel available to run the station. Thus, only the number of nets that can be operated in a standardized manner over the long term should be established. In most instances, ten 12-meter nets might be the optimal number that can be operated by one person or two people. With a larger number of personnel or fewer birds, this number might be increased to 15 or even 20 nets; the size of the station should be increased accordingly so as to maintain the appropriate net density. With a smaller number of personnel and more birds, this number might need to be decreased to six or seven nets over a smaller area. The number and distribution of nets should be such that all the nets can be checked within 15-20 minutes if there are no birds to extract (i.e., an empty net run). We have set five as the minimum number of nets permissible at a station, since it is unlikely that really useful data can be obtained from a station with fewer than five nets.

Mist nets (density): The density of nets is an important variable with regard to the precision of the data that can be obtained from mark-recapture analyses. Net density will affect both the number of different individuals captured C thus the population size sampled C and the capture probabilities of those birds. Spreading the nets as widely as possible will tend to increase the number of territories intersected, and thus the population size sampled, but will tend to decrease the capture probability for the birds on any given territory. Moving the nets closer together will do the reverse. Thus, there must be some optimal intermediate density of nets that will maximize precision by optimizing simultaneously both the capture probability and the population size sampled. This optimal density may vary from species to species and from station to station depending upon average densities and territory sizes of the various species. Analyses of MAPS data indicate that stations that produce both high capture probabilities and high capture rates

operate with net densities of about one to two nets per hectare. We suggest that the optimal density of nets for most MAPS target species may be about 1.25 to 1.5 nets per hectare. Thus, ten nets could be placed effectively in a study area of about seven or eight hectares. With nets placed at this density, the distance between adjacent nets will average about 75-100 meters. In general, except in steep, rugged terrain, visiting ten nets placed 75-100 meters apart in 15-20 minutes should present no problem. The size of the netting area may need to be reduced (and the consequent net density increased) for stations established in steep, rugged terrain so that an empty net run can be completed within the allotted 15-20 minutes.

Mist nets (placement): Nets should be placed opportunistically at sites at which birds can be captured most efficiently, such as the brushy portions of wooded areas, forest breaks




Figure 2. Diagram of an idealized MAPS station.

or edges, and in the vicinity of water. The establishment of net sites at a station should strike a balance between the conflicting needs of capturing substantial numbers of breeding adults for estimating adult survival rates and substantial numbers of dispersing young and adults for indexing productivity. This may best be achieved by placing nets in both edge and non-edge portions of the study area. To optimize both the number of birds captured and their capture probabilities, nets should be placed relatively uniformly over the available habitat at each station. Because it is not permissible to move nets after the start of the second field season, care must be taken to select optimally-efficient, permanent net sites. Care also should be taken to ensure the safety of captured birds by not placing nets low over water or at sites subject to extreme wind or heat. Nets stacked two high or placed end-to-end in batteries are acceptable but not recommended, as they double the netting effort but generally don't double the number of captures. Although artificial food and water sources are not permissible within stations, they may exist adjacent to stations on property not under the control of the MAPS operator; remember that the station boundaries extend outward 100 meters from the net. Once the net sites are established, choose and flag a fixed net-run route that will minimize travel time to all nets, and number the nets sequentially along this route. Net designations should be numeric, unique within a station, and two characters long (e.g., 01, 02, 10). Remember that nets within batteries and stacked nets must be numbered individually.

Mist nets (size, type, and mesh size): We strongly recommend that all nets used in the MAPS Program be 12-meter, 30-mm mesh, four-tier, black, tethered, nylon mist nets. Other sizes, types, and meshes may be used if local conditions so warrant, but these variables should remain constant at each net site over all periods and years that the station is operated. One 12-meter net operated for one hour represents an effort of 1.0 net hour. Thus, if nets of other sizes are used, the effort reported must be adjusted accordingly. For example, a nine-meter net operated for one hour would be counted as 0.75 net hour.

Operating a MAPS station

Station registration: Before taking up station operations, a station registration form should be submitted for each station. The information on the form provides us with contact information for the station operator or operators. It also provides us with information on the station's geographic setting, critical for accessing remotely sensed data covering the station, and information on intended station operations. Once we receive a registration form for a station, the station operator or operators are added to the mailing list for the anticipated initial banding season. Refer to pages 26-30 for detailed instructions.

Banding - dates of operation: MAPS is strictly a breeding-season study. The breeding season, in general, is considered to extend from May to August and is divided into ten 10-day periods: (1) May 1-10; (2) May 11-20; (3) May 21-30; (4) May 31-June 9; (5) June 10-19; (6) June 20-29; (7) June 30-July 9; (8) July 10-19; (9) July 20-29; and (10) July 30-August 8. The strategy for the timing of operation is that each station should be operated for all ten-day periods beginning with the first period during which (a) the great




Figure 3. Recommended starting periods for MAPS stations. Appropriate periods for stations at high elevation stations may be later than indicated on the maps. Stations in habitats adjoining the Gulf of Mexico may start in Period 1.

majority of the breeding adults of the target species have established territories; and (b) individuals of these species migrating toward more northerly breeding grounds are no longer passing through the area. The start of operation will vary, therefore, from station to station depending on the timing of the breeding season at each station, which, in turn, is dependent primarily on latitude and altitude but also, to some extent, on longitude. Refer to Figure 3 for the recommended starting period at your location. Note that the starting period for stations at higher altitudes may have to be delayed by one (or, in years of exceptionally heavy and late-melting snowpack, even two) period(s) after the period indicated in Figure 3. In years in which late-melting snowpack do cause a delay in the initiation of breeding, the operation of periods subsequent to the starting period may also have to be delayed somewhat, more so earlier than later in the season. Nevertheless, it is extremely important that the number of periods during which the station is operated be held constant at each station for all years.

It is also important not to begin the operation of a MAPS station before migrating individuals of the locally-breeding species, bound for breeding areas farther north, have finished moving through the area. These birds, if captured, will bias estimates of the proportion of residents in the adult population as well as productivity indices. The

presence of such spring migrant individuals will also tend to lower the precision of all these estimates. Furthermore, the locally-breeding adults of any given species are usually the first individuals to arrive at a given location and, if captured before the start of the MAPS data-collection period, may learn to avoid the nets. Elimination from analysis of banding data collected before the official start of the season will tend to negatively bias adult survival rates, because most breeding adults are captured early in the season and net avoidance may prevent their recapture. Thus, in general, spring-migration monitoring should not be conducted at MAPS stations. If operators must run a spring-migration monitoring program and a MAPS program in the same place, they must use different net sites for the two programs.

Collection of MAPS data should be curtailed before substantial numbers of individuals (of the locally-breeding species) that are migrating south from breeding areas farther to the north begin to pass through the station. Inclusion of these individuals in productivity analyses will produce productivity indices that will tend to be more representative of areas farther north than of the local landscape. Analyses conducted on data from the four-year (1992-1995) MAPS pilot project indicated that substantial numbers of migrating individuals (as determined from the fat contents of the banded birds) began moving through most areas, regardless of latitude, after Period 10 (July 30 - August 8) (DeSante et al. 2004a). As a result, MAPS protocol now calls for the operation of all MAPS stations through Period 10 each year, but not thereafter.

In contrast to the situation in spring, however, it is not necessary to actually curtail the operation of the station late in the season before fall migrants begin passing through the area. Rather, the station can be operated after Period 10 and data from these later periods can be removed from analysis after the fact. This is because very few, if any, breeding adults are captured for the first time late in the season and the elimination of data from these periods will not affect survival-rate estimates. Thus, a station can use the same nets for fall-migration monitoring as for MAPS monitoring without compromising the value of the MAPS data. Indeed, analyses of age ratios from successively later periods during fall-migration monitoring will provide measures of productivity from areas increasingly farther north. It is very important to note, however, that if MAPS nets are operated outside of the breeding season, such operation must be discontinued at least three months prior to the beginning of the appropriate starting period for that station. This will assure that net avoidance by breeding individuals of permanent resident species will not bias survival-rate estimates for these species.

Effort data: Effort data are critical for comparing capture rates among years and for assessing both productivity indices and population trends. Because daily activity patterns differ both by age class and species, period-by-period, net-by-net, and hour-by-hour effort data are necessary for comparing productivity levels among years and for estimating numbers of birds missed because of missing effort. These data should be summarized on the Summary of Mist-Netting Effort form. Refer to pages 31-38 for detailed instructions regarding effort at MAPS stations.

Breeding Status data: The goal of the Breeding Status List is to provide a complete assessment of the summer residency status of all species present at each station each season. In order to accomplish this, it is necessary to record observations of the nesting

20 - 2025 MAPS Manual

behavior, singing, and overall presence of each species during each visit to the station. These data should be summarized on the Breeding Status List. Refer to pages 72-79 for detailed instructions.

Habitat Structure Assessment (HSA) data: The habitat structure assessment data serve three main functions: they provide a classification for each station, permit detection of gross changes in habitat structure at the station that may explain changes in population demographics, and provide station-specific habitat data to complement remotely-sensed landscape data at a fine resolution. HSA's should be conducted every five years, unless the habitat at your station has undergone a major change (e.g., fire, hurricane, logging, construction, brush-clearing, etc.). Refer to pages 80-88 for detailed instructions.

Instructions and data forms: Operators of registered stations will receive a beginning-of-season letter from The Institute for Bird Populations in early April each year. This will also direct new operators to download a copy of the MAPS Manual. All operators will be asked to download blank copies of the following data forms: Banding, Unbanded, and Recaptures banding-data sheets, the Standard Net Opening and Closing Times form (first year of operation only), the Summary of Mist-Netting Effort, the Summary of Mist-Netting Results, the Breeding Status List, and the forms associated with the Habitat Structure Assessment. They should also download a description of our collaboration with the Center for Tropical Research and the Bird Genoscape Project, which explains the optional feather-pulling protocol.

It is the operators' responsibility each year to make as many copies of the forms as they will need that year. Unused forms should be discarded at the end of the season because they may become obsolete the following year.

Recording data and making corrections: All data should be recorded in dark, water stable (i.e., not gel pen) ink or pencil. If you make corrections on any data sheet, use a good eraser (pencil) or fast-drying correction fluid or correction tape (pen). Do NOT just write over errors.

The use of MAPSNet

MAPNet is a cloud-based computer program for editing, verification, and error tracking of MAPS data. At the end of the 2024 MAPS season, it replaced the long used data verification program MAPSPROG. It offers contributors the opportunity to edit coding problems (codes that do not conform to IBP's preferred codes set forth in this manual); and address and correct, if applicable, within-record inconsistencies (conflicts between codes within a record, such as juvenile birds with breeding condition or after hatching year birds with no skull pneumatization) and between-record inconsistencies (conflicts in species, age, or sex determinations in different capture records for a given band number). The verification procedures encoded in the program reflect the MAPS data-collection guidelines described in this manual and ageing and sexing criteria presented in Pyle (2022) for the months covered by the MAPS season.

By providing a data editing/verification program to contributors, IBP hopes to decentralize the process of data correction, returning it to the control of those who collected the data

and who should, therefore, be better situated to make necessary corrections and adjustments. As a result, we hope that the quality of MAPS data overall will improve and that contributors, by verifying their own data, will be better able to identify areas in which they can improve their data-collection techniques in future seasons. MAPSNet integrates data entry modules for Banding, Effort, Breeding Status, and Habitat Structure data.

MAPS data-use policy

Data contributed to the MAPS program will be used by IBP to further the mission and goals of the MAPS program, as detailed in the Introduction section of the MAPS Manual. These goals include providing estimates and indices of demographic parameters, linking demographic parameters to population trends and environmental variables (e.g., habitat, climate), and providing information to land managers that will help create and maintain habitats to conserve and enhance North American landbird populations. Individual MAPS contributors will be acknowledged by name in any IBP publication or report arising from the use of MAPS data whenever their data represent more than 5% of the MAPS data used in that publication or report, and will be offered co-authorship whenever their data represent a substantial proportion (more than 30%) of the MAPS data used in that publication or report.

The MAPS database is the most extensive database on landbird demographics in North America and represents an invaluable scientific resource of immense conservation potential. As such, IBP is eager to share this resource with outside researchers. With the release of the MAPS Data Exploration tool (<https://ibp-maps-data-exploration-tool.org/app/maps>) in spring 2023, it has become easier for researchers to access this wealth of data.

As part of the access to the data in the application, researchers much agree to the data policy that if 20% or more of the MAPS data requested by a researcher will come from any single MAPS contributor, IBP will forward the written request to that contributor for permission to use those data. IBP also requests that publications or reports using MAPS data acknowledge the MAPS Program and IBP as the source of the MAPS data, and acknowledge individual MAPS contributors by name whenever their data represent more than 5% of the MAPS data used in that publication or report. We request that researchers offer co-authorship of any publication or report using MAPS data to individuals whose data represent a substantial proportion (more than 30%) of the entire data set used in that publication or report. Finally, **we request that a copy of the final publication or report be sent to The Institute for Bird Populations** (a PDF version is fine) so that it can be archived and included on The Institute for Bird Populations publications page: <http://www.birdpop.org/pages/pubsDatabase.php>.

An example of some language for an acknowledgement could be:

We thank the many dedicated volunteers who have collected and donated these data to the MAPS program. We also thank The Institute for Bird Populations for developing the MAPS Program and curating the MAPS data. Data used in this analysis were made available via funding from the National Science Foundation (Grant EF 1703048).

22 - 2025 MAPS Manual

Data downloads are tracked by IBP. And, on a regular basis, MAPS Operators will be sent a report of the data request summary from when their data was downloaded. This information can be used as justification as to supporters and funders who often ask how the data are being used.

By submitting your MAPS data, you are allowing your data to be included in analyses completed by IBP and non-IBP researchers who make data requests. Although we assume that our data-use policy will be followed and you will be acknowledged and offered co-authorship when appropriate, we cannot guarantee our policy will be followed once data are distributed to non-IBP researchers.

STATION REGISTRATION

Joining the MAPS Program

If you are interested in establishing one or more MAPS stations and feel that you are able to meet our requirements, please complete the registration form which is available on the IBP website at: <https://www.birdpop.org/pages/mapsDataForms.php> and submit it to the MAPS Coordinator, Danielle Kaschube at dkaschube@birdpop.org. On the registration form you will detail the proposed location, habitat, and operation of your station (Fig. 4).

Subject to approval of your proposal, you will be added to the roster of active MAPS operators and will receive the necessary forms and instructions, as well as annual reports on the results of the program and the program's newsletter, *MAPS Chat*. Please register each MAPS station with the Institute for Bird Populations before initiating operations. This helps us plan and budget accordingly and ensures that you receive program and protocol updates.

Please feel free to send questions Danielle at dkaschube@birdpop.org or call her at 609-892-0445.

Instructions for completing the MAPS Station Registration Form

Date: Record the date the form is completed.

Station Manager Contact Information:

Name: The name of the station manager, the person in charge of the MAPS station and responsible for seeing that changes in forms and protocol are communicated to all persons helping at the station. This will be IBP's official contact person to whom mailings and phone calls with data questions will be addressed and who will be acknowledged in publications and reports. Please keep us up-to-date concerning changes in contact information or responsibilities.

Title: The job title of the station manager within the organization, if any, with which the station is affiliated.

Affiliated Organization: The organization, if any, with which the station is affiliated.

Address and phone numbers: The mailing and e-mail addresses and phone numbers for the station manager.

Federal Banding Permit #: Provide the federal bird banding permit number under which the station will be operated. If you haven't yet received your federal permit, write "in process" in this space and provide the permit number once you have been approved.

Contact Information for an Additional Station Operator: You may provide contact information for another individual with station operation responsibilities on this form. Both operators will be included in our mailing lists and receive mailings concerning station operations. Often, secondary operators are staff biologists, technicians, students, or volunteers who play a critical role in conducting the banding station field work. If more than two individuals should be associated with this station, please provide the additional names and contact information via email at the time the registration form is submitted.

24 - 2025 MAPS Manual

Location Code: A unique, four-character code which you may select to designate your set of stations. If the code you propose conflicts with established MAPS location codes, we will contact you to discuss an alternative.

Station Code: A unique, four-character code which you may select to designate your station. For single-station locations, this is typically identical to the location code. If the code you propose conflicts with established MAPS station codes, we will contact you to discuss an alternative.

Name of Station: The full name of your station; please try to keep it short (four words or fewer).

Funding Source(s): List government agencies, non-governmental organizations, foundations, and grants providing financial support for station operation. Use "private" if the station is self-financed or if private individuals provide funding.

Property Name: The name of the piece of land on which the station is located. Please be precise in listing the property name e.g., Wenatchee National Forest; Kittitas County Environmental Educational Center; or Starr Ranch Sanctuary. If the property is owned by an individual or family, just write "private property".

Land Owner: The owner of the land on which the station is located. Please be precise in listing the land owner, i.e., for a station in Wenatchee National Forest, the "United States Forest Service, Wenatchee National Forest, Naches Ranger District"; for Kittitas County Environmental Education Center, "Kittitas County School District" rather than just Kittitas County; or for the Starr Ranch Sanctuary, "Audubon California" rather than just Audubon Society. If the land is owned by an individual or family, just write "private".

Nearest Town: Indicate the nearest reasonably sized community, as the neotropical migrant flies, on state-level road map or web application such as Google Earth.

County: This equates to parishes in LA, boroughs in AK, etc. Western provinces please indicate your regional or municipal district.

Latitude and Longitude: Please provide the lat/long coordinates in degrees, minutes, and seconds **to the nearest second** for the center of the station; please convert UTM coordinates and lat/longs given in decimals (many GPS units give seconds in decimals). North American longitudes are negative (except in the outer Aleutians).

Source of lat-long coordinates: The information source from which you determined the lat/long coordinates of the center of the station (e.g., Google Earth, hard copy of topographic map, online topographic map, GPS unit, etc.).

Datum: The reference point around which latitude and longitude are structured. If using a topographic map created before 1983 this will be NAD27. If using a topographic map created after 1983, a GPS unit, or online mapping information, the datum will be available somewhere on the source. (Google Earth uses NAD83.)

Average Altitude: The station's average altitude (elevation) in meters (1m=3.280833ft).

**2025 MAPS STATION REGISTRATION FORM**Date: February 20, 2025

Please refer to the instructions in the current MAPS Manual when completing this form.

Station Manager Contact InformationName: Joe Smith Title: BiologistAffiliated Organization: Bird Park County ParkAddress: Charles County Parks and Rec.456 Main St., Grayton, MD 20678Phone numbers: Work: 301-555-4444 Home or Cell: 301-556-2222E-mail: smith@svn.org Federal Banding Permit # 12345

You may also provide contact information for another individual with station operation responsibilities here. To add additional people, please submit a separate list of their contact information when the registration form is submitted:

Name: Sam Jones Title: _____Affiliated Organization: Maryland Ornithological SocietyAddress: 123 State St.Port Tobacco, MD 20677Phone numbers: Work: 301-654-3333 Home or Cell: 301-558-4141E-mail: jones@svn.org Federal Banding Permit # 12345 - CX**Station Information**

(pick up to 4 letters for each; the two codes may be the same)

Location Code: BIPA Station Code: SMCRName of Station: Small CreekFunding Source(s): Charles CountyProperty Name: Bird Park County ParkLand Owner: Charles County Parks and RecreationNearest Town: Grayton County: Charles State/Province: MD

26 - 2025 MAPS Manual

Please include **latitude and longitude** of the center of the station:

In degrees, minutes, seconds,

to the nearest second.

Latitude: 38E 26' 20"

Longitude: -077E 10' 41"

In decimal degrees,

to at least five decimal points.

Latitude: _____ . _____

Longitude: _____ . _____

- or -

Source of lat/long coordinates: eg. GPS, Google Earth, etc. Google Earth

Datum: NAD27 – or – WGS84/NAD83 (**check one**)

Average Altitude (in m): 11

General Habitat Description (e.g., "Mixed woodland in suburbia"; "cottonwood-willow riparian corridor"): mixed deciduous forest and riparian corridor

Please include a map showing the position of the station relative to nearby towns, major roads, and other geographic features; maps that are ideal for this purpose can be produced for free using Google Earth (free at www.earth.google.com).

Station Operation

First year of operation (expected): 2025

Number of nets: 10

Number of hours of operation per day (We recommend six): 6

Number of days of operation per ten-day period (We recommend one): 1

Periods of operation: From Period 3 through Period 10.

DATA-USE POLICY

Please refer to the MAPS Manual to read the full data-use policy.

In summary, if you submit data to the MAPS program your data will be made available for analysis by both IBP and other researchers.

JS Initial here to indicate awareness of this policy

To ensure receiving a timely spring packet of data sheets and information, **submit this form by April 1** to Danielle Kaschube, dkaschube@birdpop.org. **Forms submitted after April 1 are still valid** but may delay receiving the beginning-of-season materials.

Figure 4. Completed MAPS Station Registration Form.

Please include two maps: 1) a map on which the location of the station is clearly marked relative to nearby towns, roads, and other geographic features. 2) a map on which the exact location of the station is clearly marked relative to habitat features. Maps that are ideal for this purpose can be produced for free by using Google Earth (downloadable for free at www.earth.google.com. Having the “eye alt” [bottom left of Google screen] at about 100 kms usually provides a good scale for the first map and about 1-2 kms usually provides a good scale for the second map.)

First year of operation (expected): Please indicate the year in which you plan to begin operating your station.

General Habitat Description: Using key words, provide a brief description of the habitats at the station. Some examples: “Spruce-fir forest/meadow with willow thickets” or “Cottonwood riparian corridor/desert scrub” or “Maple-basswood forest with kettlehole marsh,” etc.

Number of nets: The optimal density appears to be about one net for every two acres in the 20-acre core of the study area. The minimum number of nets permitted is five (see “Establishing a MAPS station,” above). Note: a six-meter net = 0.5 net

Number of hours/day and days/period: In order to provide maximum comparability among stations, we strongly recommend six hours of operation per day (beginning at local sunrise) and only one day of operation per 10-day period. If you are operating in an extremely hot environment, five hours of operation per day may be more appropriate.

Periods of operation: See Figure 3 for appropriate starting period. All stations should run through Period 10. High-altitude stations may start one or even two periods later as appropriate. MAPS net sites should not be used in the spring before the appropriate starting period, since spring banding activity at MAPS net sites may induce net avoidance among resident birds for the remainder of the breeding season (see “Establishing a MAPS station,” above).

Data-use Policy: Initial to indicate that you have seen and agree to the IBP data-use policy.

MIST-NETTING EFFORT

Operation of nets

The importance of standardizing effort between periods and between years cannot be overemphasized. The accuracy and precision of MAPS indices and estimates depends on effort being equal, both in quantity and in timing, in all periods and all years. Thus, the number of nets operated and the timing of their operation should be standardized for all days of operation and kept constant from year to year at each station.

The first net should be opened at official local sunrise or at 0400, whichever is later; thus, starting time will change during the course of the season. The nets should be opened in the same sequence on each day of operation. If possible, they also should be

checked in this same sequence on every net run. They should remain open, if possible, for exactly six hours and should be closed in the sequence in which they were opened. At stations being operated in hot climates, it may be necessary to close nets earlier than six hours after opening and to open nets earlier in the morning, but no more than 30 minutes before sunrise. Effort for a given day may be shifted up to 30 minutes early or late (relative to the standard at the station) if circumstances demand it. Nets never should be opened more than 30 minutes before sunrise nor be closed any later than 30 minutes after the standard closing time for the station. Nets should not be operated if the average wind speed exceeds ten knots or gusts exceed 20 knots (the tiers of the net will be blown into concave 'C's) or if other weather variables (e.g., precipitation or extreme heat or cold) are likely to endanger the lives of captured birds. Efforts to lure or drive birds into nets are not permitted.

Frequency of mist-netting effort

MAPS nets should be operated on only one day during each ten-day period, and the dates of operation in consecutive ten-day periods should be fairly far apart (in general, at least six days). While it is true that increasing the number of days of operation in each ten-day period will tend to increase the resulting capture probabilities, the payoff from this increase seems to fall off rapidly after two or three days of operation. Although two days per ten-day period may offer the best return on capture probability per effort spent, the two days certainly will be spent better by operating two different stations for one day each and thereby effectively doubling the total number of birds handled. Thus, if the personnel at a given station have the ability to operate on multiple days in each ten-day period, we strongly recommend the operation of multiple stations for one day each, even if the stations must be adjacent to each other (Burton and DeSante 2004). Only for certain experimental stations and other already-established stations that have a long history of operating on more than one day per ten-day period will multiple days of operation per period be accepted. In these cases, the number of days of operation during each period should, if possible, be standardized for all periods; the total effort and timing, however, must be kept consistent from year to year at each station.

MAPS Periods and Intended Periods

The primary assumption of MAPS effort standardization is that banding effort at any time during a given period is equivalent to effort at any other time during the same period. Refer to pages 20-22, 'Banding C dates of operation' for clarifications of the MAPS Periods. Generally, banding effort in the middle of the period is likely to best approximate the proportion of young to adults prevailing for that period. However, any banding effort within the span of ten days that comprises that period is considered to be an appropriate estimator of that proportion.

Circumstances will occasionally make it impossible to conduct banding effort within the defined ten-day period. If it is impossible to put in the effort for a given period within the period itself, the station may be operated within five days before or after the period in

question. For example, an operator who knows she/he will not be able to band during Period 3 due to prior engagements, may band for Period 3 as early as May 16, but no earlier. Even though the date May 16 falls in the range of Period 2, the effort is intended for Period 3. As another example, suppose your station starts operation in Period 3, but due to thunderstorms during this ten-day block, you are unable to band. You will need to make-up this effort by running the station no later than June 4. In this case, even though the date June 4 falls in Period 4, the effort is intended for Period 3.

This effectively lengthens the MAPS season by five days on either end; thus, all data collected at MAPS nets from April 26 to August 13 should be submitted.

At higher altitudes in western mountains (generally above about 2,000 m), the entire season may need to be shifted up to ten, or even 20, days later if a heavy late-lingering snowpack delays the onset of breeding. This is a rare situation, but operators at such stations may decide to delay the entire operation of their station by one or even two periods in that year, thus completing station operation in Period 11 or 12. These operators are, in effect, banding during Period 4 (or Period 5) for Intended Period 3, etc. In this case, the season may extend as late as August 23, or even September 2 (allowing for the five-day grace period at the end of the period). If you are unable to put in the effort for Periods 9 or 10 within the allowable time frame, please try to make up the effort as soon as possible, but certainly well before the end of August. Periods 9 and 10 are critical periods for calculating productivity indices, and we may be able to use data from later in August to make up the missing effort.

Alternatively, MAPS operators may decide to delay operations by one or two periods at the beginning of the season, then gradually “catch up” to normal or near normal station operation, and complete late in Period 10 or in Period 11. This is the preferred method of shifting effort, as neotropical migrants in years of phenological delay tend to speed up the breeding season by shortening the time between broods or between nesting attempts in order to leave the breeding range shortly after they would have left in a normal breeding season. In such a situation, banding should extend no later than Period 11, and the number of days between banding sessions should be decreased, and remain relatively equal, over the course of the season. For example, for a station that normally begins operation in Period 3 (May 21- 30), but instead begins in Period 5 (June 10-19 - essentially 20 days late) and plans to continue banding through Period 11 (August 9-18 - essentially 10 days late), there are only seven periods - or 70 days - in which to conduct eight sessions of banding. Thus, banding should occur approximately every 8-9 (8.75) days, rather than every 10 days, over the course of the season, ending in Period 11.

Making up missed effort

If nets are closed early or opened late (relative to the standard at the station) due to inclement weather or unforeseen circumstances, the missing hours should be recovered if possible. This may entail either ending the day late (**but no more than 30 minutes**) as shown in Figure 6 for Intended Period 7, or making up the missed effort on another day within the same ten-day period as shown in Figure 6 for Intended Period 5. You **must** make up this effort if the missing effort amounts to more than half of a normal day's operation. For example, assume a normal full day's operation is 60 net hours (ten 12-

meter nets open for six hours). On one particular day, however, the nets are opened at 0600 but have to be closed at 0830 due to rain. Only 25 net hours have been accumulated. If the nets can be reopened at 0900, they should be closed at 1230 to recover the lost 30 minutes (5 net hours). If, however, it continues raining beyond 0900, the remaining 35 net hours should be made up as soon as possible during the appropriate hours (0830-1200) on another day within that period. If circumstances will prevent your return later in the intended period to make up the lost effort, then reopen the nets later in the morning, conditions permitting, to accumulate as much effort as possible (and at least half a normal day's operation) for that intended period (as shown for Intended Period 7 in Figure 6).

If multiple days are operated during an intended period, not designed to make-up missed effort, the second and subsequent days are assigned a **subperiod** code, B through J, as appropriate. If a single day is operated each period, the subperiods are all A.

Minimum allowable effort

For the purposes of MAPS analyses, we divide the MAPS season into two "superperiods," an adult superperiod, during which adults usually predominate in the catch, and a young superperiod, during which young often predominate. For a given location, the dates of these superperiods depend on the recommended starting period for that location (Fig. 3). Table 1 lists the adult and young superperiods for each starting

Table 1. Adult and young MAPS superperiods.

<u>Recommended start</u>	<u>Adult superperiod</u>	<u>Young superperiod</u>
Period 1 (May 1-10)	Periods 1-6 (May 1-Jun. 29)	Periods 7-10 (Jun. 30-Aug. 8)
Period 2 (May 11-20)	Periods 2-7 (May 11-July 9)	Periods 8-10 (July 10-Aug. 8)
Period 3 (May 21-30)	Periods 3-7 (May 21-July 9)	Periods 8-10 (July 10-Aug. 8)
Period 4 (May 31-Jun. 9)	Periods 4-8 (May 31-July 19)	Periods 9-10 (July 20-Aug. 8)
Period 5 (Jun. 10-19)	Periods 5-8 (Jun. 10-July 19)	Periods 9-10 (July 20-Aug. 8)

period. For data from a given station-year to be useable in MAPS analyses, the station . must have been operated for a minimum of three periods during the adult superperiod and a minimum of two periods during the young superperiod. Please note that for stations having a recommended start in Periods 4 or 5, absolutely no periods may be missed during the young superperiod (Periods 9 and 10). Please also keep in mind that all stations should be run from the recommended starting period through Period 10.

STANDARD NET OPENING AND CLOSING TIMES

In order to assist operators in timing the operation of their nets consistently from year to year, the Standard Net Opening and Closing Times sheet (Fig. 5) provides space for filling in local sunrise times and the net-opening and -closing schedule you intend to follow. This information is essential for analyses of effort comparability across years and for corrections for missed effort. In addition, should station operations be passed on to someone else, this information will provide the new operators and their volunteers with the times at which the nets are to be operated each period.

Typically, the standard opening time is local sunrise and the standard closing time is six hours later. Operation at some stations may deviate from this schedule consistently, year after year. In hot climates, for example, nets may need to be opened before sunrise. In cold climates, nets may need to be opened after sunrise, as shown in Periods 3 and 4 in Figure 5. In either case, the standard opening time should be no more than 30 minutes earlier or later than sunrise.

To make it easier to coordinate with banding assistants, the change from period to period in the standard opening times you designate may deviate from sunrise by 10-20 minutes, as demonstrated in Periods 8 and 10 in Figure 5. Fill out this form only once and submit a copy of the completed form to IBP with your data packet at the end of the season; please resubmit the form in future seasons only if you must change your standards.

Instructions for completing the Standard Net Opening and Closing Times form

Operators Name: Record the name of the MAPS Station Manager.

Location: Record your four-character location code.

Station: Record your four-character station code.

Date: Record the date the form is completed.

Sunrise: In the 'Period' field, record the sunrise time for your station for each period that you normally operate. Enter the sunrise time for the dates indicated in the period column (roughly, the mid-period dates), not the sunrise times for the dates you ran this season. U.S. sunrise tables are available free on the Web at https://aa.usno.navy.mil/data/RS_OneYear. Once on the web-site, you can enter your station location information. Remember that US longitudes are negative and you need to adjust for the appropriate time zone in the form.

Standard Open and Standard Close: Following the guidelines given above, record the standard opening and closing times for your nets in each period. Be sure to record the standard open and standard close times you **intend** to open and close your nets each period for your station every season; these times are not necessarily the times your nets were actually operated in any given season.



The INSTITUTE for
BIRD POPULATIONS

MAPS STANDARD NET OPENING AND CLOSING TIMES

Location: BIPA Station: SMCR

Date: 08/13/2025

Station Manager: Joe Smith

Period	Standard Open	Standard Close
1 Sunrise 05/05: _____		
2 Sunrise 05/15: _____		
3 Sunrise 05/25: <u>0550</u>	<i>0550</i>	<i>1150</i>
4 Sunrise 06/04: <u>0545</u>	<i>0550</i>	<i>1150</i>
5 Sunrise 06/14: <u>0544</u>	<i>0540</i>	<i>1140</i>
6 Sunrise 06/24: <u>0546</u>	<i>0550</i>	<i>1150</i>
7 Sunrise 07/04: <u>0550</u>	<i>0550</i>	<i>1150</i>
8 Sunrise 07/14: <u>0557</u>	<i>0600</i>	<i>1200</i>
9 Sunrise 07/24: <u>0604</u>	<i>0600</i>	<i>1200</i>
10 Sunrise 08/03: <u>0613</u>	<i>0610</i>	<i>1210</i>

Figure 5. Completed MAPS Standard Net Opening and Closing Times form.

SUMMARY OF EFFORT

The Summary of Mist-Netting Effort is the only information available that allows us to analyze between-year changes in mist-netting data in a constant-effort manner. It is critical that this form be completed exactly as indicated. Please review this section of the manual carefully before filling out the Summary of Mist-Netting Effort forms. The most common problems that we encounter are unspecified net number(s), unspecified timing (to the nearest ten minutes) of the opening and closing of the net(s), and net-hour calculations. Remember, the opening and closing times you should record are the times which you started the net runs to open or close the nets and should be recorded in the same format as capture times. Also, please be sure to double-check all net-hour calculations. All effort at MAPS nets from Intended Period 1 through Intended Period 10 [including August data for stations at higher altitudes that were delayed by one (or, in years of exceptionally heavy and late-melting snowpacks, even two) period(s)] must be recorded on the Summary of Mist-Netting Effort forms (Fig. 6), including any effort conducted before the recommended starting period. Banding data submitted for Intended Periods 11 (August 9-18) and 12 (August 19-28) must also be accompanied by effort data.

MAPSNet checks ensure that all MAPS season banding records occurred on days the nets were actually operated.

Instructions for completing the Summary of Mist-Netting Effort form

Location: Record your four-character location code.

Station: Record your four-character station code.

List net numbers of all 12-m nets: Record the net designations of all 12 meter nets.

List net numbers and lengths of all other nets: Record the net designations and lengths of all other nets. If you do not operate other length nets, please indicate by recording "N/A" or "none."

If any nets are stacked, list their net numbers and how stacked: For example, nets 02 and 03 stacked: 02-low and 03-high.

Describe net changes from last year: Indicate any previously operated nets that were not operated in the current year and any new nets added. Please note that any moved nets will require new net numbers.

MAPS season shift due to heavy snowpack at high elevation stations: See page 31 for a complete explanation of when this rare season shift may be necessary.

Intended Period: Record the **intended** period for the date operated. Remember, if it is impossible to put in the effort for a given period within the period itself, it may be done within five days before or after that period. If the date operated falls outside the

standard ten-day period, include a note explaining why the operation did not occur in the standard ten-day period.

Date: Record the month and day of the date of operation.

Net Numbers: Record the net designations of the nets operated (not the quantity of nets opened). A single day's effort should be recorded on multiple lines if nets of different sizes are used or if the nets are open for varying periods of time. For example, if all ten nets were opened at 0600 and nets 08 and 09 were closed at 1000 due to sun or wind while the remaining eight were closed as planned at 1200, then the effort should be recorded on at least two lines. See also the example for Intended Period 3 in Figure 6.

Open Time and Close Time: These times should be recorded in the same format as capture times. That is, using the 24-hour clock, record, to the nearest 10 minutes, the opening and closing times of the first net opened or closed. Always enter three digits. Note that the ultimate zero is preprinted on the form; e.g., 6:24 a.m. = 062(0), 1:48 p.m. = 135(0).

Net Hours: Record the net hours accumulated (to the nearest 0.01 net hour) for the nets recorded on each line. For MAPS purposes, a 12m net is 1 unit length. So, a 12m net operated for 1 hours equals one net hour.

Period Net Hours: Record the total effort for all days in an intended period on the last line for the intended period.

Note No.: Record a note (with a note number) on the reverse (page 2) side of the form indicating why nets were opened or closed at times that deviate from the standard protocol. Record the note numbers for these notes in the Note No. column on the form.

Total net hours for all MAPS periods combined: Carefully sum the Period Net Hours for all MAPS periods operated during the year and enter the sum in the appropriate box on the reverse side (page 2) of the form. Please use a calculator to sum these Period Net Hours and please double-check your sum. A comparison of this sum to the sum obtained from the computerized effort file (in both IBP's verification procedures and MAPSNet) provides an important check that all effort data have been entered.



2025 MAPS SUMMARY OF MIST-NETTING EFFORT - pg 1

LOCATION CODE: B I P A STATION CODE: UPED

List net numbers of all 12-m nets: 01- 09

List net numbers and lengths (in meters) of all other nets: Net 10 = 9 m

If any nets are stacked, list their net numbers and how stacked (i.e., nets 02 and 03 stacked: 02-low, 03-high, etc.): No stacked nets

Describe any changes in your nets or net sites since last year. **Any moved nets require new net numbers!** No changes

MAPS PERIODS

- | | |
|--------------------------------|---------------------------------|
| Period One: May 01 - May 10 | Period Six: June 20 - June 29 |
| Period Two: May 11 - May 20 | Period Seven: June 30 - July 09 |
| Period Three: May 21 - May 30 | Period Eight: July 10 - July 19 |
| Period Four: May 31 - June 09 | Period Nine: July 20 - July 29 |
| Period Five: June 10 - June 19 | Period Ten: July 30 - August 08 |

NOTE: Heavy snowpack in some years can cause the breeding season to be delayed in the higher elevations of the western mountains. We allow the entire MAPS season to be shifted later by one or even two periods in these instances. If your station experienced a delay in breeding this year due to heavy snowpack, please indicate the number of periods you shifted this year (1 or 2): ____

Intended Period	Date (mm/dd)	Net number(s)	Open Time (nearest 10 minute)		Close Time (nearest 10 minute)		Net Hours	Period Net Hours	Note No.*
			hour	minute	hour	minute			
<i>e.g., 3</i>	<i>05/21</i>	<i>01 - 08, 10</i>	<i>06</i>	<i>0 0</i>	<i>12</i>	<i>0 0</i>	<i>54.00</i>		
<	<	<i>08</i>	<i>06</i>	<i>3 0</i>	<i>12</i>	<i>0 0</i>	<i>5.50</i>	<i>59.50</i>	<i>1</i>
<i>3</i>	<i>05/28</i>	<i>01-09</i>	<i>05</i>	<i>5 0</i>	<i>11</i>	<i>5 0</i>	<i>54.00</i>		
<	<	<i>10</i>	<i>06</i>	<i>2 0</i>	<i>11</i>	<i>5 0</i>	<i>4.13</i>	<i>58.13</i>	<i>1</i>
<i>4</i>	<i>06/05</i>	<i>01-09</i>	<i>05</i>	<i>5 0</i>	<i>07</i>	<i>5 0</i>	<i>18.00</i>		<i>2</i>
<	<	<i>10</i>	<i>05</i>	<i>5 0</i>	<i>07</i>	<i>5 0</i>	<i>1.50</i>		<i>2</i>
<i>4</i>	<i>06/07</i>	<i>01-09</i>	<i>07</i>	<i>5 0</i>	<i>11</i>	<i>5 0</i>	<i>36.00</i>		<i>2</i>
<	<	<i>10</i>	<i>07</i>	<i>5 0</i>	<i>11</i>	<i>5 0</i>	<i>3.00</i>	<i>58.50</i>	<i>2</i>
<i>5</i>	<i>06/14</i>	<i>01-09</i>	<i>05</i>	<i>4 0</i>	<i>11</i>	<i>4 0</i>	<i>54.00</i>		
<	<	<i>10</i>	<i>05</i>	<i>4 0</i>	<i>11</i>	<i>4 0</i>	<i>4.50</i>	<i>58.50</i>	
<i>6</i>	<i>06/24</i>	<i>01-09</i>	<i>05</i>	<i>5 0</i>	<i>07</i>	<i>3 0</i>	<i>15.00</i>		<i>3</i>
<	<	<i>10</i>	<i>05</i>	<i>5 0</i>	<i>07</i>	<i>3 0</i>	<i>1.25</i>		<i>3</i>
<	<	<i>01-09</i>	<i>08</i>	<i>5 0</i>	<i>12</i>	<i>2 0</i>	<i>31.50</i>		<i>3</i>
<	<	<i>10</i>	<i>08</i>	<i>5 0</i>	<i>12</i>	<i>2 0</i>	<i>2.63</i>	<i>50.38</i>	<i>3</i>

* Please write note on reverse side of this page.

Figure 6. Completed MAPS Summary of Mist-Netting Effort (page 1).

COLLECTION AND RECORDING OF BANDING DATA

All birds captured throughout the season, including recaptures, must be identified to species and must be aged and sexed if possible (use “unknown” if necessary). Age and sex birds by the extent of skull pneumatization and/or other appropriate plumage, breeding, mensural, or molt characters (Bird Banding Offices 1991, Pyle 2022). Incorrectly-identified, -aged, and/or -sexed birds are detrimental to analyses. All birds not already banded and not excluded from the operator’s banding permit (except hummingbirds, which are not included in MAPS analyses because most banders don’t band them) must be banded with a numbered aluminum band issued by your country’s banding office.

We consider Pyle (2022) as the authority for in-hand age, sex, and difficult species determinations of North American passerines and near-passerines. The information in this book is accepted by the banding offices as well. MAPS banders are expected to be using this book in the field, and the validity of your determinations will be checked using it as well. Please ensure that you understand the concepts presented in Pyle (2022) and apply them appropriately. Please also ensure to correct your copy of Pyle (2022) with the latest version of the errata (<https://www.slatecreekpress.com//pages/errata.php>).

General procedures for recording banding data

Primary MAPS data: Primary MAPS data are the data upon which all analyses of productivity indices, survival-rate estimates, and population trends are based. Thus, it is crucial that complete primary MAPS data be taken on all birds captured, including recaptures. Primary MAPS data include the following data fields (see pages 43-69): capture code (e.g., newly banded, recaptured, band changed), band number, species (as given by the species alpha code), age, how aged (if age determined), sex, how sexed (if sex determined), status (as required on banding schedules submitted to the banding office), date, capture time, station, net number, disposition, and feather pull. It is important to note that the primary MAPS data fields are the only ones that may be subject to modification based on other information obtained during the capture or by comparisons with other capture records of the same band number. Date, capture time, and net number will allow us to screen out records that cannot be used for multi-year comparisons. Ageing and sexing criteria will allow us to screen out improperly-aged or -sexed birds and to evaluate the reliability of these criteria.

Supplemental data: MAPS operators are also asked to collect supplemental data on all birds captured, including recaptures: extent of skull pneumatization, breeding condition (presence or absence of a cloacal protuberance or brood patch), extent of body and flight-feather molt, extent of primary-feather wear, extent of juvenile plumage, existence of molt limits and information on feather generation for selected feather tracts or groups of feather tracts, wing chord, body mass, and fat class. These data are used in verification programs to assure the accuracy of the species, age, and sex determinations. They can also provide invaluable information regarding spatial (geographic) and temporal variation in the timing and extent of breeding and molt and

the physiological condition of the bird. Because supplemental data reflect information taken directly from the bird in the hand, supplemental data must never be modified and should mirror precisely what was recorded in the field.

Optional data: Additional data, such as exposed culmen, tail length and/or tarsus length, may also be taken but are not required, although they sometimes are useful in verifying the primary data. If taking these measurements, please insure you know the correct procedure to take them as described in Pyle (2022)

Codes, scales, and forms: All data should be taken according to the standardized guidelines and utilizing the standardized codes described in these instructions and should be recorded on copies of the standardized 8 ½" by 14" MAPS banding-data sheets (Figs. 7 and 8).

We realize that some contributors to the MAPS Program have long been recording many of these data according to slightly different codes and scales. The codes suggested in these guidelines are the result of thousands of hours of field work and subsequent analysis by researchers at the U.S.D.A. Forest Service, the U.S. Fish and Wildlife Service, the Point Blue Conservation Science (previously Point Reyes Bird Observatory), and The Institute for Bird Populations and are summarized in Ralph *et al.* 1993. In an effort to aid in the standardization of the capture and banding data now being collected in North America, we suggest that you adopt the scales and codes presented here. If you find it impossible to adopt these scales and codes, you must provide us with an explanation of how your codes correspond to MAPS codes so that they can be converted to MAPS codes. Note that codes other than MAPS codes are not accepted by the MAPS entry/verification program, MAPNet.

Please use the MAPS banding-data sheets for recording all MAPS banding data. We have examined well over 70 different banding-data sheets and have designed these sheets to increase the ease, logic, and accuracy of recording banding data in the field, as well as the efficiency and accuracy of entering those data into a computer for analysis and the production of banding schedules.

There are three types of MAPS banding-data sheets: the **MAPS Banding Sheet** for recording the use of new bands; the **MAPS Recaptures Sheet** for recording recaptures; and the **MAPS Unbanded Sheet** for recording birds that are captured but left unbanded.

Multiple-station locations: If more than one station is operated at a location, band strings are shared among stations, and banding is not conducted simultaneously at multiple stations, then the data from these stations should be combined on a single set of banding-data sheets to avoid gaps in the band sequences on the forms. If more than one set of banding-data sheets must be used, please use a different page-numbering sequence for each set (A1, A2,.; B1, B2...) so that each location/year/band-size/page combination is unique.

Non-MAPS data: Banding data from non-MAPS sites (e.g., nestlings, traps, feeder stations) or collected before or after the MAPS season (April 26-August 13, or through September 2 if effort for periods 9 and 10 is being made up) can be submitted to the MAPS Program if it makes clerical/administrative work easier for your banding

operation, but is not required. We recommend non-MAPS data should be recorded on separate forms and, if possible, with separate band strings so as to avoid breaks in the band sequences on the MAPS Banding Sheets. However, occasionally, individual records that technically cannot be considered MAPS data are included on MAPS banding-data sheets. These records might include birds found dead on a trail or they might consist of a bird captured accidentally not in a MAPS net or at your MAPS station. These records must be identified as non-MAPS in order to avoid including them in constant-effort analyses. **Mark these records by recording "NM" in the NOTE NUMBER field.**

Page headings and other notations: Be sure to fill out the headings - **Location, Year, Band Size (for new bands), and Page #** - on each banding-data sheet. Use the four-character location code determined during station registration. Number the pages sequentially for each band size, starting with page 1 every year; in other words, there will be a page 1 for each band size used each year. This is very important, as it will allow us (and you) to see at a glance that all data are submitted. Please write "End of year" at the bottom of the last page of each band size each season.

New bands: To ensure that band numbers are recorded and computerized correctly and to facilitate band inventory and scheduling, it is of the utmost importance that original banding data for only a SINGLE STRING OF BANDS be included on any single MAPS Banding Sheet and that the bands be recorded (and, as much as possible, used) in sequence. Please write "End of string" below the last record for each band string.

Lost and destroyed bands: Lost and destroyed bands should be recorded in sequence on the MAPS Banding Sheets. Record only code, band number, species name as "Band Lost" or "Band Destroyed," date, and station.

Recaptures: Every capture of a banded bird is a "recapture". Recaptures thus include returns (first captures in the current year of birds banded previously in the same place on the same permit), repeats (subsequent captures, even on the same day, of birds banded or recaptured in the same place earlier in the current year), and foreign recaptures (first captures of birds banded in a different place or on a different permit). Birds banded outside of MAPS operation and recaptured during MAPS operation are considered recaptures. Previously-banded birds that escape or are inadvertently released before the band number is read should also be recorded as recaptures. Complete data should be taken for **all** recaptures and should be recorded only on MAPS Recaptures Sheets. It is crucial that new and recapture banding data NOT be entered on the same sheets. Do NOT separate recaptures by band size. **Recaptures of birds with non-USGS bands from another country**, i.e., not from the USA or Canada, are handled differently than recaptures of birds with federal USGS bands. Record the band number from the foreign country on the recapture page and then add a USGS aluminum band to the other leg. Record this new band on the newly banded page for the appropriate band size. Leaving the original band on the bird will allow the other country to keep tracking the history of the bird and adding the new band will allow tracking of the bird in the USGS databases and, if the bird is recaptured in the country of origin, allow those banders to know that it was captured in the US or Canada as well. Cross reference the

band numbers in the note fields on the back of each page and both records should be given capture code "A."

Changed bands: If a band is replaced, record the capture on both the MAPS Banding Sheet (new band) and MAPS Recaptures Sheet (recapture record). Record the old band number on the Recapture Sheet, with the new number as a note on the back. Record the new band number on the Banding Sheet, with the old number as a note on the back. The old band should be sent to the banding office with the schedule on which the new band is reported. Both records should be given capture code "C." NEVER re-use a band you have taken off a bird; it makes tracking individuals exceedingly difficult and, because the structural integrity of the band is compromised, increases the risk of injury to the bird. Importantly, a changed band should be counted only as a single recapture on the Summary of Mist-netting Results (see below), because it involves only one bird.

Added bands: Occasionally, birds wind up with a band on each leg. Usually, this is the result of a bander not realizing that the bird is a recapture and applying a band to the other leg or due to a foreign recapture of a bird banded in a country other than the United States or Canada (see recapture section above). Accidental double banding can be avoided by ensuring that all banders at your location are banding on the same leg. If both bands are readable and neither is endangering the bird's welfare, it is best, because of the risk of injury to the bird, not to attempt to remove one of the bands. If the bird was captured with two bands, enter a record for each band, both with code "A" (for "Added Band"), on the Recapture Sheet. If you have applied the second band, record it (again as code "A") on the Banding Sheet, with the original band number in a note, and record the original band on the Recapture Sheet (also with code "A") with the added band number in a note (analogous to changing a band, except that no band was removed). As with changed bands, added bands should be counted only as single recaptures on the Summary of Mist-netting Results (see below).

Unbanded birds: As much information as possible (including all primary MAPS data) must be recorded on the MAPS Unbanded Sheets for all birds that are captured but not banded (escapes, releases, and mortalities) regardless of the size of band they would have received had they been banded. Data on these birds is essential for calculating capture rates, and mortalities can be used in productivity analyses. A bird is considered an "escape" if it was touched prior to escape; a bird that bounces out of or escapes from a net before it is touched should not be recorded. "Releases" might include species that a bander is not authorized to band (gallinaceous species or hummingbirds) and birds for which the recommended band size is unavailable. See Table 2 for species alpha codes for gallinaceous birds.

Mortalities: Even if all reasonable precautions are taken, mortalities do occur occasionally in the course of mist netting. If a bird dies before it is banded, it should be recorded on the MAPS Unbanded Sheet. If a bird dies just after it is banded, remove and destroy the band. Record the bird's data on the Unbanded Sheet to account for the capture and the band number on the Banding Sheet as destroyed (code "D") to account for the destroyed band. In either case, the bird should receive "000" in the "STATUS" field and a "D" or "P" in the "DISP" field for "death due to cause other than predation" or "predator-caused mortality," respectively. If the mortality is a recapture, record the

individual's band number and all normally collected data on the Recapture Sheet and then remove the band. Destroy the band unless it is a recovery, in which case you should submit the information electronically at <http://www.reportband.gov>. As before, enter "000" in the STATUS field and "D" or "P" in the DISP field.

Banding-data fields

The front of the banding-data sheet is broken into 36 fields each containing one or more columns. Each of these fields is described separately below. Please write out completely the first record on each sheet each day. After that, use a "greater than" (>) or "less than" (<) in the BANDER'S INITIALS, SPECIES NAME, STATUS, DATE, CAPTURE TIME, and STATION fields if the entry is repeated (on the same day only) on the next line; do not use ditto marks or vertical lines that can be mistaken for '1's and do not use these symbols in any other fields. If data for a given field are not collected, leave the field blank; do not use zeroes, nines, hyphens, slashes, or any other symbols to designate data not taken.

Please record all data taken, even if the values are "0," and do not make assumptions. For example, if you have what you believe is a female with a brood patch, please verify that there is no cloacal protuberance. Once you have done so, enter "0" in the CP field; if you leave it blank, we cannot assume that you checked to make sure there was not a CP.

BANDER'S INITIALS – Place the initials of the bander or person taking the data in this field. Write the initials and full names of all the banders on the page in the spaces provided at the bottom of the form.

CODE – Capture Code. Use the codes shown at the top of the banding-data sheet. Use "N" for all newly-banded birds; "L" for lost bands; "D" for destroyed bands; "U" for unbanded birds; "C" for changed bands (refer to the section on changed bands [above] for instructions on code- "C" captures); "A" for added bands (refer to the section on added bands [above]); and "R" for all other recaptures, regardless of whether they are repeats, returns, or recoveries. Note that the only capture codes acceptable on a given sheet are those presented at the top of the sheet (e.g., "N" does not appear on the Recapture Sheet). The code for unbanded birds, "U," is already filled in on the Unbanded Sheet for these birds.

BAND NUMBER – For new, lost, and destroyed bands, enter the complete band number for the first band on the first line of each page. Do not use a hyphen to separate the prefix from the rest of the band number. Please double-check to be sure that this first band number is entirely correct. Thereafter, enter only the last three digits right-justified for all other band numbers on the page. For all recaptures, however, be sure to enter the full band number each time. Furthermore, please double-check the band numbers on all recaptured birds before releasing them. Incorrect band numbers on recaptures are the most serious errors of all because correct band numbers on recaptured birds are the basis for all mark-recapture analyses. We strongly recommend the use of some form of optical magnification, preferably a magnification visor, to read the band numbers of

recaptured birds and to examine skull pneumatization of all birds. The best one we have found is the OptiVISOR, an optical glass binocular magnifier that fits over your head, tilts up when not in use, and leaves both hands free to band and examine the bird. We recommend the DA-5 model (2.5 power at a focal length of 8"; price about \$30). It is distributed by lapidary and jewelers'-supply houses or on Amazon.

It is extremely important that **all band numbers be nine characters long**. Three-digit prefixes must be recorded prefaced with a "0" (e.g., 972 becomes 0972). Two-digit prefixes must be recorded prefaced and followed by "0" (e.g., 81 becomes 0810). For unbanded birds, leave BAND NUMBER blank.

SPECIES NAME – Enter at least an abbreviation of the species name (e.g., "Blk-cap Chick" for Black-capped Chickadee). This abbreviation will not be entered in the MAPS database but will serve as a check against the error-prone SPECIES ALPHA CODE (below), such as Barn Swallow ("BARS") and Bank Swallow ("BANS"), both of which are often written incorrectly as "BASW". Write "Band Lost" or "Band Destroyed" in this space where appropriate.

SPECIES ALPHA CODE – Enter the four-letter code for the species (e.g., "BCCH" for Black-capped Chickadee) from *Four-letter and six-letter alpha codes for birds recorded from the American Ornithologists' Union check-list area* (Pyle and DeSante 2003). This list (further updated in Pyle and DeSante [up through 2023]) can be downloaded from The Institute for Bird Populations' website at <https://www.birdpop.org/pages/birdSpeciesCodes.php>. Very few discrepancies that are likely to be encountered by banders exist between this list and the current BBL codes. In the very few cases where the species codes differ, and until the BBL is able to update their codes, MAPSNet will convert the alpha codes provided to current BBL codes when producing the export file for Band Manager. Species codes for gallinaceous birds are given in Table 2; these species do not fall under the jurisdiction of the federal banding offices, which consequently do not provide alpha codes for them.

Table 2. Species alpha codes for gallinaceous birds.

<u>Species</u>	<u>Alpha codes</u>	<u>Species</u>	<u>Alpha Codes</u>
Plain Chachalaca	PLCH	Dusky Grouse	DUGR
Chukar	CHUK	Sooty Grouse	SOGR
Himalayan Snowcock	HISN	Sharp-tailed Grouse	STGR
Gray Partridge	GRAP	Greater Prairie-Chicken	GRPC
Ring-necked Pheasant	RNEP	Lesser Prairie-Chicken	LEPC
Common Peafowl	CPEA	Wild Turkey	WITU
Ruffed Grouse	RUGR	Mountain Quail	MOUQ
Greater Sage-Grouse	GRSG	Scaled Quail	SCQU
Gunnison Sage-Grouse	GUSG	California Quail	CAQU
Spruce Grouse	SPGR	Gamble's Quail	GAQU
Willow Ptarmigan	WIPT	Northern Bobwhite	NOBO
Rock Ptarmigan	ROPT	Montezuma Quail	MONQ
White-tailed Ptarmigan	WTPT		

42 - 2025 MAPS Manual

Occasionally, notes associated with a record indicate that the species determination for a recapture or an unbanded bird was uncertain. **Mark these records by recording "QS" in the NOTE NUMBER field.**

AGE -- Enter a single-digit numeric code for the age class of the bird, as shown at the top of the banding-data sheets. Alternate, single-character, alpha codes for each age class also are presented below. We strongly recommend using the numeric codes, however, because of the difficulty we have experienced in distinguishing between a printed 'H' and a printed 'A', which together comprise the vast majority of the age classes. Note: MAPSNet will only accept the numeric codes. These codes are:

- 4 - Local (L): A young bird incapable of sustained flight. (These birds always should be processed as quickly as possible and released very close to the net in which they were captured.)
- 2 - Hatching Year (H): A bird capable of sustained flight and known to have hatched during the calendar year in which it is captured. If these birds are in full juvenile plumage, consider releasing them near the net in which they were captured. Parents may still be feeding these young.
- 1 - After Hatching Year (A): A bird known to have hatched before the calendar year in which it is captured; year of hatching otherwise unknown.
- 5 - Second Year (S): A bird known to have hatched in the calendar year preceding the year in which it is captured (known to be in its second calendar year of life).
- 6 - After Second Year (O): A bird known to have hatched earlier than the calendar year preceding the year in which it is captured (known to be at least in its third calendar year); year of hatching otherwise unknown.
- 7 - Third Year (T): A bird known to have hatched two calendar years prior to the year in which it is captured (known to be in its third calendar year).
- 8 - After Third Year (Z): A bird known to have hatched more than two calendar years prior to the year in which it is captured (known to be at least in its fourth calendar year); year of hatching otherwise unknown.

Some species, for example woodpeckers, can occasionally be aged to fourth year (4Y) or after-fourth year (A4Y). There are no numeric codes for these ages, so the age code will need to remain ATY (8) which is less precise, but still correct. Include a note in the NOTES field of the more specific age.

- 0 - Indeterminable (U): Age unknown because age indeterminable; i.e., age determination attempted but not possible with confidence.
- 9 - Not attempted (X): Age unknown because age determination not attempted.

Please attempt (without relying on previous capture data) to age adult birds as second year (SY) or after second year (ASY). It should be possible to reach this level of precision with at least some individuals of roughly 95% of North American passerine and near-passerine species. In addition, many near-passerines (including woodpeckers) and a few passerines may be aged to third year (TY) and after third year (ATY). Our

Page # 1

MAPS BANDING SHEET

THE INSTITUTE FOR BIRD POPULATIONS
www.birdpop.org

Location **B I P A**

Year **2025**

Band Size **0**

Year **2025**

Page # **1**

CODE		AGE		HOW AGED AND HOW SEXED		WING CODE		SKILL		CL. PROT.		BR. PATCH		FAT		BODY MLT.		FF MOLT.		MOLT LIMITS & PLUMAGE		DISP.	
USE	SEX	REACH	HATCH	YEAR	SEX	WING	WING	WING															
Band Destroyed	Band Lost	Band Changed*	Band Adhered*	Band Adhered*	Band Adhered*	Band Adhered*	Band Adhered*	Band Adhered*	Band Adhered*	Band Adhered*	Band Adhered*	Band Adhered*	Band Adhered*	Band Adhered*	Band Adhered*	Band Adhered*	Band Adhered*	Band Adhered*	Band Adhered*	Band Adhered*	Band Adhered*	Band Adhered*	
C	C	1	9	3	0	3	8	8															
P	B	N																					
C	C	N																					
A	B	N																					
C	C	N																					
A	B	N																					
A	B	N																					
P	B	N																					
P	B	N																					
C	C	N																					
A	B	N																					
A	B	N																					
P	B	N																					
END OF STRING																							

BANDERS INITIALS

DATE

STATION

NET

DISP. NOTE

FTHR. PULL NUMBER

Bander: C C Cirris Countersong P B Peter Bridgestone A B Andrea Basset/hound

Figure 7. Completed MAPS banding-data sheet for Band Size '0' (front). Associated notes are shown in Figure 8.

NOTE NO.	NOTE
1	band changed, original band number = 122045678
2	molt limit within lesser and med covs
3	aged by adult plumage; escaped before fully processed
4	Unable to fly on release but couldn't recapture
5	couldn't see molt limit betw/ tertts and middle ss (but Pyle says can be hard to see)
6	pox on legs: pp 1-2 growing
7	male seen near net during extraction. Rects looked ~truncate but pp covs looked ~tapered - unsure of SY/ASY, aged AHY
8	molting pp 3-4. unreplaced outer pp covs & rects truncate
9	gr covs w/ 3 feather gens - 3 alternate coverts; rects with limited yellow (like Fig. 263 B)
10	molting pp 2-3, ss 8-9, & pp covs, L & R 1, unsure of brood patch 5
11	preformative nearly complete; head & upper breast feathers black w/ yellow edging

Figure 8. Note section for example banding page.

ability to index juvenile survival rates and estimate recruitment rates of young and immigration rates of adults hinges on your ability to discriminate between SY and ASY age classes. Since the presence of juvenile or first-alternate feathers indicates SY, whereas the lack of such feathers often is not definitive, it is likely that more SYs than ASYs will be identified.

WRP – Wolfe-Ryder-Pyle age classification. **This age classification system underwent a revision in 2022.** We recommend reading the paper detailing the revisions by Pyle et al. 2021 (full reference in the Literature Cited) and the 2022 and 2023 MAPS Chats (<https://www.birdpop.org/pages/mapsChatArchive.php>) which provides a summary of the paper as well as examples of how to use the system for MAPS data.

Enter the age code for the cycle-based age class of the bird. This code describes the plumage and molt stage of the individual. Entry of the cycle based codes is currently optional but we hope operators will quickly adopt their usage.

The first of the three core characters indicates the cycle:

- F - first cycle
- S - second cycle (usually used only for woodpeckers)
- T - third cycle (used rarely for woodpeckers)
- D - definitive cycle

The second of the three core characters indicates the molt status:

- C - mid-cycle, no molting feathers on bird
- P - bird in molt. For HYs, or any adult bird undergoing a pre-alternate molt, the individual can be in body-feather molt and the body molt should be more than just a few feathers, i.e., $BM \geq 2$; for AHYs during pre-basic molts, the molt generally, but not necessarily, should include primary feather molt. This code can also be used for a bird not actively molting if it is in the suspended phase of a suspended molt, e.g., REVI, but this situation will be rare.
- U - unknown

The third of the three core characters indicates the plumage. I.e., the plumage a non-molting individual is currently in or the plumage in which a molting individual is molting into:

- J - juvenile plumage
- F - formative plumage
- A - alternate plumage.
- B - basic plumage
- X - auxiliary formative
- U - unknown plumage

46 - 2025 MAPS Manual

An adjunct code – placed before the three-character core code – is used to indicate when the core code cannot be specifically defined, either because of how the species molts or because the plumage of an individual bird isn't certain.

- M - indicates that the bird has reached, at minimum, the plumaged indicated by the core WRP code
- H or A - specific to WRP code FCF. Used to separate HYs that have finished the molt (H-FCF) from SYs that have not begun the second prebasic molt yet (A-FCF). These codes will be needed for certain species in July and August in which HYs may complete the preformative molt before SYs have started the second prebasic molt; thus, we need to distinguish these FCFs with an adjunct code.

Not all of the code combinations are valid. Below is a list of the codes most frequently used in MAPS, followed by the calendar based age code that usually applies in North America, but note that there is no direct correspondence between WRP and calendar-based age codes. The most common codes are listed first and in the order they are encountered in a typical passerine:

- FPJ - **F**irst **p**rejuvenile molt: The individual is molting into its juvenile plumage and must have molting body and/or flight feathers. AGE=2/HY in North America and during the MAPS season.
- FCJ - **F**irst **c**ycle juvenile plumage: The individual is in full juvenile plumage, i.e., only juvenile feathers are present and the bird has no molting feathers. JP=3 and AGE=2/HY during the MAPS season. Only J codes should be in the molt limits and plumage fields and the JUV. BDY PL. should equal 3.
- FPF - **F**irst **p**reformative molt: The individual is molting into its formative plumage. The bird must have molting body and/or flight feathers. JP=1 or 2 and AGE=2/HY during the MAPS season.
- FCF - **F**irst **c**ycle **f**ormative plumage: The individual is in full formative plumage, i.e., the bird has no molting feathers and most often has a mixture of retained juvenile feathers and replaced formative feathers. **During the MAPS season, we use the WRP adjunct codes; please enter an adjunct code here for all known FCFs - either H or A.**

H-FCF - Hatch year, first cycle formative plumage. This code is applied to fresh HYs that have finished the preformative molt for species in which adult birds might be in the same plumage, e.g., Wilson's Warbler, Canada Warbler, etc. AGE=2/HY during the MAPS season.

A-FCF – After hatch year, first cycle formative plumage. This code is applied to worn SYs that have not begun the second prebasic molt, for species in which hatch year birds might be in the same plumage, e.g., Wilson's Warbler, Canada Warbler, etc. AGE=5/SY during the MAPS season.

- M-FCF - **Minimum first cycle formative:** This code will be used for species that have a complete preformative molt, e.g., WREN, BUSH, NOCA, GRSP, HOSP, swallows, and other species –**and– for unknown plumage adults of other species.**
AGE=1/AHY during the MAPS season.
- FPA - **First prealternate molt:** The individual is molting into its first alternate plumage. The bird must have molting body and/or flight feathers. AGE=5/SY during the MAPS season.
- FCA - **First cycle alternate plumage:** The individual is in full first-alternate plumage, i.e., the bird has no molting feathers and has a mixture of juvenile, formative, and alternate feathers. AGE=5/SY during the MAPS season.
- M-FCA - **Minimum first cycle alternate:** The individual is in alternate plumage but it can't be determined if it is in first or definitive cycle alternate plumage.
AGE=1/AHY during the MAPS season.
- SPB - **Second prebasic molt:** The individual is molting into its second basic plumage. In most cases, it should have molting primaries or secondaries and still retain some juvenile, formative and/or first alternate feathers. AGE=5/SY during the MAPS season.
- M-SPB - **Minimum second prebasic molt:** The individual is molting into its basic plumage. In most cases, it should have molting primaries or secondaries and no definitively juvenile feathers remain so it is uncertain if this is second or subsequent prebasic molt. AGE=1/AHY during the MAPS season.
- DCB - **Definitive cycle basic plumage:** The individual is in full basic plumage, i.e., the bird has no molting feathers and has only basic feathers. Often AGE=6/ASY during the MAPS season.
- DPA - **Definitive prealternate molt:** The individual is molting into its definitive alternate plumage. The bird must have molting body and/or flight feathers. AGE=6/ASY during the MAPS season.
- DCA - **Definitive cycle alternate plumage:** The individual is in full alternate plumage, i.e., the bird has no molting feathers and has a mixture of basic and alternate feathers. AGE=6/ASY during the MAPS season.
- DPB - **Definitive prebasic molt:** The individual is molting from one basic plumage to definitive basic plumage. In most cases, it should have molting primaries or secondaries and show both retained and replaced definitive basic and/or alternate feathers. Often AGE=6/ASY during the MAPS season.
- FCU - **First cycle unknown plumage:** The individual is its first cycle because it retains juvenile feathers, but it is unknown if it is in formative or alternate plumage. Used often for species where the prealternate molt can vary from absent to limited and/or difference between formative and alternate plumages is subtle, e.g., OVEN or female COYE, etc. AGE=5/SY during the MAPS season.

Less common codes:

- SCB - **Second cycle basic** plumage: The individual is in second basic plumage with a few retained juvenile feathers and has no molting feathers. This and the following two codes are used almost exclusively for woodpeckers during MAPS banding. AGE=7/TY during the MAPS season.
- TPB - **Third prebasic** molt: The individual is molting into its third basic plumage. It must have molting primaries and retains at least some juvenile and second basic feathers. AGE=7/TY during the MAPS season.
- TCB - **Third cycle basic** plumage: The individual is in third basic plumage with retained juvenile and second basic feathers and has no molting feathers. This code is only rarely used for woodpeckers. AGE=4Y during the MAPS season.
- 4PB - **Forth prebasic** molt: The individual is molting into its fourth basic plumage. It must have molting primaries and have at least four generations of feathers (including the newly molted ones). AGE=8/ATY during the MAPS season.

There are times when the codes UPB, UPU, and UUU are needed because not enough information was seen on the bird, often because the bird was released early. However, whenever possible we ask you try to use one of the above codes to both provide IBP with the most information possible on each bird and to solidify your understanding of molts and plumages. Please contact IBP if you have any questions!

The WRP determination must be supported by data in the body molt, flight feather molt, juvenile body plumage, and/or the first seven MOLT LIMITS & PLUMAGE fields.

HOW AGED -- The how-aged codes indicate the criteria that you used to determine the age of the bird. Use only the appropriate code(s) shown at the top of the banding-data sheets. Use two codes if possible and enter them from left to right in order of importance for your age determination. If you use only one code, enter it left-justified. You must record at least one criterion unless the age is unknown (i.e., unless AGE = 0 or 9). The how-aged codes are as follows:

- S - Skull: The degree of skull pneumatization.
- C - Cloacal Protuberance: The presence of a cloacal protuberance on adults.
- B - Brood Patch: The presence of a brood patch on adults.
- J - Juvenile Plumage: The presence of juvenile body plumage on juveniles. To use this code, the JUV. BDY PL. box needs to have a number greater than zero.
- L - Molt Limit: The presence of two generations of feathers **within** a feather tract (e.g., within the greater coverts) **or between** two adjacent feather tracts (e.g., between the primary coverts and greater coverts). If the Molt Limit code is used, at least one of the first seven MOLT LIMITS & PLUMAGE fields (see below) must be filled in.

- P - Plumage: The appearance, if reliable for ageing, of plumages other than juvenile body plumage. Feather color, shape, quality, and wear are plumage characteristics; measurements are not. Contrasts in color, shape, quality, or wear between two generations of feathers or groups of feathers should generally be treated as a molt limit characteristic (L), not a plumage characteristic (P). If the Plumage code is used, at least one of the first seven MOLT LIMITS & PLUMAGE fields (see below) must be filled in.
- M - Molt: The presence and characteristics, if reliable, of **active molt**, indicated by pinfeathers and/or missing flight feathers in a symmetric pattern.
- F - Feather Wear: The degree, if reliable, of flight-feather wear.
- I - Mouth/Bill: The external and/or internal appearance, if reliable, of the bill or the presence of a fleshy gape on very young birds.
- E - Eye color: The color of the iris, if reliable. This does not include the eye ring.
- V - Age updated after review of photographs of the individual
- O - Other: Any criterion not listed above (e.g., date, orbital apterium, talon-flange serration, tail fork, egg visible, etc.). If you use this code, you must explain how the bird was aged in a note on the back of the sheet.

Note that W (Wing Length) and T (Tail Length) are not valid how-aged codes. If tail length (or tail fork) are used to age Barn Swallows or Great-tailed Grackles use "O" and provide a note to that effect.

Please do not age recaptures based upon previous captures. Each capture should be treated in the field as if it were a new bird in order to avoid perpetuating previous errors and to enable us to see what is possible from field observations at that time of year.

Please remember that you must record at least one ageing criterion unless the age is unknown (indeterminable or unattempted). In many cases, especially with adults, more than one criterion is available; RECORD TWO! Try to look at and indicate features such as plumage, eye color, and bill/mouth that are not recorded elsewhere on the form. Don't forget that CPs and BPs can be used for ageing adults, since HY birds don't get them. Please study the sample banding sheet (Fig. 7) to better understand how this field should be used.

You must record data in at least one of the first seven MOLT LIMITS & PLUMAGE fields whenever "L" or "P" is used as a how-aged code. You also must record data in at least one of the MOLT LIMITS & PLUMAGE fields whenever you age an adult bird more specifically than AHY (i.e., SY, ASY, TY, or ATY). Remember, however, that when you age an adult bird as SY by Molt Limit, you must also indicate, with an additional how-aged code, what you used to determine that it was **not** a HY bird (e.g., skull, cloacal protuberance, brood patch, plumage, molt, feather wear, mouth/bill, eye color, or other). This additional how-aged code should generally be recorded as the left-justified one.

SEX -- Enter "M" for male, "F" for female, "U" for indeterminable (sex unknown because sex indeterminable, i.e., sex determination attempted but not possible with certainty), or

“X” for not attempted (sex unknown because sex determination not attempted). If you must use numeric codes (we strongly recommend against them), use “4” for male, “5” for female, “0” for indeterminable, and “9” for not attempted.





HOW SEXED -- Use the codes below as in HOW AGED above. As with age, do not sex recaptures in the field based on previous captures. Note that S (skull), L (molt limit), M (molt), and F (feather wear) are not valid how-sexed codes.

- C - Cloacal Protuberance: The presence of a cloacal protuberance, if reliable, on adult males.
- B - Brood Patch: The presence or degree of a brood patch, if reliable, on adult females.
- J - Juvenile Plumage: The appearance of juvenile body plumage, if reliable, on juveniles.
- P - Plumage: The appearance, if reliable, of all plumages other than juvenile plumage. Does not include measurements.
- I - Mouth/Bill: The appearance, if reliable, of the bill.
- E - Eye Color: The color, if reliable, of the iris.
- W - Wing Length: The wing chord, if reliable.
- T - Tail Length: The length, if reliable, of the tail.
- O - Other: Any criterion not listed above (e.g., singing, tail fork, egg visible, etc.). Use of this code requires an explanatory note.


We commonly see data for species that cannot be sexed by plumage with “CP” or “BP” in the HOW SEXED field. This invariably is the result of a bander instructing a recorder to enter “male by CP” or “female by BP” and the recorder not realizing that “P” in this case stands for plumage. It is much safer (and faster) to say the codes (e.g., “M by C”) rather than the words.

SKULL -- Skull Pneumatization. In order to determine the degree of skull pneumatization, it is necessary to part the feathers of the head to get them out of the way (wetting them slightly may help), then gently rock the skin back and forth over the skull while looking through the skin to the skull. The best procedure is to start at the back of the skull and proceed toward the front looking for the pattern of the line that separates the pneumatized area from the area that is not pneumatized. A pneumatized skull consists of two layers of bone connected by tiny “struts” and filled with air, much like the wing of a plane. A pneumatized skull appears opaque and grayish with tiny whitish dots. In contrast, an un-pneumatized skull, consisting of a single, thin layer of bone, appears pinkish and somewhat translucent and never shows the minute dots characteristic of a pneumatized skull. We very strongly recommend the use of a binocular magnifier such as the OptiVISOR for determining the degree of skull pneumatization (see above under BAND NUMBER). See Yunick 1979, Ralph *et al.* 1993, and Pyle 2022 for more complete information (including diagrams) on the determination of age by skull pneumatization.


Skull pneumatization should be recorded by means of the scale shown below. We strongly recommend using the numeric codes, although corresponding alpha codes that were developed by Ralph *et al.* (1993) also are shown below; these alpha codes may be used in the field if necessary, but we request that they be converted to numeric codes prior to submission.

- 0 - (N = none): Skull not pneumatized; that is, only a single thin layer of bone covers the entire brain, which shows through the thin covering of bone and appears as an unmarked, pinkish color. Beware of thick-skinned species such as corvids and parids, whose skull can be very difficult to see because the skin itself tends to be rather opaque; and heavily-muscled species such as grosbeaks and cardinals, whose jaw muscles can obscure the rear of the skull. 
- 1 - (T = trace): A trace of skull pneumatization can be seen at the very back of the skull, usually appearing as an opaque, grayish crescent or a very-small, triangular area. Somewhere from 1 to 5% of the skull is pneumatized. 
- 2 - (L = less than 1/3): Skull less than 1/3 pneumatized but some pneumatization is obvious. Thus, somewhere from 6 to 33% of the skull is pneumatized. Generally, the posterior part of the cranium has an inverted 'u'- or 'v'-shaped area of pneumatization that is usually distinctly grayish and contrasts with the unpneumatized area. The grayish area typically shows the characteristic, small, whitish dots of a pneumatized skull. 
- 3 - (H = half): Skull greater than 1/3 but less than 2/3 pneumatized. In typical birds, most of the rear half of the skull is pneumatized, as is a small portion of the front part extending back around the eyes. This front part of the skull is usually very difficult to see because the feathers of the forehead are dense and short and difficult to move out of the way. In most cases, a bird given a "3" skull will show a pneumatized area extending up the midline or sides of the skull. 
- 4 - (G = greater than 2/3): Skull at least 2/3 pneumatized but at least small areas of skull not pneumatized. Thus, somewhere from 67 to 94% of the skull is pneumatized. The un-pneumatized areas generally show either as two oval, pinkish spots on either side of the cranium or (rarely) as a single spot in the center of the skull. 

- 5 - (A = almost complete): Somewhere from 95 to 99% of the skull is pneumatized. These birds have virtually a fully-pneumatized skull that shows one or two tiny, dull-pinkish areas where the pneumatization is incomplete. It should be noted that some birds, including many flycatchers, thrushes, and vireos, never develop a fully pneumatized skull, even when adult, but retain a "5" skull throughout life. Thus, a "5"-skull bird cannot necessarily be called a HY/SY bird because it could be an AHY/ASY bird whose skull never completely pneumatized.




- 6 - (F = fully complete): Skull fully pneumatized.



- 8 - (I = invisible): Skull examined but extent of pneumatization not visible. Do not use this code if you have determined that pneumatization is incomplete but are unsure of the appropriate score; in this case, make your best guess!

CL. PROT. – Cloacal Protuberance. As the breeding season approaches, the cloaca of most male birds (and female Wrentits) begins to enlarge and forms an obvious protuberance which serves a role in sperm storage. The development of the cloacal protuberance is recorded according to the system shown below. Again, we strongly recommend using the numeric codes, although corresponding alpha codes are also given.

- 0 - (N= none): Cloaca not enlarged.
 1 - (S= small): Cloaca somewhat enlarged and noticeably swollen. The shape of the protuberance is generally such that it is widest at the base and narrowest near the tip (conical). Since small cloacal protuberances (CPs) can be hard to discern, caution should be used in ageing or sexing birds on the basis of a CP of 1 alone. A CP of 1 can **not** be used to age or sex the thrushes of the genera *Catharus*, *Hylocichla*, *Turdus*, or *Ixoreus*. We have found that unenlarged



cloacas in these species have often been designated CP = 1, which regularly has led to incorrect age or sex determinations.


- 2 - (M = medium): Cloacal protuberance large, with a diameter fully as large near the tip as at the base (cylindrical).
- 3 - (L = large): Cloacal protuberance very large and with a diameter considerably larger in the middle than at the base (bulbous).

Unlike a brood patch (see below), a regressing CP simply goes back down the scale: 3-2-1-0. CPs vary greatly in size and shape among species, being largest and most prominent in sparrows and thrushes and much less prominent in jays and Wrentits. It may be possible to sex species that rarely show prominent CPs by examining the angle of the CP with respect to the body axis. In males, the CP seems to point straight out, more or less perpendicular to the body axis. In females, the cloaca seems to point toward the rear of the bird, somewhat more parallel to the axis of the body. This same tendency can be used with caution to distinguish the occasionally slightly enlarged cloacal region of a female from a true class- 1 CP of a male. Class- 2 and - 3 CPs of males, of course, cannot be confused with those of females in any species except, possibly, in Wrentits.

Please note that all cloacas, whether enlarged or not, stick out. A true CP is characterized by firmness and lateral swelling. Note also that immature birds DO NOT get CPs.

BR. PATCH -- Brood Patch. Just prior to and during the time that the female (and in some species, the male as well) is incubating eggs in a nest, the feathers of the lower breast and abdomen are lost, vascularization increases just below the skin, and considerable fluid collects below the skin. The purpose of these changes is, of course, to facilitate the transfer of heat from the incubating bird's body to the eggs. The scale shown below should be used to record the sequence of events in the development and regression of a brood patch. Again, we strongly recommend using the numeric codes, although corresponding alpha codes are also given.

NOTE: In hummingbirds and in juveniles of most species, the lower breast and abdomen are normally unfeathered. This can cause it to look like a brood patch of 1 or 4, but the area is darker red and unwrinkled and usually has a less distinct margin.



- 0 - (N = none): No brood patch is present. The lower breast and abdomen are more-or-less feathered. Unfeathered areas of the breast and abdomen are smooth, without evident vascularization.
- 1 - (S = smooth): The lower breast and abdomen feathers are dropped and some vascularization can be seen, but most of the area is still rather smooth and dark red.
- 2 - (V = vascularized): Vascularization is evident, some wrinkles are present, and some fluid is present under the skin, giving the area a pale, opaque, pinkish color as opposed to the normal, dark-red muscle color.
- 3 - (H = heavy): The vascularization is extreme, the brood patch becomes thickly wrinkled, and much fluid is present under the skin, often giving the region a yellow blistery appearance. This is the maximum extent of the brood patch and corresponds closely to the time during which the bird is incubating eggs.
- 4 - (W = wrinkled): The vascularization mostly has disappeared and the fluid under the skin is mostly gone. The skin, however, retains many thin, dry-looking, contracted wrinkles.
- 5 - (M = molting): The vascularization and fluid and most of the wrinkles are gone. New pinfeathers are present as the area begins to become re-feathered. Most birds do not reach class 5 BPs until the nesting season is over and the prebasic molt has begun.

The sequence of 0 to 5 is rather symmetric. Classes 1 and 5 resemble each other, class 5 being distinguished most easily by the growth of new feathers. Similarly, classes 2 and 4 resemble each other but class 4 can be distinguished by its dry, thin wrinkles, as opposed to the thick, fluid-filled wrinkles of class 2.

FAT – Fat Content. Subcutaneous fat is a yellow or orange substance that is stored just under the skin and is used as fuel for migratory flights and for maintenance during the colder winter months. Fat generally is stored in three discrete areas that usually begin

filling in the following order: (1) the hollow in the furculum (wishbone) just below the throat at the top of the breast muscles; (2) the hollow directly under the wing, essentially in the "wingpit"; and (3) the lower abdomen just anterior to the vent area. The stored fat can be seen clearly through the nearly-transparent skin and contrasts with the dull, dark-reddish color of the breast muscles. It is seen most easily by holding the bird on its back while placing the index and middle fingers on the front and back of the bird's neck, stretching the head slightly forward along a line parallel to the body, and gently blowing the feathers away from the upper breast to expose the furculum. Then check under the wing and on the abdomen, again by blowing the feathers gently out of the way. Fat content generally can be assessed quite easily while checking for breeding condition and body molt. The placement of the field on the banding-data sheets reflects this fact. The codes shown below should be used to record fat content. Again, the use of the numeric codes is strongly recommended.

- 0 - (N = none): No fat in the furculum or anywhere on the body.
- 1 - (T = trace): A very small amount of fat in the furcular hollow (< 5% filled) but not enough to cover the bottom of the furculum, and no fat or just a trace of fat is present under the wing, on the abdomen, or anywhere else on the body; **or**, if there is no fat in the furcular hollow, at least a trace of fat is present under the wing, on the abdomen, or both.
- 2 - (L = light): The bottom of the furculum is completely covered but the furcular hollow is less than 1/3 filled, and a small amount of fat may be present under the wing, on the abdomen, or both; **or**, if there is no fat in the furcular hollow, a covering pad of fat is definitely present under the wingpit and, usually, on the abdomen.
- 3 - (H = half): The furcular hollow is about half full (actually anywhere from 1/3 to 2/3 filled), and a covering pad of fat is definitely present under the wingpit and, usually, on the abdomen; **or**, if there is no fat in the furcular hollow, a thick layer of fat occurs under the wing and on the abdomen.
- 4 - (F = filled): The furcular hollow is full (actually anywhere from 2/3 full to level with the clavicles) and a thick layer of fat also occurs under the wing and on the abdomen; **or**, if the fat in the furcular hollow is not full, the fat under the wing as well as on the abdomen is well mounded.
- 5 - (B = bulging): The furcular hollow is more than full; that is, the fat is bulging slightly above the furculum. The fat under the wing as well as that on the abdomen is also well mounded.
- 6 - (G = greatly bulging): Fat is bulging greatly above the furculum. Large mounds of fat occur under the wings and on the abdomen.
- 7 - (V = very excessive): The fat pads of the furculum, "wingpit," and abdomen are bulging to such an extent that they join. Nearly the entire ventral surface of the body is thus covered with fat, and fat even extends onto the neck and head. Such birds are nicknamed "butterballs."

NOTE: The upper fat classes (5-7) are seen most often just prior to and during migration.

BODY MLT – Body Molt. Body molt should be determined by examining the bases of all the contour feathers on the bird's body, including all the body feathers as well as the upper- and underwing coverts and the upper- and undertail coverts. The bases of the feathers can be exposed by blowing lightly but continuously over the body. The presence of pinfeathers is a sure sign of the early stages of molt. Later stages can be recognized by a remnant, scaly sheath at the base of each growing feather. These sheaths persist until the feathers are fully grown. You should integrate several factors in making your rating, including the number of feather tracts in molt and the proportion of feathers in molt in each feather tract. Body molt should be rated according to the scale shown below. Again, numeric codes are preferred.

- 0 - (N = none): No body molt. No feathers in sheath or growing.
- 1 - (T = trace): Only a very few feathers molting anywhere on the bird's body, usually in no discernible pattern.
- 2 - (L = light): A few feathers are molting from a few feather tracts, or some feathers (fewer than ½) are molting from only one tract. In general, fewer than 1/3 of the contour feathers on the bird are molting.
- 3 - (M = medium): Some feathers (generally fewer than ½) are molting from most tracts, or many feathers (generally more than ½) are molting from one tract or a few tracts. In general, from 1/3 to 2/3 of a bird's contour feathers are in molt. This class also should be used for a bird in spring whose pre-alternate molt normally includes only the head but that has nearly all head feathers in molt. Such a bird would be given a class "3" even though fewer than a of all its contour feathers are molting.
- 4 - (H = heavy): Many feathers (generally more than 1/2) are molting from many or most tracts. In general, more than 2/3 of the contour feathers on the bird are in molt.

FF MOLT– Flight-feather Molt. "Flight feathers" is a collective term for primaries, secondaries, and rectrices but when of **recording flight feather molt on the MAPS banding data sheet, only consider the primaries and secondaries.**

Most adult passerines in North America undergo a complete molt following the breeding season. This molt usually occurs from July to September and most often occurs on the breeding grounds, although there are some notable exceptions (see Pyle 2022). We refer to this complete molt in adults as the "**prebasic molt**" (= "adult prebasic molt" in Pyle 2022). At the same time of year (July to September), juvenile birds also undergo a molt which we refer to as the "**preformative molt**". In contrast to the complete prebasic molt of adults, the preformative molt in juveniles of most passerine species is "partial"; that is, it includes the body feathers but not the flight feathers, except sometimes the innermost rectrices (the "decks") and the innermost secondaries (the "tertials"). Thus, the presence or absence of symmetric flight-feather replacement in a bird undergoing

molt in the late summer and early fall often provides another good indicator of the age of the bird. First, be sure to check Pyle (2022) to make sure that the species does not replace flight-feathers during the preformative molt. Then, examine the primaries and secondaries for the presence or absence of flight-feather molt, and examine both the left and right sides to be sure that the replacement is symmetric and not adventitious (the accidental, generally asymmetric, loss of flight feathers or body feathers anywhere on a bird). Record flight-feather molt with the codes shown below. In this case, we recommend using alpha codes since the codes are categorical and do not represent a sequence that can be expressed numerically.

- N - (0 = none): No flight-feather molt.
- A - (1 = adventitious): Accidental, adventitious, usually asymmetric flight-feather molt.
- S - (2 = symmetric): Normal, symmetric or nearly-symmetric flight-feather molt, indicative of prebasic molt in adult birds and preformative molt in some young birds. A few species also exhibit prealternate flight-feather molt (see Pyle 2022).
- J - (3 = juvenile growth): Growth of juvenile flight feathers in fledgling birds (only to be used for very young birds, just out of the nest, growing their first flight feathers).

IMPORTANT NOTE: If a bird is exhibiting flight-feather molt, record, as a note, the particular group(s) of feathers (primaries or secondaries) in which molt is occurring. If possible, record which feathers are molting, e.g., PP1-4new, PP56 & S12 molting both sides. This information will aid greatly in the verification of age data.

FF WEAR – Flight-feather Wear. The juvenile generation of flight feathers (primaries, secondaries, and rectrices) is structurally weaker than later (adult) generations of feathers and thus may wear and fade at a faster rate. Furthermore, because young birds grow their juvenile flight feathers considerably before adults molt their flight feathers, at any given time during the following 12 months, juvenile flight feathers are older than the new generation of adult flight feathers. The result is that juvenile feathers in the following spring are likely to show greater degrees of feather wear than do adult feathers at that time. If so, these data may help to determine the age of birds first captured in the spring. In addition, hatching year birds in the summer have very fresh, new flight feathers, while adult flight feathers, before they are molted, are very old and worn. This also helps facilitate the determination of age in mid-summer birds. Examine only the outer 4-5 primaries to determine wear. Flight-feather wear should be classified according to the scale shown below. Again, numeric codes are preferred.

- | | | |
|---|--|---|
| 0 - (N = none): No wear at all. The feather edges are perfect. A light-colored edge exists all the way around the feathers, including the tips. |  | 0 |
| 1 - (S = slight): Very little wear. Feather edges are only slightly worn and no actual fraying or nicks have occurred. Often, a light-colored edge exists around the sides of the feathers but not at the tips. |  | 1 |
| 2 - (L = light): Relatively light wear. The feathers are definitely worn but with very little fraying and very few actual nicks. |  | 2 |
| 3 - (M = moderate): The feathers show considerable wear and some very definite fraying. Nicks and chips are obvious along the vanes. |  | 3 |
| 4 - (H = heavy): The feathers are very heavily worn and frayed. The tips are often worn completely off. |  | 4 |
| 5 - (X = excessive): The feathers are extremely ragged and torn up, and the shafts are usually exposed well beyond the vanes. All the tips are usually completely worn or broken off. |  | 5 |

JUV. BDY PL. -- Extent of Juvenile Body Plumage. Most fledgling birds wear a juvenile plumage that is distinct, at least in texture, from any other plumage of the species. Juvenile plumage is generally distinguished from adult plumages by loosely-textured ("fluffy") contour feathers, often with streaks or spots not found on corresponding adult feathers. It is important to examine individual feathers in assessing the extent of juvenile plumage. This plumage may be worn from only a few days to several months, depending on species and fledging date, until the **preformative molt**, at which time it is molted into "**formative plumage**". The extent of juvenile body plumage on a young bird, therefore, is often a good indicator of how long the individual has been out of its nest. Note that flight feathers (primaries, secondaries, and rectrices) are generally not replaced during the preformative molt and should not be considered when assessing the extent of juvenile plumage. The extent of juvenile body plumage should be recorded according to the scale shown below. Again, we recommend using the numeric codes rather than the alpha codes. **For the purpose of this field, only juvenile birds have juvenile body plumage!** Most birds do retain some juvenile wing coverts through the first breeding season, but these second-year birds are considered to have no juvenile body plumage because they have completed the preformative (and in some cases prealternate) body molt.

60 - 2025 MAPS Manual

- 3 - (F = full): Full juvenile body plumage. The bird has not yet begun its preformative molt.
- 2 - (G = greater): More than half of the juvenile body plumage remains, although the bird already has begun its preformative molt. The individual looks mostly like a juvenile, but some formative body plumage is evident.
- 1 - (L = less): Less than half of the juvenile body plumage remains. The bird has molted primarily into formative body plumage, but some juvenile body plumage remains.
- 0 - (N = none): No juvenile body plumage. The individual has molted already into full formative body plumage. All adult birds, including SYs, therefore, have "0" juvenile plumage, even if they have some retained juvenile coverts or flight feathers.

In summary, a bird is in full (3) juvenile plumage from fledging until the onset of the preformative molt. During this molt, juvenile plumage is replaced by formative body plumage. Thus, birds in partial (2 or 1) juvenile plumage must be in molt. Recently-fledged birds still may be growing their juvenile feathers but should be classed as "3" juvenile plumage. Similarly, birds in the final stages of the preformative molt may have shed all of their juvenile body feathers but still be growing their formative feathers; such birds have "0" juvenile plumage.

MOLT LIMITS & PLUMAGE - These fields are to be used for adult birds aged more specifically than AHY (i.e., SY, ASY, TY, or ATY), as well as for any birds (including those aged HY or AHY) aged by molt limit or plumage (i.e., any time "L" or "P" is used as a how-aged code). Up to eight fields, which describe individual (or multiple) feather tracts or non-feathered body parts, may be considered for any individual bird. At least one of the first seven fields must be filled in if the bird is aged by molt limit or plumage, and at least one of the fields must be filled in if the bird is aged SY, ASY, TY, or ATY. Refer to Pyle (2022), Froehlich (2003), and Saracco (2004) for additional discussion and examples of the use of molt limits and plumage criteria for ageing landbirds. The eight MOLT LIMITS & PLUMAGE fields are:

PRI. COVS - Primary coverts.

SEC. COVS - Secondary coverts (i.e., greater, median, lesser, carpal, and alula coverts and alula).

PRIMARIES - Primaries.

SECONDS - Secondaries, not including the tertials.

TERTIALS - Tertials.

RECTRICES - Rectrices.

BODY PLUM. - Includes all feather tracts of the head, upperparts and underparts (including the underwing coverts).

NON-FEATH - Includes all non-feather parts including bill, mouth, eye, legs, and feet. Typically, because the change in these features is gradual, rather than a

relatively swift change like molting feathers, the only two codes generally used in this field are J (characteristic suggests this is a juvenile bird) and N (characteristic suggests this is not a juvenile bird).

The codes entered in these fields should reflect the feather generation(s) present within the particular feather tract (or multiple feather tracts in the case of body plumage). Adventitiously (accidentally) replaced feathers should be ignored (except to provide context to the other feathers in the tract) because recognizing them as a separate feather generation will lead to mis-aging birds. Similarly, brand new or actively molting feathers should be ignored when coding tracts containing actively molting feathers.

The use of any of the following three codes during the MAPS season on an adult bird prior to completion of its prebasic molt indicates that it is a SY bird; the use of any of these codes on a young bird after its preformative molt confirms that it is a HY bird.

- J - Juvenile: Feather tract comprised entirely of retained juvenile (or a mix of juvenile and alternate) feathers, but no formative feathers. This code should also be used for NON-FEATH if non-feathered body parts show characteristics indicative of a young bird.
- L - Molt limit: Molt limit between juvenile and formative feathers exists **within** the feather tract, regardless of whether or not alternate feathers are also present in the tract.
- F - Formative: Feather tract comprised entirely of formative (or a mix of formative and alternate) feathers, but no juvenile feathers.

The use of any of the following three codes during the MAPS season indicates that the bird is not a SY or HY bird:

- B - Basic: Feather tract comprised entirely of basic (or a mix of basic and alternate) feathers (note that basic feathers = “adult basic feathers” in Pyle 2022), but no juvenile or formative feathers. The use of this code during the MAPS season on an adult bird prior to its prebasic molt indicates that it is an ASY bird; the use of this code during the MAPS season to describe feather tracts on an adult bird after its prebasic molt indicates only that it is an AHY bird.

Individuals of some near-passerine species (e.g., woodpeckers) can be aged to TY or ATY during the MAPS season (see discussion in Pyle 2022, pgs. 41-42) due to incomplete molts, which result in feathers that are retained through the next prebasic (not preformative) molt. Such individuals can have up to three generations of juvenile and basic feathers present within the same feather tract (these species do not acquire alternate feathers). Two codes are to be used to distinguish cases in which juvenile and basic (rather than juvenile and formative) feathers are present, from situations in which two generations of basic (rather than formative and basic) feathers are present:

- R - Retained: Both juvenile and basic (rather than juvenile and formative) feathers are present within the tract (e.g., see Figs. 25 and 26 in Froehlich 2003). The

use of this code during the MAPS season on an adult bird prior to its prebasic molt indicates that it is a TY bird; the use of this code during the MAPS season to describe feather tracts on an adult bird after its prebasic molt indicates that it is a SY bird. **Typically, only used for woodpeckers.**

- M - Mixed: Multiple generations of basic feathers are present in the tract (e.g., see Fig. 27 in Froehlich 2003). The use of this code during the MAPS season on an adult bird prior to its prebasic molt indicates that it is an ATY bird; the use of this code during the MAPS season to describe feather tracts on an adult bird after its prebasic molt indicates only that it is an ASY bird. **Typically, only used for woodpeckers.**

The following two codes, which can be used during the MAPS season (prior to the prebasic molt) to distinguish adult (AHY) from young (HY) birds, are generally not useful for ageing adult birds to more specific age classes (i.e., SY, ASY, TY, ATY):

- A - Alternate: body feather tracts contain at least some alternate feathers – **or** – the **entire tract of wing coverts** is of alternate plumage. If **ANY** juvenile, formative, or basic feathers are present in a wing covert tract, the alternate feathers should be ignored and the code for the feather tract should be based on the other feathers, that is “J”, “L”, “F”, or “B”.
- N - Non-juvenile: Feathers in this tract are definitely not juvenile feathers (or the non-feathered body part is not characteristic of a young bird), but whether or not they are formative or basic feathers cannot be determined with confidence. Note that if primary coverts are coded “J” and a molt limit exists between the primary coverts and the secondary coverts, the secondary coverts must be formative feathers and, thus, must be coded “F”, not “N”, even though formative and basic secondary coverts might be indistinguishable from each other. The code “N” should only be used as a last resort; every effort should be made to identify appropriate feather tracts to formative or basic. Often, this is best accomplished by considering the tract in the context of other tracts which, for example, have perhaps been reliably aged juvenile. This code should also be used for NON-FEATH if non-feathered body parts show characteristics indicative of an adult bird.

The following code should be used for feather tracts on HY birds of those species that have an auxiliary preformative molt and who have molted out juvenile feathers into the auxiliary formative feathers:

- X - Auxiliary preformative: This code should be used for feather tracts that were replaced during the preformative auxiliary molt. These will be mostly body feathers and is generally used only for species in the Cardinalidae family, and potentially for some species in the Passerellidae family.

The following code should be used for feather tracts examined, but not meeting any of the above criteria:

- U - Unknown: This code should be used for any feather tract or non-feathered body part that is examined, but that shows ambiguous characteristics or that cannot be coded with confidence.

Finally, LEAVE BLANK any field representing a feather tract or non-feathered body part that was not examined for any reason, including cases where that feather tract provides no useful information for ageing the bird.

As an example of the use of these fields, consider the age determination of a SY bird (i.e., AGE = 5) prior to its prebasic molt. The age of SY birds can be determined by the retention of juvenile feathers, which will be evident in some feather tracts but not others (depending on the extent of the preformative molt). Any feather tract for which retained juvenile feathers are evident will have either a "J" or "L" entered in its field, depending on whether molt limits are between or within feather tracts, respectively. If the molt limit is between feather tracts, the tract with juvenile feathers would be coded "J" and the tract with formative feathers would be coded "F." If the molt limit is within the feather tract, the tract would be coded "L." In each of these cases where a molt limit between juvenile and formative feathers can be discerned, the bird should be aged by molt limit (HOW AGED = L). If, however, a molt limit cannot be discerned, but the juvenile feathers present can be distinguished as juvenile (as opposed to basic) feathers by their appearance alone (i.e., color, shape, quality, or wear), the bird would be aged by plumage (HOW AGED = P). Remember, any feather tract or non-feathered body part that was examined, but for which a code could not be determined, should have a "U" entered in its field.

As another example, consider an ASY bird (i.e., AGE = 6) prior to its prebasic molt. Birds of this age are typically distinguished by having undergone a complete prebasic molt B adjacent feather tracts generally show little if any contrast in quality or wear. Such birds should have a "B" entered in all fields for which the basic feathers present can be distinguished as basic (as opposed to juvenile) feathers by their appearance alone (i.e., color, shape, quality, or wear), and should be aged by plumage (HOW AGED = P). They should not be aged by molt limit (HOW AGED = L) because there is no molt limit. Note that any alternate feathers present provide no information as to whether the individual is a SY or ASY bird.

As a third example, consider a species that can undergo a complete preformative molt (e.g., a Northern Cardinal). When examining an adult of these species during the breeding season, you may find that all of the feathers are of a single generation (i.e., no molt limits). Because formative and basic feathers appear identical in this species, you will not be able to age the bird specifically to SY or ASY and so the bird must be aged AHY (i.e., AGE = 1). Such birds should have "N" entered in all fields for which the formative or basic feathers present can be distinguished as non-juvenile feathers by their appearance alone (i.e., color, shape, quality, or wear), and should be aged by plumage (HOW AGED = P). If a molt limit is present in these species, the limit must be between juvenile and formative feathers and "N" should not be used in any field because evidence for the existence of formative feathers is provided in context by the presence

64 - 2025 MAPS Manual

of juvenile feathers. Therefore, the bird must be aged SY (AGE = 5). Birds of species that can undergo a complete preformative molt can never be aged ASY (AGE = 6) in the field.

Finally, it is possible that various feather tracts in an individual bird will show conflicting characteristics (i.e., characteristics that indicate different age classes). When making an age determination for such a bird, give more weight to tracts that are more reliable or have the most obvious reliable features. Although it is not necessary that all tracts in a record agree, you should be confident in your ultimate age designation. During the MAPS season, a bird with no reliable feather tracts or a bird for which conflicting characteristics make age determination difficult should be aged as AHY (AGE = 1) prior to the prebasic molt and as indeterminable (AGE = 0) after the prebasic (or preformative) molt.

WING – Wing Chord. Record wing chord (the length of the unflattened wing) to the nearest mm. See Pyle (2022) or Ralph *et al.* (1993) for an explanation of the technique. Unless there is little or no overlap in wing lengths between sexes (e.g., icterids), DO NOT sex birds by wing length alone in the absence of population-specific wing-chord data.

BODY MASS – Using an electronic balance, record the mass of the bird to the nearest tenth of a gram. If an electronic balance is not available, record the weight of the bird to the nearest 0.5 gram using a Pesola (or other spring-operated) scale.

STATUS – Record status as a single, three-digit code. These codes are available at the Bird Banding Offices Bander Portal (https://www.pwrc.usgs.gov/bbl/Bander_portal) under the “Reference” tab as one of the drop down “Lookup Tables”. You must sign in to have access to this table. The most-frequent codes are “300” - normal wild bird captured, banded, and released; “301” - normal wild bird captured, banded and color-banded, and released; and “500” - injured bird (see DISP). In addition to these standard status codes, please use code “000” for all birds that were not banded or that died prior to release. Birds that die prior to release should also have a D or P in the DISP field. Please note that status “000” birds are now requested to be included in schedules submitted to the banding offices if they are mortalities. Live released unbanded birds with status “000” are not to be included in Bander Portal submissions.

DATE (MO/DAY) – Month/day. Record the date of capture as month and day, all in numbers. The year is entered once on the top of the form. Record all months and days as two-digit numbers (i.e., June is written “06”). The first entry on a page for each date must be written out completely; subsequent entries for that date may be entered as “>.”

CAPTURE TIME – Using the 24-hour clock, record, to the nearest 10 minutes, the starting time of the net run on which the bird was extracted. Thus, all birds extracted (or escaping) on a given net run will have the same capture time. This is necessary for standardizing effort between years. Do not enter the time at which the bird was extracted, processed, or released. Always enter three digits. Note that the ultimate zero is preprinted on the form; e.g., 6:24 a.m. = 062(0), 1:48 p.m. = 135(0).

STATION – Record the four-character code for the MAPS station as determined during station registration.

NET – Enter a two-digit, numeric code (e.g., “06”) for the net site at which the bird was captured. It is important that net codes not include alpha characters nor be more than two characters long. Please enter “?” into this field if the net number is unknown.

DISP – Disposition. Enter a code from the list below indicating the final disposition of any injured or dead bird. A bird is considered “injured” if its survival probability is thought to be compromised, or for healed injuries, could previously been compromised; thus, a minor flesh wound or loss of a few feathers is generally not worthy of note. Any injured or dead bird also should have a status code of “500” or “000,” respectively.

- B - Body injury.
- D - Death due to a cause other than predation.
- E - Eye injury.
- F - fouled feathers, typically from oil
- I - Ill or diseased.
- L - Leg injury.
- M - Malformed (deformity such as crossed mandibles)
- O - Old (healed) injury.
- P - Predator-caused mortality.
- R - Band removed from bird and then bird released bandless. Only should occur for leg injuries where other leg cannot be banded.
- S - Stress or shock.
- T - Tongue injury.
- W - Wing injury (unable to fly).

NOTE NUMBER – Enter a number (starting with “1” on each page) if additional information needs to be recorded, and record this information with the corresponding note number in the NOTE field on the back of the banding-data sheet. Occasionally, individual records that technically cannot be considered MAPS data are included on MAPS banding-data sheets. These records must be identified as non-MAPS in order to avoid including them in constant-effort analyses. **Mark these records by recording “NM” in the NOTE NUMBER field.** Occasionally notes associated with a record indicate that the species determination for a recapture or an unbanded bird was uncertain. **Mark these records by recording “QS” in the NOTE NUMBER field.**

FTHR. PULL – Enter a code from the list below indicating which feathers were pulled during this capture event. Only record this information when the feathers are actually pulled, not on a recaptured bird that has previously had feathers pulled. If no feathers were pulled, leave the field blank.

- O - Outer two rectrices were pulled (i.e., rectrix 6 from both the left and right side of the tail). Previously, this was indicated by FTHR. PULL = P.

66 - 2025 MAPS Manual

- X - R3 was pulled from both sides of the tail.
- I - An inner and an outer rectrix were pulled (i.e., rectrix 1 from one side and rectrix 6 from the other side were pulled).
- C - contour feathers only were pulled for feather sampling purposes. If both flight rectrices and contour feathers were pulled, use the code pertaining to the rectrices pulled.

NOTE -- Record notes on the back of the banding-data sheet. These include characterizations of examined feather tracts in adult birds (see AHY/SY/ASY/TY/ATY above). Other examples of notes include measurements of difficult-to-identify species such as *Empidonax* flycatchers; documentation of rarities or extralimital species; suspected age or sex determinations of birds given age code "0" or "9" or sex code "U" or "X"; details of any "O" (other) code for HOW AGED or HOW SEXED; explanations for injured, dead, and unbanded birds; and sequence of color bands, if present. Please be liberal in your note-taking, especially to indicate which, if any, flight feathers are missing, erupting, or growing.

Tail and tarsus measurements are not explicitly asked for on the data sheet but you can enter them in the NOTES field. In MAPSNet there are fields where they can be entered. Please ensure you are following the directions on how to take these measurements from Pyle (2022).

A strategy for ageing and sexing birds

A useful strategy for ageing and sexing is to complete the skull, cloacal protuberance, brood patch, molt, wear, and juvenile plumage fields first (that is, after identifying the bird and banding it or, if a recapture, reading and recording the band number). In most cases, ageing and sexing should be straightforward if you keep the following breeding-season "rules" in mind:

- A skull of 0 to 4 indicates a hatching year bird (except in a few species such as cuckoos that may never progress beyond 4); a skull of 5 or 6 indicates an adult.
- Presence of a CP indicates an adult male (except in Wrentits).
- Presence of a BP indicates an adult and, in most species, a female (for exceptions see "Sex" in individual species accounts (Pyle 2022) or "List of species in which males can develop brood patches" at <http://www.birdpop.org/pages/mapsDataForms.php>) on the IBP website. A full (class 3) BP indicates a female (except in cuckoos, kingfishers, woodpeckers, Clark's Nutcracker, and Wrenit, in which males develop full brood patches).
- Heavy body molt in the absence of symmetric flight-feather molt generally indicates a hatching year bird. Conversely, symmetric flight-feather molt, especially of the primaries, indicates an adult in most species (see "Molt" in individual species accounts (Pyle 2022) for exceptions).
- Flight-feather wear of 3 or greater generally indicates an adult.

- Presence of juvenile body plumage indicates a hatching year bird.

If your data conflict with one another, look again! **And remember that the absence of CP, BP, molt, or wear is not, by itself, conclusive evidence of anything!**

If you use how-aged codes of "L" or "P", you must fill in at least one of the first seven MOLT LIMITS & PLUMAGE fields. Also, if you have an adult bird prior to its prebasic molt, please go on and attempt to age it more specifically to SY, ASY, etc., by recording the feather generations of the relevant feather groups (or soft-part features) in the appropriate MOLT LIMITS & PLUMAGE fields.

SUMMARY OF MIST-NETTING RESULTS

This summary serves as a check to ensure that all of your capture records C for newly banded, unbanded, and recaptured birds C have been submitted and that the dates and station are correct for each capture. Please complete this form using your raw banding-data sheets at the end of each banding day. Please do not fill out this sheet from your computer file! Remember to count changed and added bands only once (as recaptures) and to omit lost and destroyed bands. Refer to Figure 10 for an example.

Instructions for completing the Summary of Mist-Netting Results form

Location: Record your four-character location code.

Station: Record your four-character station code.

Intended Period: Record the intended period for the date operated.

Date: Record the month and day of the date operated.

New: Record the number of new individuals banded. Remember, if a bird dies before processing, this individual should be recorded on the data sheet as an unbanded bird. The band that was applied to the individual should be taken off the bird and be recorded as destroyed.

Unbanded: Record the number of birds captured but not banded.

Recaps: Record the number of recaptures. Remember that previously-banded birds that escape or are inadvertently released before the band number is read should be recorded and counted as recaptures (BAND NUMBER remains blank on the banding data sheet).

Total: Tally the number of new, unbanded, and recaptured birds for each day of operation. At the end of the season, record the totals of these three categories at the bottom of the form.

BREEDING STATUS LIST

Many of the individual adult birds captured in the MAPS program are transients that do not breed at the MAPS station. They include floaters that have not yet acquired a breeding territory, failed breeders searching for a new mate or new breeding territory, and post-breeding individuals dispersing from breeding territories to molting and pre-migration staging areas. Because the presence of such transient individuals negatively biases adult survival rates, we use a transient modification (Pradel *et al.* 1997) of Cormack-Jolly-Seber mark-recapture models (Pollock *et al.* 1990) to estimate the survival rate of resident individuals and the proportion of residents among newly captured birds. We suggest, however, that the proportion of residents in the adult population may be more than a simple nuisance parameter. Rather, we suggest that this proportion may vary in a predictable manner as a function of population change and, thus, may be of fundamental importance to avian population dynamics. In order to obtain the most useful measure of the proportion of residents, we pool data for a given species only from those stations at which the species is a regular or usual breeder, that is, only from those stations where at least one individual of the species was known to be a summer resident attempting to breed during more than half of the years the station has been operating.

As part of MAPS protocol, therefore, we ask MAPS operators to record breeding status information on all species seen or heard during each visit to each station. We ask operators to record anecdotal observations of active nests; birds carrying nesting material, food, or fecal sacs; distraction displays; courtship; copulation; and territorial singing or drumming using a protocol similar to that employed in Breeding Bird Atlas projects. These observations are recorded on the stations' Breeding Status List. Using these data, coupled with capture data, MAPS operators are asked to determine the breeding (summer residency) status of all species at each station each year.

In addition to providing unbiased data on the proportion of residents in the population, this protocol provides a unique and extremely valuable database, one which allows the construction of temporally and spatially explicit species-habitat relationships based on actual breeding status at each of the hundreds of MAPS stations. This database can overcome many of the limitations of traditional species-habitat relationships derived from point-count data. Such limitations are caused by including species as breeders that are in fact only transients at the location in question; and by excluding actual breeding species at the location in question because they are not encountered within the short duration of most point counts.

It is important to understand that what we are asking you to determine by asking for breeding status is **whether or not any portion of at least one breeding territory or home range of a given species includes any portion of the area of your MAPS station.** Remember, breeding status is determined only for the area contained within the boundaries of your MAPS station, NOT the preserve, county, or any other area in which your station is located. Remember also that the boundaries of your station include all of

the area extending outward for 100 m from your outermost nets. In general, typical MAPS stations include an area of about 20 ha (50 acres).

Station-specific Breeding Status Lists (Fig.10) are included with the spring email sent to MAPS operators in mid-April. Operators of stations from which no previous data have been received by IBP will receive a blank Breeding Status List with the four-letter alpha location code, four-letter alpha station code, unique five-number numerical station code assigned by IBP ("STA"), and the current year preprinted on the form. Before the season begins, such operators should list in the SPECIES CODE field, the four letter code for the species (e.g., "BCCH" for Black-capped Chickadee) from the most current *Four-letter and six-letter alpha codes for birds recorded from the American Ornithological Society checklist area* (Pyle and DeSante 2003). This list (further updated in Pyle and DeSante [up through 2023 from Chesser et. al 2023]) can be downloaded from The Institute for Bird Populations' website at <https://www.birdpop.org/pages/birdSpeciesCodes.php>. Operators of stations from which previous MAPS data have been received by IBP will receive a preprinted Breeding Status List that lists all species ever encountered at the station in American Ornithological Society (AOS) checklist order. (If data were not submitted using MAPSNet, species recently added to the list by the operator may not be included on the preprinted list because of a backlog in data entry and processing at IBP). Species that are encountered at the station during the MAPS season, but that are not on the preprinted (or anticipated) list, should be added to the end of the list. At the end of the season, the completed Breeding Status List will thus include all species **detected** at the station during **all** breeding seasons that the station has been operated, not just those species detected during the current year or only those species for which individuals have been captured.

Breeding Status Lists for stations from which previous MAPS data has been received will also contain filled-in BRSTAT codes for each species. The BRSTAT code is the cumulative breeding status at the station for all previous years of operation, a kind of breeding status summary. First time operators' Breeding Status Lists will have a dash preprinted in BRSTAT. For each species, the BRSTAT code represents a summary of all of the yearly breeding status codes. The following BRSTAT codes are in use:

- B - Regular breeder. Summer resident or suspected summer resident during all years the station was operated.
- U - Usual breeder. Summer resident or suspected summer resident for more than ½ of the years the station was operated, but not all years.
- O - Occasional breeder. Summer resident or suspected summer resident for ½ or fewer of the years the station was operated.
- T - Transient. The station lies within the species' breeding range, but no individual of the species was a summer resident at that station during any year.
- A - Altitudinal disperser. A species which breeds only at lower elevations than that of the station, and which disperses to higher elevations after breeding.

- M - Migrant. The station falls outside of the species' normal breeding range.
- ? - Unidentified. Individuals of the taxon were not identified to species; no breeding status was assigned.

These codes are provided to inform you of the overall breeding status for each species. It is important that you **determine each new year's breeding status independently of the BRSTAT.**

PERIOD STATUS: The Period Breeding Status (i.e., Confirmed - C; Probable - P, Observed - O) of each species encountered during each period of operation at each station must be recorded on the list for that station, either during the day as the birds are detected or at the end of the day before leaving the field. Record, using upper case letters, the highest hierarchical Period Breeding Status (Table 3 or Fig. 10) detected for each species that period; and, using lower case letters, the appropriate Daily Behavior Sub-

Table 3. Hierarchical categories of Period Breeding Status (upper case) and associated Daily Behavior Sub-Codes (lower case) for MAPS Breeding Status List

The criteria used to designate Period Breeding Status are as follows:

Confirmed (C): The following criteria confirm a species as a breeder:

- (n) current year's nest found in the study area with eggs or young, in the process of being built, or already depredated or abandoned;
- (m) adult seen gathering or carrying nesting material to a likely nest site in the study area;
- (v) egg seen in the oviduct with bird in hand
- (f) adult seen carrying food or fecal sac to or from a likely nest site in the study area;
- (d) distraction display or injury feigning by an adult bird;
- (l) capture of a young bird incapable of sustained flight (a "local"), or very young (stub-tailed) fledglings being fed by parents in the study area.

Probable (P): The following criteria suggest, but do not confirm a species as a breeder:

- (c) copulation or courtship observed of a species within its breeding range;
- (t) other territorial behavior observed in the study area;
- (s) territorial song or drumming heard.

Observed (O): The following criteria indicate the species was detected, but with no evidence of local breeding:

- (b) bird captured or banded. NOTE: The presence of a brood patch or cloacal protuberance on a single individual is not valid evidence of local breeding;
- (e) bird encountered (seen or heard) in the study area but with no territorial behavior;
- (o) bird was encountered flying over the study area
- (z) bird was both banded and encountered

Absent (--): The species was not encountered during that period.

Codes (Table 3 or Fig. 10) Associated with that Period Breeding Status. Note that sub-codes can only be combined with other sub-codes at the same breeding status level. For example, 'Cf' and 'Obe' are acceptable combinations; 'Cs' and 'Obs' are not. Use a '-' to indicate that a species was not observed in a given period. Note that the certainty of Period Breeding Status codes likely will decrease for most species as the season progresses and breeding behavior diminishes. For example, a species recorded as a probable breeder in May and confirmed as a breeder in June may drop back to a probable breeder in July and show no signs of breeding (or disappear altogether) by August. Please also include observations at the station that were not from the banding day when determining the code for the period. E.g., if a species was only observed during the banding day but during a subsequent visit to the station the next day for non-banding purposes the species was heard singing, the period code should be Ps, not Oe.

YEAR STATUS: The current year breeding status. At the end of the season, review your period status codes and enter the apparent breeding status for the current year for each species in the right-hand column, using one of the following nine categories:

Breeder (B): Summer resident. A Breeder is a species within its normal breeding range that is confirmed or determined to be a breeder or summer resident within the station (i.e., at least one individual was determined to reside at least partly within the station boundary during the breeding season of the year under consideration). It needn't be proven that the species actually bred, or even found a mate. Summer residents outside their normal breeding range should be given the code "E" (see below).

A species automatically qualifies for a Breeder ("B") Year Status if it was given a Confirmed ("C") Period Breeding Status in one or more periods. Thus, a current year's nest found in the study area with eggs or young, in the process of being built, or already abandoned or depredated qualifies the species for a "B" Year Code, as does the sighting of an adult carrying nesting material, food, or a fecal sac to or from a likely nesting site, or doing a distraction display or feigning injury within the station. The sighting of very young (stub-tailed) fledglings being fed by parents within the study area also qualifies the species for a "B" Year Code. Probably the most common means of classifying a species as a Breeder is by the presence of at least one territorial (singing or drumming) male in the study area throughout the breeding season. Note that such territorial behavior is coded "Ps" (probable breeder) for individual periods, but if it occurs over much of the season, the species should be considered a Breeder' ("B") rather than a Likely Breeder ("L"). Multi-period observations of courtship, copulation, or other territorial or mating behaviors also qualifies the species as a Breeder, especially if coupled with song or drumming in other periods. In summary, note that it is acceptable to assign a year status of "B" to a species that exhibits persistent territorial singing during the height of the breeding season, as well as to those confirmed by nest sightings, fledglings or other "hard evidence" of breeding activity.

Banding data are also useful for determining breeding (summer residency) status. Within-year recaptures or resightings of an adult, at least seven days apart and with at least one occurrence during the height of the breeding season, indicates a summer resident. The capture of a single adult in breeding condition (i.e., with a greatly



The INSTITUTE for BIRD POPULATIONS

2025 MAPS BREEDING STATUS LIST

List all species ever encountered at the station!

Location: **B I P A** Station: **U P E D** Sta: **1 6 6 9 8**

Period Status Codes: Please record only the highest hierarchical breeding status observed during each period (C supersedes P and O; P supersedes O), and the appropriate daily sub-codes (hierarchically listed) describing the behavior or observation indicating that status (e.g., Cn, Pc, or Ob). If a species wasn't observed, use a ---. If a station was operated two or more days in a period, record only the highest breeding status observed.

<p>C = Confirmed Breeder n = current year's nest found m = carrying nest material v = egg in oviduct f = carrying food or fecal sac d = distraction display I = local bird</p>	<p>P = Probable Breeder c = courtship/copulation t = other territorial behavior s = song/drumming</p>	<p>O = Observed b = banded/captured e = encountered o = flyover z = both banded and encountered</p>	<p>--- = Absent</p>
--	---	--	---------------------

ENTER DATE (mm/dd) FOR INTENDED PERIOD BELOW

SPECIES CODE	BRSTAT	DATE										2025 YEAR STATUS
		/	/	05/28	06/05	06/14	06/24	07/06	07/13	07/25	08/04	
HAWO	T			C	C	Ps	Ps	C	C	C	C	L
ACFL	U			Ps	Ps	Ps	Ps	Ob	C	C	Oe	B
YTVI	O			C	C	C	Ps	Ps	C	C	C	L
REVI	B			Ps	Ps	Ps	Cn	Cnf	Ps	Ob	C	B
TRES	T			C	C	Oo	C	C	C	C	Oo	T
CACH	U			Ps	Ps	Ps	Ps	Ps	Ps	Ps	Ps	B
WOTH	B			Ps	Ps	Ps	Ps	Ps	Ps	Oe	Oe	B
GCTH	M			Ob	C	C	C	C	C	C	C	M
AMRO	B			C	Ps	Ps	C	C	Oe	C	C	L
GRCA	B			Ps	Ps	Ps	Ps	Ps	Oe	Oe	C	B
NOPA	O			Oe	Ps	Ps	Ps	Oe	Ps	Ob	Obe	B
YWAR	T			C	C	C	C	C	C	Ob	Obe	T
AMRE	O			C	C	C	C	C	C	Ob	Oe	T
OBSERVER'S INITIALS				CC	CC/ PB	CC	CC/ AB	CC	CC	CC	CC/ AB	

BRSTAT: Cumulative breeding status for all previous years of operation
B = Regular Breeder (all years) **A** = Altitudinal Disperser
U = Usual Breeder (>, not all, years) **M** = Migrant
O = Occasional Breeder (≤ years) **?** = Uncertain Species ID
T = Transient

2025 YEAR STATUS: Current year breeding status
B = Breeder **M** = Migrant
L = Likely Breeder **E** = Extralimital Breeder
T = Transient **?** = Uncertain Species ID
A = Altitudinal Disperser --- = Absent
H = Higher Altitude (than usual) Breeder

Figure 11. Completed MAPS Breeding Status List

enlarged cloacal protuberance or a heavily vascularized brood patch) is not sufficient evidence to classify the species as a breeder because failed breeders often wander widely before losing their BPs and CPs. However, several individuals showing breeding characteristics at various times during the season could warrant a “B” designation for the species for the year.

Likely Breeder (L): Probable summer resident. A species within its normal breeding range that was suspected to be a breeder or summer resident but was encountered somewhat infrequently during the breeding season of the year under consideration is classified as a Likely Breeder. This code permits a degree of uncertainty when determining single-year breeding status and need be used only for species that were suspected summer residents *but were encountered infrequently* during the field season. We suspect this may happen with species that reside on the fringes of the station or are difficult to detect. To avoid the uncertainty associated with the status “L,” all efforts should be made to assign one of the more definitive status codes to each species whenever possible.

Please note that while continual territorial singing throughout the breeding season would merit a “B” status, singing on only two or three different days may indicate an “L,” or even a “T.” In such cases, one should consider the likelihood of summer residency in terms of habitat suitability for that species, the dates on which the singing occurs, and any behavioral knowledge of that species. As shown in Figure 11, two instances of a Hairy Woodpecker drumming in June at a station within its breeding range (and with no additional records) likely (but not definitively) indicates summer residency, and thus merits a year status of “L.” However, two instances of Swainson's Thrushes singing in May within their breeding range, again with no additional records, would *not* suggest a single-year status of “L,” as this species is known to sing during migration and to sing well into the season if a summer resident. The year status, in this latter case, should be “T.” When the cumulative breeding status (BRSTAT) is calculated, years coded “L” are treated as if they were coded “B.”

Transient (T): A species that breeds in the general area of the station (perhaps even less than a kilometer away) but, because of habitat or patchy distribution, does not breed at the station is classified as a Transient. In order to qualify as a Transient, the station must lie within the breeding range of the species, but no individuals of the species can be thought to be breeders or summer residents within the station (see above definition of “Breeder”). Transient individuals may be adults within their normal breeding range that move through the station during the breeding season but do not establish a territory or home range within the station boundaries. Early in the season, such adult individuals could be birds still in migration, birds that have completed migration but not yet established territories, or birds that might never establish territories that year (floaters). For example, capturing one or two individuals of a rarely observed species in June with well-developed CPs or BPs would not permit categorizing that species as anything other than a transient. Mid-season transients could be failed breeders from beyond the station boundary that are simply moving through the station. Later in the season, transient individuals could be adults or young in post-breeding

dispersal, or even very early individuals in fall migration. A species may be a Transient at one station within a location and a Breeder at another station.

Altitudinal Disperser (A): A species which breeds only at lower elevations than that of the station and which disperses to higher elevations after breeding. In the Sierra and, to a lesser extent, the Cascades and other western montane areas, this is a common phenomenon for Orange-crowned Warblers, Nashville Warblers, and House Wrens. In order to qualify as an altitudinal disperser, the station must lie upslope from the breeding range of the species.

Higher Altitude (than normal) Breeder (H): An altitudinal disperser that has resided during the height of the breeding season (not just during the post-breeding period) in a given year above its normal breeding elevation. When the cumulative breeding status (BRSTAT) is calculated, years coded "H" are treated as if they were coded "A."

Migrant (M): The station does not lie within the breeding range of the species, and the species did not reside at the station during the breeding season. Migrant species may pass through the station on migration, or reside through the winter. Specifically, the species' breeding range, as delineated by range maps and descriptions, does not include the specific geographic location of the station. (The primary references we use are the range maps from *Birds of North America Online*, *The Sibley Field Guide to Birds of Eastern North America* and *The Sibley Field Guide to Birds of Western North America*. We also use National Geographic's *Field Guide to the Birds of North America*, Peterson's *Field Guide to Western Birds* and *Field Guide to Eastern Birds*, *Birds of North America Online* (Rodewald 2015) and range descriptions in the A.O.U. *Checklist of North American Birds* [1957 for subspecies, 1983, and 1998]; and status codes in DeSante and Pyle's *Distributional Checklist of North American Birds*, 1986).

There is one important exception to the above definition of Migrant. If a station lies within a mountain range at a higher elevation than a given species breeding range, but adults and young of the species habitually move through it during post-breeding and juvenile dispersal, respectively (as in a foothill species that disperses upslope), the species should be classified as an Altitudinal Disperser (A; see above) at the station rather than a Migrant (M). The inclusion of capture data for such species from such stations can provide important information for regional productivity indices. A Migrant status for such a species would cause it to be overlooked during productivity analyses. Finally, do not confuse the terms "Migrant" and "migratory"; migratory species can be classified as Migrants, Altitudinal Dispersers, Transients, Likely Breeders, or Breeders.

Extralimital Breeder (E): A summer-resident species that is outside of its normal breeding range. As with species given a code of "B," it need not be proven that the species actually bred, or even found a mate; merely residing at the station during the breeding season is sufficient to warrant a code of "E." These vagrant individuals are not given a code of "B," as they are unlikely to return in subsequent years; in pooled analyses, a "B" code could bias survivorship estimates for the species. When the cumulative breeding status (BRSTAT) is calculated, years coded "E" are treated as if they were coded "M."

Unidentified (?): This code is used primarily for observed (non-captured) individuals that were not identified to species. Examples include UNGU, UNCR, UNSW. This code is also used if an unidentified individual was captured but not banded, for example UNHU or an unbanded UEFL. A species that has been banded would receive a breeding status of “?” if, during verification, the species identification of an individual became uncertain and no other individuals of that species were encountered in that year.

Absent (--): No individual of the species was detected at the station C neither captured, heard nor seen C for the duration of the MAPS field season in the year under consideration. This code applies both to species previously captured/encountered at the station in a past field season and to those anticipated but not yet encountered.

OBSERVER’S INITIALS: Record the initials of the people involved with collecting breeding status data for each intended period.

Please note that it is very important to classify each species correctly, to the best of your ability, within the boundaries of your station (i.e., within 100 m of nets). Inclusion of data for a species in mark-recapture analyses from stations where it does not regularly or usually breed will deflate estimates of proportion of residents and lower the precision of survival-rate estimates as well. However, transients can be included in the calculation of productivity indices. Bear in mind that the Breeding Status List is annual in nature (i.e., you will consider breeding status of each species on a year-by-year basis) and that a species’ year status may change from one year to the next. Generally, such changes will be a species changing from Breeder to Likely Breeder or Transient or vice versa, but occasionally a species can change from an Extralimital Breeder or Migrant to a Breeder or Transient (or vice versa) as its breeding range changes.

It is important to remember to assign a breeding status each year to all species ever captured or encountered **at the station, and not just to those that were captured during the most recent field season.**

Occasionally, during the verification process, recapture data or other information come to light that require re-determination of Breeding Status Codes for various species at a station. IBP biologists may change some Year Codes and will want you to examine their changes.

Additionally, if non-MAPS data is included with your submission, the following three additional codes may be used in your breeding status files:

- D = The species was only encountered at the station outside of the MAPS season, but the station lies *within* breeding range of the species.
- W = The species was only encountered at the station outside of the MAPS season, and the station lies *outside* of the breeding range of species.
- @ = The Breeding Status List is missing or incomplete for this species this year.

HABITAT STRUCTURE ASSESSMENT (HSA)

The Monitoring Avian Productivity and Survivorship (MAPS) program is focused on monitoring avian demographic parameters and relating their spatial and temporal variation to landscape-scale ecological and environmental data. For this purpose, we utilize geographic information systems (GIS), remote sensing of habitat types, and existing vegetation maps from various sources, at a variety of spatial resolutions (e.g., from 1km to 30m cells). Many of these spatial data have greatly improved in quality in the last few years and we are depending on them for much of our vegetative analyses. However, there certain pieces of data, such as the specific plant communities and vertical structure of the vegetation associated with the habitat types that must be collected directly at the station.

The horizontal pattern, vertical structure, and type of the vegetation within a MAPS station can affect the number and diversity of breeding birds present, as well as the efficiency with which birds can be monitored by mist-netting. Many ecological studies show that habitat structure is an important factor for predicting avian diversity and abundance. It has been shown, in general, that spatial habitat patterns and vertical habitat structure are good predictors of the presence and relative demographic success of component species in avian communities. Furthermore, changes in the vegetation at a station may cause changes in the breeding bird community or affect how attractive the site is to dispersing birds. For all these reasons, the Habitat Structure Assessment (HSA) protocol is designed to describe the type and distribution of the vegetation at each monitoring station.

MAPS banding data can be combined with remotely sensed datasets, habitat structure assessment data, breeding status lists (BSL), breeding/overwintering range data, and various weather and other environmental data to tackle some aspects of avian population conservation. Analyses of these data will help guide efforts towards the ultimate goal of providing management solutions to the problems of declining avian populations, and to make these solutions available to a variety of land stewards. The success of this approach will depend in large measure upon the participation of station operators in applying the HSA protocol.

HABITAT TYPES

The goal is to describe the arrangement and coverage of up to five (but **usually one or two**) discrete and recognizable habitat types at your station and, within each of those habitat types, to describe the arrangement and coverage of component vertical layers. We provide a blank grid map to help you map and classify the main habitat types within your station but this can also be done using GIS software, if you have it.

The methods outlined in these instructions are designed to provide useful information with a minimum of effort. Please keep in mind that the purposes of these maps and descriptions are:

- a) to provide a general classification and characterization of the habitat of the study area to allow for broad comparisons and groupings among stations;
- b) to provide a method for monitoring major changes in the vegetation that occur as a result of natural successional change, new management practices (logging, grazing, development), or the occurrence of major "catastrophes" such as fire or flood; and
- c) to provide a relatively rapid assessment of the habitat structure and spatial patterns of vegetation.

Once the results of these assessments are analyzed, we will be able to group stations in terms of features to which a chosen target species may respond.

Habitat Structure Assessments should be conducted every five years, unless the habitat at the station has undergone a major change (e.g., fire, hurricane, logging, construction, brush-clearing, etc.). We ask that contributors take a copy of their station map and completed HSA forms into the field every 5 years at the appropriate time (see below) and verify that the information is correct, and has not significantly changed. These vegetation assessments **should be made during the fourth period of recommended operation** (usually the time of maximum canopy and shrub cover), e.g., stations beginning in Intended Period 4 should do their HSA in Intended Period 7.

It may be helpful to follow the step-by-step approach to preparing your station map and habitat description(s) as outlined on the following pages.

A STEP-BY-STEP APPROACH:

Step1: Prepare a station map. The station map should depict the locations of your nets and the major physical features of the study area. A typical MAPS station consists of about ten 12-m mist nets dispersed rather uniformly over an 8-hectare (20-acre) core area. Plot the exact location, orientation, and designation (net number) of each of your nets in the central portion of a copy of the 24 x 24 cell grid map (Form H0: MAPS Station Map) provided in the HSA protocol. The scale of each cell on this map should normally be 30m on a side. If your station is very long you might need to increase the size of a map cell from the standard 30m to perhaps 40m. If your station is very small (because you have your nets spaced closely together; see below), you might need to decrease the size of a map cell down to 20m. Be sure to indicate the scale on your station map by using the scale bar provided. In addition, plot also the exact locations of all the important natural and human-made physical features such as lakes, ponds, streams, ditches, roads, trails,

buildings, and other structures. Be sure also to indicate magnetic north on your map. If possible, obtain an aerial photograph or GIS output of the site, use it to help locate the nets and other physical features on your station map, and submit it to IBP in addition to the station map. It is important to mark clearly the exact location and orientation of each net on the station map. This will assure that the location and orientation of individual nets will remain consistent if the personnel operating a station changes from year to year.

Step 2: Identify the boundary of your station and draw it on the station map. The boundary of a MAPS station is defined to include all areas that lie within 100m of the outermost nets. Remember, 100m is equal to over eight standard (12m) net lengths; you may be surprised at how much area this encompasses! On the map, however, 100m is a distance equivalent to three and one-third cells on the 30m grid (see heavy dashed line on Figure 12). The station boundary itself can be determined by first drawing a circle with radius equal to 100m centered at the outermost end of each net. This should produce a set of overlapping circles. Then connect the outermost points of the outermost circles with a smooth, straight or slightly outwardly convex line. This will give you the boundaries of your station. If you have nets that are more than 200m removed from all other nets (producing non-overlapping circles), a corridor about 100m wide between these nets and encompassing the route traveled to reach them should be included within the station boundary. In this way, all stations will be single units regardless of layout of their nets.




Figure 12. Completed Station Map (truncated)

Note that, in general, this procedure will create an approximately 20-hectare (50-acre) station if ten nets are dispersed over an 8-hectare (20-acre) core area. In the ideal case, the periphery of the 8-hectare core area would approximately be a square 280m on a side or a 160m radius circle, while the boundary of the 20-hectare station

would approximately be a square 450m on a side or a 250m radius circle. When plotted on the 30m grid map, such stations will encompass about 225 squares (a grid of 15 x 15 cells for a square station). The shape of your station, however, may not be a square or a circle. Nets, for example, might be located along a riparian corridor, within and around a montane meadow, or around part of an ox-bow lake. To allow you to map variously shaped stations, Form H0 depicts a 24 x 24 cell grid. In general, you should try to include between about 200 and 250 cells in your station. Even if your station is long and narrow, it should be drawn at least eight squares wide; a map using a grid of 9 x 24 cells might be ideal for such a case. Figure 12 presents an example of a completed station map based on an actual map submitted to us by a MAPS operator. Note that the nets are clustered somewhat more closely than normal for a typical MAPS station and, as a result, the total area of the station is only about 13 hectares (rather than 20 hectares).

Step 3: Define and delineate the habitats within the station. Habitat definitions should be based primarily on vegetation type and structure, and hydrology. Generally, anything smaller than about one ha (about 2.5 acres) in area - roughly equivalent to about ten or eleven 30m grid cells - should not be considered a separate habitat. Water features, such as lakes or rivers, should not be considered a separate habitat, regardless of size. Consider also the bird communities present in deciding whether to define and delineate separate habitats; different habitats should be reflected in some way in the bird communities they support. However, do not define your habitats too narrowly; if you wind up with more than five, you must reconsider your definitions. **Typically, stations have one, two or three habitat types; only rarely are there more**, so do not feel obliged to recognize five habitat types. Each habitat must be at least 5% of the station area. If not, it isn't considered a separate habitat for MAPS purposes.

Provide a concise name for each habitat type you identify within your station. In addition, use the National Vegetation Classification Standard (2021) provided in the MAPS HSA Code Supplement to identify each habitat type to the ALLIANCE level (see below under 'National Vegetation Classification Standard Alliance'). Please ensure that no two habitat types present within your station have the same alliance code – if they do, they are the same habitat. **NOTE: The classification system has been updated and habitat codes defined prior to the 2022 season will need to be re-determined. Call IBP if you need help with this.**

Clearly indicate the habitat delineations on your station map by drawing solid lines around them and shading the area if necessary. However, do not let your shading mask the clarity of net locations and other physical features.

Step 4: Complete the MAPS Habitat description forms. One form should be used for each habitat type (Form H1: Habitat Assessment Form). For each habitat type,

82 - 2025 MAPS Manual

consider only the portion of that habitat that falls within the station boundaries (area within 100 m from each net).

Cover: When describing habitat types, cover applies to the percentage of the station that is covered by that particular habitat type. When cover is used to describe upperstory, midstory, and understory vegetative layers, it is defined as the sum of the areas delimited by the vertical projections of plant perimeters onto the ground and not to light passing through the foliage of a single tree, shrub or fern. Consider any area inside the “drip-line” of the individual plant or tree as fully covered. Any overlap of cover between neighboring individual plants or trees is only considered once. In this case a completely closed canopy forest with overlapping crowns can only total 100% cover.

Habitat Structure Assessment form H1

We provide below a guide to filling in each field of those forms. We also provide, based on the example station map (Figure 12), a completed form H1 for the dominant forest habitat (Figure 13).

DATE: Indicate the month, day and year on which the habitat assessment is undertaken. We recommend you do this during the fourth visit to your station (not necessarily Period 4).

HABITAT DESIGNATION: The letter (A, B, C, D or E) should correspond to the labeling of the habitat types marked on your station map H0.

HABITAT NAME: The name should correspond to the labeling of the habitat types marked on your station map H0.

PERCENTAGE OF STATION COMPRISED OF THIS HABITAT TYPE: What percentage of the area of the station is occupied by the habitat in question? The sum of the percentages for each of the habitat types defined at the station (up to five) should generally be 100%. Small patches (<1 hectare, or <5% of the total area) of distinct habitat types must be lumped into the habitat type they appear to be associated with and described in the vegetative (or non-vegetative) layers. For instance, the presence of a few bushes around the edges of a meadow can be described in the shrub layer of the meadow habitat description and not treated as a separate habitat.

NATIONAL VEGETATION CLASSIFICATION STANDARD DIVISION: This is obtained from the National Vegetation Classification Standard (2021) which we have summarized in a separate document, the MAPS HSA Code Supplement: https://www.birdpop.org/docs/misc/HSA_code_supplement_2022.pdf. The supplement provides the background and instructions on how to use the hierarchical



Form H1: MAPS Habitat Structure Assessment (HSA) form

Location: HALF Station: HAFR Habitat Designation: A

Survey Date: 07 / 23 / 2025 Surveyed by: Jane Intern

Habitat Name (as indicated on station map H0): Sycamore, Tulip Poplar, Sweetgum Forest

Percentage of station comprised of this habitat type (from station map): 62 %

National Vegetation Classification Standard Division: I.B.3.Na

National Vegetation Classification Standard Alliance (optional): A3702

General description of habitat type. Include habitat age, major species, disturbance history, etc: Second growth woods on gently sloping hillside, on either side of a small riparian corridor containing a few greater than 100 year old Sycamore trees. Lies on well-drained (mostly) calcareous substrate. The lower portion of the station was clearcut in 1954 and they whole station was heavily flooded by a hurricane in 1983 but hasn't flooded since.

Vegetative Layers	% cover of station
Upperstory: >15m	80
Midstory: 5 -15m	40
Understory: 0.5 - 5m	40

Number of snags (>1m tall, >10cm diam.)

Circle one: 0-5 | 5-15 | >15

Average height of:

Tree canopy: 20 m

Shrubs: 2.0 m

Herbaceous vegetation: 1.2 m

Geographic Features	Options
	Circle one per line.
Drainage:	well-drained poorly-drained
Slope:	flat <u>gentle</u> undulating steep
Geography:	bottomland <u>hillside</u> ridgetop plain

Aquatic Features (if applicable):

Permanence

<u>seasonal</u>	permanent	occasional	other
-----------------	-----------	------------	-------

Features	% cover of station	If applicable, circle one or more aquatic features				
Running water	1	seep/trickle	very small brook (<0.5m)	<u>small stream (0.5-2.0m)</u>	large stream (2.0-5.0m)	river (>5m)
		canal				
Standing water	2	pond/lake <50m ² >50m ²	for livestock <50m ² >50m ²	<u>marsh/bog</u>		
				<50m ² <u>>50m²</u>		

Management/Disturbance history (if applicable):

Year(s)	Disturbance Type (circle one)
1945	fire wind flood drained icestorm logging: <u>clear-cut</u> strip selective logging
1983	fire wind <u>flood</u> drained icestorm logging: clear-cut strip selective logging
	fire wind flood drained icestorm logging: clear-cut strip selective logging

Figure 13. Completed MAPS HSA Form H1

classification system and provides a key to the NVCS codes (e.g., 1.B.2.Ne North American Great Plains Forest & Woodland Division); and provides a glossary of the terms used in the classification system.

NATIONAL VEGETATION CLASSIFICATION STANDARD ALLIANCE: The MAPS HSA Code Supplement:

https://www.birdpop.org/docs/misc/HSA_code_supplement_2022.pdf also provides regional lists of NVCS alliance level codes and one-line descriptions of the dominant species in those alliances. Use the NVCS Formation Code to limit the options, then choose the Alliance code that best describes the habitat being assessed. This space is for only the Alliance number (located in column H of the Excel database (e.g., A3725 or A3627). This code can be difficult to determine so it is optional.

GENERAL DESCRIPTION OF HABITAT TYPE INCLUDING HABITAT AGE: A description of the habitat type in your own words that includes major species, habitat age, disturbance history, etc. It is essential that you give a general description of the habitat type in the space provided. The following example was adapted from a station description provided by a MAPS station operator:

“Located along a small ridge between two valleys (approx. altitude in valleys is 150 m). This wooded area consists of oak (southern red predominates with some black and white), hickory, and poplar. Other species are Carolina buckthorn, ash, cherry, sweet gum, black gum, dogwood, etc. Vines are muscadine, poison ivy and rattan vine along with Japanese honeysuckle. Very little shrub undergrowth, mostly huckleberry/blueberry types.”

VEGETATIVE LAYERS PERCENT COVER: Divide the vegetation within each habitat into three main layers; upperstory, midstory, and understory, based solely on their height as described below, and **enter the percentage cover**. The cover percentage should represent the percentage cover of the layer **within** the habitat type and **not** the percentage cover of the area of the entire station.

For example, imagine a meadow that has shrubs dotted over it and reshape the area into a square, pushing the shrubs into one corner. Let us say, for instance, they cover approximately 7% of the entire area, then you would enter 7% onto the data sheet for the understory.

Upperstory: This vertical layer encompasses all vegetation above 15m from the ground, including coniferous or broad-leaved trees, vines, and epiphytic plants and lichens.

Midstory: This vertical layer encompasses all the vegetation between 5 and 15m above the ground, including saplings and tall shrubs as well as vines, epiphytic plants and lichens, and vegetation hanging down from the upperstory level.

Understory: This vertical layer includes vegetation found between 0.1 and 5m above the ground and includes mainly shrubs and small saplings. In addition, this layer may also contain herbaceous vegetation extending up from the ground cover layer.

AVERAGE HEIGHT OF TREE CANOPY: Enter the average height to the nearest 5m (or nearest 2m if the average height is less than 15m) of the tree canopy in the habitat type under consideration. Remember, trees are defined as woody, generally single-stemmed plants that are at least 5m tall at maturity. Do this regardless of whether the tree canopy extends into the upperstory, midstory, or only into the understory (see below). This can be achieved by drawing an imaginary line running through the tops the canopy whereby the area of the outlines of trees above the line equals the area of the gaps below it (*ignore gaps that reach the ground*). Estimates obtained using a clinometer are preferred but we understand that few stations have one. If you are working in a managed forest or woodland it is likely that the responsible land manager(s) can provide this information.

AVERAGE HEIGHT OF SHRUBS: Enter the average height to the nearest 0.5m of the shrubs (woody, generally multi-stemmed plants with a bushy appearance) in the habitat type under consideration. Note that when considering the understory vegetative layer (below), you will estimate the cover and pattern of all vegetation between 0.5 and 5m, regardless of whether it is comprised of trees (except trunks), shrubs, or herbaceous vegetation. Here, we are asking only for the average height of the shrubs.

AVERAGE HEIGHT OF HERBACEOUS VEGETATION: Enter the average height to the nearest 0.1m of the herbaceous (non-woody, vascular) vegetation, which includes graminoids (grass-like vegetation including grasses, sedges, rushes, etc.), forbs (broad-leaved herbaceous vegetation), ferns, and or non-vascular vegetation (mosses and lichens).

NUMBER OF SNAGS: Circle the number (0, <5, 5-15, >15) that best represents the number of snags present in the habitat type under consideration. Snags are defined as dead woody stems greater than 1m in height and greater than 10cm in diameter.

GEOGRAPHIC FEATURES: Circle the features present within this habitat type.

Drainage – circle one option (well-drained or poorly-drained) that best represents the drainage characteristics of the habitat.

Slope – circle one option that best represents the topography of the habitat: flat, gently sloping, undulating or steep. This is a judgment call but not a difficult one.

Geography – indicate if the habitat is associated with a bottomland, hillside, ridgetop or plain.

AQUATIC FEATURES: Enter the percent cover of these features within each habitat type. Please note that these features should be indicated clearly on your station map. Let us consider the types of features individually:

Permanence: whether the water body remains present throughout the season or not. Such water bodies may be permanent, seasonal (vernal pools) or occasional (flooded field).

Running water: Running water courses range in width from a seep/trickle, to a very small brook (<0.5m), a small stream (0.5-2.0m), a large stream (2.0-5.0m), a river (>5.0m) or canal.

Standing water: Standing water includes ponds and lakes (<50m² or >50m²), water catchment for livestock (<50m² or >50m²), or marsh/bog areas (<50m² or >50m²).

For example, if ponds occur within a meadow and normally have water in them year-round, then you would circle natural pond, permanent and the size category <50m² or >50m² (if both size classes occur circle both). If there is standing water at your station that does not fit any of these categories, circle other and write a brief description in the habitat description section.

MANAGEMENT/DISTURBANCE HISTORY: Circle the appropriate option(s) describing the type of management practice or disturbance, if any apply (e.g., Year(s): 1945-55 logging: clear-cut)). The logging patterns may be indicated by circling clear cut (of at least 1 hectare), selective (small patches <1 hectare in extent, or individual trees), or a strip of trees (removed for a powerline, for example).

A description of the management history should be included in the general description of the habitat above (e.g., I estimate that the area was probably logged in the late 1940's/early 1950's.) especially if the disturbance is not listed as one of the choices to be circled.

DATA SUBMISSION

Making sure the required data from each station become a part of the compiled MAPS database is the final and crucial step in operating a MAPS station. To maximize the use that can be made of the data, all elements listed below must be included. It is also important to ensure that data submission occurs within a reasonable amount of time; delays hold up analyses, prevent us from providing you with timely feedback, and require us to spend time rounding up outstanding data.

What data to submit

Each year, for each location, MAPS operators must submit the following data:

- Banding data for newly banded birds
- Banding data for recaptured birds
- Banding data for unbanded birds
- Summary of mist-netting effort data for each station
- Breeding status data for each station

Habitat Structure Assessment (HSA) data (including the station map) must also be submitted for each station during its first year of operation and every five years following (i.e., sixth year, eleventh year, etc.). However, if substantial habitat change has occurred within five years as a result of rapid succession or catastrophic events (anthropogenic or natural), a revised HSA, along with the creation of a revised station map, should be completed. A revised station map should also be submitted anytime that nets are moved. Be sure to show the locations of the old, as well as the new, nets. Refer to the HSA Protocol (Nott *et al.* 2003; which can be downloaded from the IBP website) for directions on how to create the station map.

In addition, submit a completed Standard Net Opening and Closing Times sheet for each station at the end of the first season of operation and after any season in which any of the standard operating times are changed, that is, whenever you have changed any of the times at which you plan to operate your station in the future.

How to submit MAPS data

Currently, data may be submitted to the MAPS program in two ways: electronically using MAPSNet or non-electronically by submitting hard (paper) copies of all data.

Submitting data through MAPSNet: Coinciding with the Bird Banding Offices' effort to institute electronic data submission, IBP has developed a cloud-based data verification/editing, and error-tracking program called MAPSNet for submitting MAPS data to IBP. We strongly encourage all operators to submit their 2025 data using MAPSNet, which was introduced on pg. 20. MAPSNet includes modules to verify your banding, effort, breeding status, and habitat structure assessment data collected during the 2025 MAPS season. MAPSNet is designed to mimic the data verification procedures that have been developed by IBP over more than 30 years and have been applied to

every set of MAPS data contributed to the MAPS program. The checks embedded in the program will allow you to see and correct any errors or inconsistencies that occur in your own data and will help you to improve your data collection. Moreover, submission of data through MAPSNet will, in the long run, reduce the amount of time IBP biologists must spend in verifying data from the over 300 MAPS stations operated each year, thus allowing them more time and resources to focus on analyses aimed at understanding the causes of population declines in landbirds and at formulating management and conservation strategies for them.

Information on how to use MAPSNet is available on the MAPSNet web page <http://www.birdpop.org/pages/mapsMAPSNet.php>.

For stations that have run for more than one-year, proper use of MAPSNet requires that recapture records from the current year be checked against banding data from previous years in order to correct discrepancies among recaptures and to screen recaptures for possible misread band numbers. So, IBP must pull past years of data into MAPSNet before it is accessible for the current year.

It is extremely important to proof the file you wish to import against your raw data sheets before using MAPSNet to verify it. Please email the MAPS Data Manager at mapsdatamanager@birdpop.org if you have questions regarding the use of MAPSNet.

The Bird Banding Offices is now requiring the use of their Bander Portal banders. As of the release of this manual (April 2025), MAPSNet creates a file that easily imports into the Bander Portal, including the conversion of codes from MAPS codes to BBL codes. Additionally, in the few cases where the species alpha codes in Pyle (2022) differ from those used by the BBL, the alpha codes will be converted to BBL codes in the export.dbf file.

Following are explanations of the fields listed in Table 4; for further details on the codes used, consult the "Collection and Recording of Banding Data" section of this manual. All character fields should be entered left-justified and numeric fields right-justified.

- LOC - Location. Enter your four-character location code.
- BI - Bander's Initials. Enter the two-character bander's initials.
- BS - Band size. The purpose of this and the following field is to enable us to find original data easily. Records on 'Unbanded' and 'Recapture' sheets should be entered with band sizes 'U' and 'R', respectively. Unbanded birds on new-band sheets should be given the band size for the sheet
- PG - Page number.
- C - Capture code.
- BAND - Band number (always nine digits long).
- SPEC - Four-letter species alpha code.
- AGE - Age.
- HA - How aged.
- WRP - Wolfe-Ryder-Pyle plumage and molt code.
- SEX - Sex.
- HS - How sexed.

90 - 2025 MAPS Manual

- SK - Skull.
- CP - Cloacal protuberance.
- BP - Brood patch.
- F - Fat.
- BM - Body molt.
- FM - Flight-feather molt.
- FW - Flight-feather wear.
- JP - Juvenile body plumage.
- WNG - Wing chord.
- WEIGHT - Body mass.
- STATUS - Status.
- DATE - In dBase, enter as MM/DD/YYYY. In a text file, enter as YYYYMMDD.
- TIME - Omit the final '0'.
- STATION - Station code.
- NET - Original net designation. Enter your net number (preferably two digits) or "?" if net number is unknown, left justified
- DISP - Disposition.
- NOTE - Enter the note number if the record has a note. Otherwise, leave blank.
- PPC - Primary coverts.
- SSC - Secondary coverts.
- PPF - Primaries.
- SSF - Secondaries.
- TT - Tertiaries.
- RR - Rectrices.
- BPL - Includes all feather tracts of the head, upperparts and underparts (this field is to be used for MAPS data in and subsequent to 2004).
- NF - all non-feather parts including bill, mouth, eye, legs, and feet. A note is required if this column is used.
- TAIL - Tail length. Not required for MAPS
- TARSUS - Tarsus length.
- NOTES - Enter a brief note.
- FTHR. PULL - Enter 'O' if the outer two rectrices were pulled or 'I' if an inner and outer rectrix were pulled. If no feathers were pulled, leave this field blank.

Before submitting electronic banding data, it is extremely important that you proof your electronic file against the banding-data sheets for data-entry errors.

Submitting paper copies of banding data: Operators who are unable to use MAPSNet and are unable to submit electronic data must submit copies (photocopies or PDF scans, or photos, including notes, are acceptable) of all of the forms mentioned above, **including completed Banding Data Sheets for newly banded birds, unbanded birds, and recaptured birds; completed Summary of Mist-Netting Effort forms for each station, completed Summary of Mist-Netting Results forms for each station, and completed Breeding Status Lists for each station each year.** In addition, such operators must submit paper copies of their completed HSA forms and the associated station map for each station (once every five years or more often if substantial habitat change has

occurred), and their Standard Net Opening and Closing Times form (after the first season and whenever they change their standard operation). Please do not staple data sheets together or put them in binders when submitting data.

Due date

MAPS operators are requested to return their completed data and map(s) to The Institute for Bird Populations as soon as possible after the completion of the season. In general, the due date is September 15. The due date for operators using MAPSNet is October 15. Data will be accepted after these dates, but late data, especially from long-standing stations, compromise our ability to conduct analyses and prepare reports on schedule. However, we would rather receive complete, proofed, carefully-compiled data packets a little late than incomplete packets submitted on schedule.

Hard copy data packets should be addressed to: MAPS Data Manager, The Institute for Bird Populations. Our mailing address is PO Box 518, Petaluma, CA 94953 USA.

We will acknowledge receipt of your data; if you do not hear from us within a month of sending your data, chances are we did not receive them!

Table 4. MAPS Banding-data file structure for 2025 data

<u>Field</u>	<u>Field Name</u>	<u>Type</u>	<u>Width</u>	<u>Dec</u>	<u>Description</u>
1	LOC	Character	4		Location code
2	BI	Character	2		Bander's initials
3	BS	Character	2		Band size
4	PG	Character	3		Data page number
5	C	Character	1		Capture code
6	BAND	Character	9		Band number
7	SPEC	Character	4		Four-letter species alpha code
8	AGE	Character	1		Age
9	HA	Character	2		How aged
10	WRP	Character	4		Wolfe-Ryder-Pyle plumage code
11	SEX	Character	1		Sex
12	HS	Character	2		How sexed
13	SK	Character	1		Skull pneumatization
14	CP	Character	1		Cloacal protuberance score
15	BP	Character	1		Brood patch score
16	F	Character	1		Fat content score
17	BM	Character	1		Body molt score
18	FM	Character	1		Flight feather molt score
19	FW	Character	1		Flight feather wear score
20	JP	Character	1		Juvenile plumage score
21	WNG	Numeric	3	0	Wing chord
22	WEIGHT	Numeric	5	1	Body mass
23	STATUS	Character	3		Status upon release
24	DATE	Date	8		Capture date
25	TIME	Character	3		Time of capture
26	STATION	Character	4		Station code
27	NET	Character	4		Net

92 - 2025 MAPS Manual

28	DISP	Character	1		Disposition on release
29	NOTE	Character	2		Notes on data sheet
30	PPC	Character	1		Feather generations in primary coverts
31	SSC	Character	1		Feather generations in secondary coverts
32	PPF	Character	1		Feather generations in primaries
33	SSF	Character	1		Feather generations in secondaries
34	TT	Character	1		Feather generations in tertials
35	RR	Character	1		Feather generations in rectrices
36	BPL	Character	1		Feather generations in body plumage
37	NF	Character	1		Generation indicated by non-feather parts
38	FP	Character	1		Feather pull status
39	TAIL	Numeric	3	0	Tail length measurement
40	TARSUS	Numeric	5	1	Tarsus length measurement
41	NOTES	Character	50		Notes taken in the field

LITERATURE CITED

- American Ornithologists' Union. 1957. *Check-list of North American Birds*. 5th edition. American Ornithologists' Union, Washington, DC.
- American Ornithologists' Union. 1983. *Check-list of North American Birds*. 6th edition. American Ornithologists' Union, Washington, DC.
- American Ornithologists' Union. 1998. *Check-list of North American Birds*. 7th edition. American Ornithologists' Union, Washington, DC.
- Anders, A.D., D.C. Dearborn, J. Faaborg, and F.R. Thompson III. 1997. Juvenile survival in a population of neotropical migrant birds. *Conservation Biology* 11:698-707.
- Baillie, S.R. 1990. Integrated population monitoring of breeding birds in Britain and Ireland. *Ibis* 132:151-166.
- Baillie, S.R., R.E. Green, M. Boddy, and S.T. Buckland. 1986. *An Evaluation of the Constant Effort Sites Scheme*. British Trust for Ornithology, Thetford, UK.
- Bird Banding Offices. 1991. *North American Bird Banding Manual, Vol. II*. U.S. Fish and Wildlife Service, Washington, DC, and Canadian Wildlife Service, Ottawa, ON.
- Brown, L.R. 1991. *State of the World, 1992*. Worldwatch Institute, Washington, DC.
- Burton, K.M., and D.F. DeSante. 2004. Effects of mist-netting frequency on capture rates at monitoring Avian Productivity and Survivorship (MAPS) stations. Pp. 7-11 in Ralph, C.J., and E.H. Dunn, eds. *Monitoring Bird Populations Using Mist Nets*. *Studies in Avian Biology* 29.
- Butcher, G., and S. Droege. 1992. *Needs Assessment: Monitoring Neotropical Migratory Birds*. Neotropical Migratory Birds Conservation Program. 58 pp.
- Chesser, R.T., S.M. Billerman, K.J. Burns, C. Cicero, J.L. Dunn, B. E. Hernández-Baños, R.A. Jiménez A.W. Kratter, N.A. Mason, P.C. Rasmussen, J. V. Remsen, Jr, K. Winker. 2023. Sixty-fourth Supplement to the American Ornithological Society's Check-list of North American Birds. *Ornithology* 140(3)
- Clobert, J., J.-D. Lebreton, and D. Allaine. 1987. A general approach to survival rate estimation by recaptures or resightings of marked birds. *Ardea* 75:133-142.
- DeSante, D.F. 1992. Monitoring Avian Productivity and Survivorship (MAPS): a sharp, rather than blunt, tool for monitoring and assessing landbird populations. Pp. 511-521 in McCullough, D.C., and R.H. Barrett, eds. *Wildlife 2001: Populations*. Elsevier Applied Science, London, UK.
- DeSante, D.F. 1995. Suggestions for future directions for studies of marked migratory landbirds from the perspective of a practitioner in population management and conservation. *Journal Applied Statistics* 22:949-965.
- DeSante, D.F. 1996. General Evaluation of the Monitoring Avian Productivity and Survivorship (MAPS) Program in: *An Evaluation of the Monitoring Avian Productivity and Survivorship (MAPS) Program*. The Institute for Bird Populations, Point Reyes Station, CA. 128 pp.
- DeSante, D.F. 2000. Patterns of productivity and survivorship from the MAPS Program. Pp. 166-177 in Bonney, R., D.N. Pashley, R.J. Cooper, and L. Niles, eds. *Strategies for Bird Conservation: the Partners in Flight Planning Process*. Proceedings of the Third

- Partners in Flight Workshop; 1995 October 1-5; Cape May, NJ. Proceedings RMRS-P-16. Ogden, UT: USDA, Forest Service, Rocky Mountains Research Station.
- DeSante, D.F., K.M. Burton, J.F. Saracco, and B.L. Walker. 1995. Productivity indices and survival rate estimates from MAPS, a continent-wide programme of constant-effort mist netting in North America. *Journal Applied Statistics* 22:935-947.
- DeSante, D.F., K.M. Burton, and D.R. O'Grady. 1996. The Monitoring Avian and Productivity and Survivorship (MAPS) Program fourth and fifth annual report (1993 and 1994). *Bird Populations* 3:67-120.
- DeSante, D.F., and D.R. Kaschube. 2006. The Monitoring Avian Productivity and Survivorship (MAPS) program 1999, 2000, and 2001 report. *Bird Populations* 7:23-89.
- DeSante, D.F., and D.R. Kaschube. 2007. The Monitoring Avian Productivity and Survivorship (MAPS) program 2002 and 2003 report. *Bird Populations* 8:46-115.
- DeSante, D.F., and D.R. Kaschube. 2009. The Monitoring Avian Productivity and Survivorship (MAPS) program 2004, 2005, and 2006 report. *Bird Populations* 9:86-169.
- DeSante, D. F., D. R. Kaschube, and J. F. Saracco. 2015. Vital Rates of North American Landbirds. www.VitalRatesOfNorthAmericanLandbirds.org. The Institute for Bird Populations.
- DeSante, D.F., D.R. Kaschube, and T.S. Sillett. 2003a. *Evaluation of the first ten years of the Monitoring Avian Productivity and Survivorship (MAPS) program in Alaska and adjacent Canada, Part 1. General results*. The Institute for Bird Populations, Point Reyes Station, CA. 226 pp.
- DeSante, D.F., D.R. Kaschube, and T.S. Sillett. 2003b. *Evaluation of the first ten years of the Monitoring Avian Productivity and Survivorship (MAPS) program in Alaska and adjacent Canada, Part 2. Statistical power to detect temporal trends and spatial differences in survival of landbirds*. The Institute for Bird Populations, Point Reyes Station, CA. 50 pp.
- DeSante, D.F., D.R. Kaschube, J.F. Saracco, and J.E. Hines. 2009. Power to detect differences and trends in apparent survival rates. *Bird Populations* 9:29-41.
- DeSante, D.F., M.P. Nott, and D.R. O'Grady. 2001. Identifying the proximate demographic cause(s) of population change by modelling spatial variation in productivity, survivorship, and population trends. *Ardea* 89 (special issue): 185-207.
- DeSante, D.F., M.P. Nott, and D.R. Kaschube. 2004b. *MAPS stations on National Wildlife Refuges in the USFWS Pacific Region*. The Institute for Bird Populations, Point Reyes Station, CA. 57 pp.
- DeSante, D.F., M.P. Nott, and D.R. Kaschube. 2005. Monitoring, modeling, and management: Why base avian management on vital rates and how should it be done? Pp. 795-804 in Ralph, C.J., and T.D. Rich, eds. *Bird Conservation Implementation and Integration in the Americas*. USDA Forest Service Gen. Tech. Rep. PSW-191.
- DeSante, D.F., and D.R. O'Grady. 2000. The Monitoring Avian Productivity and Survivorship (MAPS) Program 1997 and 1998 report. *Bird Populations* 5: 49-101.
- DeSante, D.F., D.R. O'Grady, K.M. Burton, P. Velez, D. Froehlich, E.E. Feuss, H. Smith, and E.D. Ruhlen. 1998. The Monitoring Avian Productivity and Survivorship (MAPS) Program Sixth and Seventh Annual Report (1995 and 1996). *Bird Populations* 4:69-122.

- DeSante, D.F., D.R. O'Grady, and P. Pyle. 1999. Measures of productivity and survival derived from standardized mist-netting are consistent with observed population trends. *Bird Study* 46(suppl):S178-188.
- DeSante, D., and P. Pyle. 1986. *Distributional Checklist of North American Birds*. Volume 1: United States and Canada. Artemisia Press, Lee Vining, CA.
- DeSante, D.F., and D.K. Rosenberg. 1998. What do we need to monitor in order to manage landbirds? Pp. 93-106 in Marzluff, J.M., and R. Sallabanks, eds. *Avian Conservation: Research and Management*. Island Press, Washington, DC.
- DeSante, D.F., and J.F. Saracco. 2009. Power of the MAPS program to detect differences and trends in survival and a vision for program growth. *Bird Populations* 9:42-75.
- DeSante, D.F., J.F. Saracco, D.R. O'Grady, K.M. Burton, and B.L. Walker. 2004a. Methodological considerations of the Monitoring Avian Productivity and Survivorship (MAPS) program. Pp. 28-45 in Ralph, C.J., and E.H. Dunn, eds. *Monitoring Bird Populations Using Mist Nets*. *Studies in Avian Biology* 29.
- DeSante, D.F., J.F. Saracco, P. Pyle, D.R. Kaschube, and M.K. Chambers. 2008. *Integrating the MAPS program into Coordinated Bird Monitoring in the Northeast (U. S. Fish and Wildlife Service Region 5)*. The Institute for Bird Populations, Point Reyes Station, CA. 99 pp.
- DeSante, D.F., O.E. Williams, and K.M. Burton. 1993. The Monitoring Avian Productivity and Survivorship (MAPS) Program: overview and progress. Pp. 208-222 in Finch, D.M., and P.W. Stangel, eds. *Status and Management of Neotropical Migratory Birds*. USDA Forest Service Gen. Tech. Rep. RM-229.
- Froehlich, D. 2003. *Ageing North American Landbirds by Molt Limits and Plumage Criteria: a Photographic Companion to the Identification Guide to North American Birds, Part I*. Slate Creek Press, Bolinas, CA. 51 pp
- Geissler, P. 1997. Review of the Monitoring Avian Productivity and Survivorship (MAPS) Program in: *An Evaluation of the Monitoring Avian Productivity and Survivorship (MAPS) Program*. The Institute for Bird Populations, Pt Reyes Sta, CA.
- George, T.L., R.J. Harrigan, J.A. LaManna, D.F. DeSante, J.F. Saracco, and T.B. Smith. 2015. Persistent impacts of West Nile virus on North American bird populations. *Proceedings of the National Academy of Sciences* 112(46):14290–14294.
- Glass, A. J., A.J. Caven, D. Kim, M.O. Sutton, and N. Arcilla. 2020. Climate change and land management implications for a declining Neotropical migratory songbird breeding in the North American Great Plains. *Avian Conservation and Ecology* 15(1):4.
- Hines, J.E., W.L. Kendall, and J.D. Nichols. 2003. On the use of the robust design with transient capture-recapture models. *Auk* 120:1151-1158.
- Holmes, R.T., and T.W. Sherry. 1988. Assessing population trends of New Hampshire forest birds: local vs. regional trends. *Auk* 105:756-768.
- Howell, S.N.G., C. Corbin, P. Pyle, and D.I. Rogers. 2003. The first basic problem: a review of molt and plumage homologies. *Condor* 105:635-653.
- Hutto, R.L. 1988. Is tropical deforestation responsible for the reported decline in neotropical migrant populations? *American Birds* 42:375-379.
- The Institute for Bird Populations. 2023. MAPS Data Exploration Tool: Processed MAPS results. Version 1.0. <https://ibp-maps-data-exploration-tool.org>

- Johnson, E.I., J.D. Wolfe, T.B. Ryder, and P. Pyle. 2011. Modifications to a molt-based ageing system proposed by Wolfe et al. (2010). *Journal of Field Ornithology* 82:422-424.
- Kaschube, D.R., J.F. Saracco, C. Ray, C.M. Godwin, K.R. Foster, and P. Pyle. 2022. Minimum capture-recapture rates and years of banding station operations to obtain reliable adult annual survival estimates. *Journal of Field Ornithology* 93:7.
- Knopf, F.L. 1994. Avian assemblages on altered grasslands. *Studies in Avian Biology* 15:247-257.
- Lebreton, J.-D, K.P. Burnham, J. Clobert, and D.R. Anderson. 1992. Modeling survival and testing biological hypotheses using marked animals: a unified approach with case studies. *Ecol. Monogr.* 62:67-118.
- Morton, E.S., and R. Greenberg. 1989. The outlook for migratory songbirds: "future shock" for birders. *American Birds* 43:178-183.
- National Geographic Society. 1999. *Field Guide to the Birds of North America. 3rd edition.* National Geographic Society, Washington, DC.
- Ng, A.M.B., M. Pontius, S.L. De Ruiter, and D.S. Proppe. 2020. Noise, avian abundance, and productivity at banding stations across the Continental United States. *Avian Conservation and Ecology* 15(2):4.
- Nott, M.P. 2000. *Identifying management actions on DoD installations to reverse declines in Neotropical birds.* The Institute for Bird Populations, Pt. Reyes Sta., CA. 18 pp.
- Nott, M.P., and D.F. DeSante. 2002. Demographic monitoring and the identification of transients in mark-recapture models. Pp. 727-736 in J.M. Scott, P.J. Heglund, M.L. Morrison, et al., (eds.), *Predicting Species Occurrences: Issues of Scale and Accuracy.* Island Press, New York
- Nott, M.P., D.F. DeSante, R.B. Siegel, and P. Pyle. 2002. Influences of the El Niño/Southern Oscillation and the North Atlantic Oscillation on avian productivity in forests of the Pacific Northwest of North America. *Global Ecology and Biogeography* 11:333-342.
- Nott, M.P., D.F. DeSante, and N. Michel. 2003a. *Management Strategies for Reversing Declines in Landbirds of Conservation Concern: a Landscape-Scale Analysis of MAPS Data.* The Institute for Bird Populations, Point Reyes Station, CA 357 pp.
- Nott, M.P., D.F. DeSante, and N. Michel. 2003b. *Monitoring Avian Productivity and Survivorship (MAPS) Habitat Structure Assessment (HSA) Protocol.* The Institute for Bird Populations, Point Reyes Station, CA. 42 pp.
- Nott, M.P., D.F. DeSante, P. Pyle, and N. Michel. 2005. *Managing Landbird Populations in Forests of the Pacific Northwest: Formulating Population Management Guidelines from Landscape-scale Ecological Analyses of MAPS Data from Avian Communities on Seven National Forests in the Pacific Northwest.* The Institute for Bird Populations, Point Reyes Station, CA 163 pp. + appendices.
- Peach, W., S. Baillie, and L. Underhill. 1991. Survival of British Sedge Warblers *Acrocephalus schoenobaenus* in relation to west African rainfall. *Ibis* 133:300-305.
- Peach, W.J., S.T. Buckland, and S.R. Baillie. 1990. Estimating survival rates using mark-recapture data from multiple ringing sites. *The Ring* 13:87-102.
- Peach, W.J., S.T. Buckland, and S.R. Baillie. 1996. The use of constant effort mist-netting to measure between-year changes in the abundance and productivity of common passerines. *Bird Study* 43:142-156.

- Peterjohn, B.G., J.R. Sauer, and C.S. Robbins. 1995. Population trends from the North American Breeding Bird Survey. Pp. 3-39 in T.E. Martin and D.M. Finch (eds.), *Ecology and Management of Neotropical Migratory Birds*. Oxford University Press, New York, NY.
- Peterson, R.T. 1980. *Peterson Field Guide: Eastern Birds*. 4th edition. Houghton Mifflin Comp., Boston, MA.
- Peterson, R.T. 1990. *Peterson Field Guide: Western Birds*. 3rd edition. Houghton Mifflin Comp., Boston, MA.
- Pollock, K.H., J.-D. Nichols, C. Brownie, and J.E. Hines. 1990. Statistical inference for capture-recapture experiments. *Wildlife Monographs* 107.
- Pradel, R. 1996. Utilization of capture-mark-recapture for the study of recruitment and population growth rate. *Biometrics* 52:703-709.
- Pradel, R., J.E. Hines, J.-D. Lebreton, and J.D. Nichols. 1997. Capture-recapture survival models taking account of transients. *Biometrics* 53:60-72.
- Pyle, P. 2022. *Identification Guide to North American Birds*, Part 1. Second Edition. Slate Creek Press, Bolinas, CA.
- Pyle, P. 2004. New terminology will help you understand molt. *MAPS Chat* 7:2,8.
- Pyle, P., and D.F. DeSante. 2003. Four-letter and six-letter alpha codes for birds recorded from the American Ornithologist's Union check-list area. *North American Bird Bander* 28:64-79.
- Pyle, P., and D.F. DeSante. 2005. Updates to four-letter and six-letter alpha codes based on Revisions by the American Ornithologists' Union. *North American Bird Bander* 30:70-72.
- Pyle, P., and D.F. DeSante. 2006. Updates to four-letter and six-letter alpha codes based on Revisions by the American Ornithologists' Union. *North American Bird Bander* 31:193-195.
- Pyle, P., D.F. DeSante, M.P. Nott, and D.R. Kaschube. 2005. *The MAPS program in the Pacific Northwest: Current status and future directions*. The Institute for Bird Populations, Point Reyes Station, CA. 106 pp.
- Pyle P, K.R. Foster, G.M Godwin, D.R. Kaschube, and J.F. Saracco. 2020. Yearling proportion correlates with habitat structure in a boreal forest landbird community. *PeerJ* 8:e8898
- Pyle, P., M. Gahbauer, E.I. Johnson, T.B. Ryder, and J.D. Wolfe. 2021. Application of a global age-coding system ("WRP"), based on molts and plumages, for use in demographic and other studies of birds. *Ornithology* 139(1):1-12. URL: <https://doi.org/10.1093/ornithology/ukab063>
- Pyle, P., J.F. Saracco, and D.F. DeSante. 2018. Evidence of widespread movements from breeding to molting grounds by North American landbirds. *The Auk: Ornithological Advances* 135:506-520.
- Ralph, C.J., G.R. Geupel, P. Pyle, T.E. Martin, and D.F. DeSante. 1993. *Handbook of Field Methods for Monitoring Landbirds*. USDA Forest Service Gen. Tech. Report PSW-GTR-144.

- Robbins, C.S., J.R. Sauer, R.S. Greenberg, and S. Droege. 1989. Population declines in North American birds that migrate to the neotropics. *Proc. Nat. Acad. Sci. (USA)* 86:7658-7662.
- Robinson, R.A., R. Julliard, and J.F. Saracco. 2009. Constant effort: studying avian population processes using standardised ringing. *Ringing & Migration* 24:199-204.
- Rodewald, P. (Editor). 2015. The Birds of North America Online: <http://bna.birds.cornell.edu/BNA/>. Cornell Laboratory of Ornithology, Ithaca, NY.
- Rosenberg, D.K., D.F. DeSante, K.S. McKelvey, and J.E. Hines. 1999. Monitoring survival rates of Swainson's Thrush *Catharus ustulatus* at multiple spatial scales. *Bird Study* 46(suppl): S198-208.
- Ruegg, K., E. Anderson, K. Paxton, V. Apkenas, S. Lao, R.B. Siegel, D.F. DeSante, F. Moore, and T. Smith. 2014. Mapping migration in a songbird using high-resolution genetic markers. *Molecular Ecology* 23:5726–5739.
- Ruegg, K.C., R.J. Harrigan, J.F. Saracco, T.B. Smith, and C.M. Taylor. 2020. A genoscape-network model for conservation prioritization in a migratory bird. *Conservation Biology* 34 (6): 1482-1491.
- Rushing, C.S., T.B. Ryder, A.L. Scarpignato, J.F. Saracco, and P.P. Marra. 2016. Using demographic attributes from long-term monitoring data to delineate population structure. *The Journal of Applied Ecology*. Early View published online at <http://onlinelibrary.wiley.com/doi/10.1111/1365-2664.12579/abstract>.
- Saracco, J. 2004. Changes in the use of the "adults only" (now called "molt limits and plumage" fields: a more intuitive and informative approach to ageing birds. *MAPS Chat* 7:3-5.
- Saracco, J.F., and D.F. DeSante. 2008. *Identifying proximate causes of population trends in migratory birds*. The Institute for Bird Populations, Point Reyes Station, CA. 44 pp.
- Saracco, J.F., and M. Rubenstein. 2020. Integrating broad-scale data to assess demographic and climatic contributions to population change in a declining songbird. *Ecology and Evolution* 2020;10:1804-1816.
- Saracco, J.F., D.F. DeSante, D.R. Kaschube, J.E. Hines, M.P. Nott, and R.B. Siegel. 2006. *An analysis of the Monitoring Avian Productivity and Survivorship (MAPS) program and a vision for its integration into North American Coordinated Bird Monitoring*. The Institute for Bird Populations, Point Reyes Station, CA. 87 pp.
- Saracco, J.F., D.F. DeSante, and D.R. Kaschube. 2008. Assessing landbird monitoring programs and demographic causes of population trends. *Journal of Wildlife Management* 72:1665-1673.
- Saracco, J.F., D.F. DeSante, M.P. Nott, W.M. Hochachka, S Kelling, and D. Fink. 2009a. Integrated bird monitoring and the Avian Knowledge Network: Using multiple data resources to understand spatio-temporal variation in demographic processes and abundance. Pp. 659-661 in Rich, T.D., C. Arizmendi, D.W. Demarest, and C. Thompson (eds.), *Proceedings of the 4th International Partners in Flight Conference: Tundra to Tropics*. 13-16 February 2008. McAllen, Texas. Partners in Flight.
- Saracco, J.F., D.F. DeSante, M.P. Nott, and D.R. Kaschube. 2009b. Using MAPS and MoSI to monitor landbirds and inform conservation. Pp. 651-658 in Rich, T.D., C. Arizmendi, D.W. Demarest, and C. Thompson (eds.), *Proceedings of the 4th International Partners*

- in Flight Conference: Tundra to Tropics*. 13-16 February 2008. McAllen, Texas. Partners in Flight.
- Saracco, J.F., J.A. Royle, D.F. DeSante, and B. Gardner. 2010. Modeling spatial variation in avian survival and residency probabilities. *Ecology* 91(7):1885-1891.
- Saracco, J.F., J.A. Royle, D.F. DeSante, and B. Gardner. 2012. Spatial modeling of survival and residency and application to the Monitoring Avian Productivity and Survivorship program. *Journal of Ornithology* 152 (Supplement 2):S469-S476
- Saracco, J.F., R.B. Siegel, L. Helton, S. Stock, and D. DeSante. 2019. Phenology and productivity in a montane bird assemblage: trends and responses to elevation and climate variation. *Global Change Biology* 2019:25:985-086.
- Sibley, D.A. 2003. *The Sibley Field Guide to Birds of Eastern North America*. Alfred A. Knopf, New York.
- Sibley, D.A. 2003. *The Sibley Field Guide to Birds of Western North America*. Alfred A. Knopf, New York.
- Temple, S.A., and J.A. Wiens. 1989. Bird populations and environmental changes: can birds be bio-indicators? *American Birds* 43:260-270.
- Terborgh, J. 1989. *Where Have All the Birds Gone? Essays on the Biology and Conservation of Birds that Migrate to the American Tropics*. Princeton University Press, Princeton, NJ.
- Vega Rivera, J.H., J.H. Rappole, W.J. McShea, and C.A. Haas. 1998. Wood Thrush postfledging movements and habitat use in northern Virginia. *Condor* 100:69-78.
- USNVC [United States National Vegetation Classification]. 2021. United States National Vegetation Classification Database, V2.031. Federal Geographic Data Committee, Vegetation Subcommittee, Washington DC. [usnvc.org] (accessed 04/14/2022)
- Wolfe, J.D., T.B. Ryder, and P. Pyle. 2010. Using molt cycles to categorize the age of tropical birds: an integrative new system. *Journal of Field Ornithology* 81:186-194.
- White, G.C. 1983. Numerical estimation of survival rates from band-recovery and biotelemetry data. *J. Wildlife Management* 47:716-728.
- White, G.C., and K.P. Burnham. 1999. Program Mark: survival estimation from populations of marked animals. *Bird Study* 46(suppl.):120-138.
- Wilcove, D.S. 1985. Nest predation in forest tracts and the decline of migratory songbirds. *Ecology* 66:1211-1212.
- Youngflesh, C., G.A. Montgomery, J.F. Saracco, D.A.W. Millerd, R.P. Guralnicke, A.H. Hurlbert, R.B. Siegel, R. LaFrancee, and M.W. Tingley. 2023. Demographic consequences of phenological asynchrony for North American songbirds. *Proceedings of the National Academy of Sciences* 28:e2221961120.
- Youngflesh, C., J.F. Saracco, R.B. Siegel, and M.W. Tingley. 2022. Abiotic conditions shape spatial and temporal morphological variation in North American birds. *Nature Ecology and Evolution*. DOI 10.1038/s41559-022-01893-x.
- Yunick, R.P. 1979. Variation in skull pneumatization patterns of certain passerines. *North American Bird Bander* 4:145-147.