

**THE 2007 REPORT OF THE
MONITORING AVIAN PRODUCTIVITY AND SURVIVORSHIP
(MAPS) PROGRAM ON FORT BRAGG**

Phil Nott, Peter Pyle, and Danielle Kaschube

**THE INSTITUTE FOR BIRD POPULATIONS
P.O. Box 1346
Point Reyes Station, CA 94956-1346**

(415) 663-1436

pnott@birdpop.org

March 12, 2008

Introduction

Since 1989, The Institute for Bird Populations has been coordinating the Monitoring Avian Productivity and Survivorship (MAPS) Program, a cooperative effort among public and private agencies and individual bird banders in North America, to operate a continent-wide network of over 500 constant-effort mist-netting and banding stations. MAPS was designed to provide information on the vital rates (productivity or birth rate, and survivorship or death rate) of landbirds that is critically needed for efforts to identify demographic causes of the severe and sometimes accelerating population declines documented (Robbins et al. 1989, Terborgh 1989, Peterjohn et al. 1995) for many species of North American landbirds (DeSante 1992, DeSante et al. 1995, 1999, [2001a](#)). Such data on vital rates are also critically needed in efforts to identify management strategies to reverse such population declines (DeSante 1995, DeSante and Rosenberg 1998).

MAPS is organized to fulfill three sets of goals and objectives: monitoring, research, and management. The specific **monitoring** goals of MAPS are to provide, for over 100 target species, including Neotropical-wintering migrants, temperate-wintering migrants, and permanent residents: (a) annual indices of adult population size and post-fledging productivity from data on the numbers and proportions of young and adult birds captured; and (b) annual estimates of adult population size, adult survival rates, proportions of residents, and recruitment into the adult population from modified Cormack- Jolly-Seber analyses of mark-recapture data on adult birds.

The specific **research** goals of MAPS are to identify and describe: (a) temporal and spatial patterns in these demographic indices and estimates at a variety of spatial scales ranging from the local landscape to the entire continent; and (b) relationships between these patterns and ecological characteristics of the target species, population trends of the target species, station-specific and landscape-level habitat characteristics, and spatially-explicit weather variables.

The specific **management** goals of MAPS are to use these patterns and relationships, at the appropriate spatial scales, to: (a) identify thresholds and trigger points to notify appropriate agencies and organizations of the need for further research and/or management actions; (b) determine the proximate demographic cause(s) of population change; (c) suggest management actions and conservation strategies to reverse population declines and maintain stable or increasing populations; and (d) evaluate the effectiveness of the management actions and conservation strategies actually implemented through an adaptive management framework.

All of these goals are in agreement with the Department of Defense (DoD) Partners-in-Flight (PIF) strategy. Moreover, because birds are excellent indicators of the health of ecological systems, their reproductive success and survival can serve as sensitive barometers of the overall effectiveness of efforts to maintain the biodiversity and ecological integrity of military installations. Accordingly, the MAPS program was initiated on select military installations beginning in 1992 and soon became one of the focus projects of the DoD PIF program. It was expected that information from the MAPS program would be capable of aiding research and management efforts on these military installations to protect and enhance the installations'

avifauna and ecological integrity, while allowing them to fulfill their military mission.

Accordingly, in 1995, six MAPS stations were established and operated on Fort Bragg. The operation of these stations during the summers of 1995 and 1996 and the subsequent analyses of data from those years were accomplished through funding from U.S. Army Fort Bragg. Operation of these six MAPS station and associated data analyses during the three years 1997-1999 was accomplished by means of funding from the DoD Legacy Resource Management Program. The operation of the six stations was continued during the summers of 2000 through 2007 by means of funding from Fort Bragg, while the comprehensive analyses of data from 1995-2002 was funded by the DoD Legacy Resource Management Program.

The initial objective of the MAPS Program on DoD installations such as Fort Bragg has been to identify generalized management guidelines and formulate specific management actions that could be implemented on military installations and elsewhere to reverse the population declines of target landbird species and to maintain the populations of stable or increasing species. The identification and formulation of these management guidelines and actions was to be achieved by modeling the vital rates (productivity and survivorship) of the various landbird species as a function of landscape-level habitat characteristics and spatially explicit weather variables. The goal was to identify relationships between adult population size, numbers of young produced, productivity (ratio of young to adults), and trends in those parameters and these habitat and weather variables. The resultant management strategies were designed to involve efforts to modify the habitat from characteristics associated with low population size, population trend, or productivity to characteristics associated with high population size, population trend, or productivity (especially for species for which low productivity was found to be driving the population decline).

The Legacy Resource Management Program allowed us to undertake these analyses and formulate management strategies. These analyses were completed in 2003 and management guidelines were formulated for ten bird species of conservation concern that breed in the southeastern United States ([Nott et al. 2003](#)). With additional funding from the Legacy Resource Management Program, we are currently implementing these guidelines through management actions on eight military installations (including Fort Bragg) in conjunction with efforts to increase military Readiness and Range Sustainment ([Nott and Michel 2005](#)). The strategy for implementing these guidelines includes the establishment of new MAPS stations to monitor the effectiveness of such proposed or on-going management, the discontinuance of an equal number of old stations, and the continued operation of others of the old stations to serve as controls for the new management stations. In this way, the total number of stations operated will remain the same.

At Fort Bragg Wood Thrush and Prairie Warbler, were identified as management species of concern. Following the recommendations of [Nott et al. \(2003\)](#), the I102 station was discontinued in 2003 to reduce the probability of capturing endangered Red-cockaded Woodpeckers that breed within the boundaries of that station. The I102 station was replaced by the Sandstone Hill station in a mosaic of upland patchy forest, shrubland, and grasslands that are frequently managed to reduce fire risks.

A complete summary of the results of the MAPS Program on Fort Bragg from 1993-1999, as well as on 12 other installations or groups of nearby installations in the eastern United States, was presented by DeSante et al. (2001b), a summary of 2000-2005 results was presented by DeSante et al. (2002, 2004, 2005), Pyle et al. (2006), and Nott et al. (2007). This report briefly updates these earlier reports and documents the operation of the six MAPS stations on Fort Bragg during the 2007 breeding season.

Methods

Six MAPS stations were operated in 2007, in the same locations where they were first established in 1995 (five stations) or 2003 (Sandstone Hill station). Each of these six MAPS stations was operated in accordance with the highly standardized banding protocols established by The Institute for Bird Populations for use by the MAPS Program throughout North America and spelled out in detail in the MAPS Manual ([DeSante et al. 2006](#)). On each day of operation each year, one 12-m long, 30-mm mesh, 4-tier nylon mist net was erected at each of ten fixed mist-netting sites within the interior eight hectares of each 20 hectare station. These ten nets at each station were operated for six morning hours per day (beginning at local sunrise), and for one day in each of nine consecutive 10-day periods between May 14 and August 7 (Table 1). The operation of stations occurred on schedule in each of the ten-day periods and was carried out by IBP field biologist intern Emily Hockman, who was trained by IBP field biologist Danielle Kaschube.

With few exceptions, all birds captured during the course of the study were identified to species, age, and sex and, if unbanded, were banded with USGS/BRD numbered aluminum bands. Birds were released immediately upon capture and before being banded or processed if situations arose where bird safety would be compromised. The following data were taken on all birds captured, including recaptures, according to MAPS guidelines using standardized codes and forms ([DeSante et al. 2006](#)):

- (1) capture code (newly banded, recaptured, band changed, unbanded);
- (2) band number;
- (3) species;
- (4) age and how aged;
- (5) sex (if possible) and how sexed (if applicable);
- (6) extent of skull pneumaticization;
- (7) breeding condition of adults (i.e., extent of cloacal protuberance or brood patch);
- (8) extent of juvenal plumage in young birds;
- (9) extent of body and flight-feather molt;
- (10) extent of primary-feather wear;
- (11) presence of molt limits and plumage characteristics;
- (12) wing chord;
- (13) fat class and body mass;
- (14) date and time of capture (net-run time);
- (15) station and net site where captured; and

(16) any pertinent notes.

Effort data (i.e., the number and timing of net-hours on each day of operation) were also collected in a standardized manner. In order to allow constant-effort comparisons of data to be made, the times of opening and closing the array of mist nets and of beginning each net check were recorded to the nearest ten minutes. The breeding (summer residency) status (confirmed breeder, likely breeder, non-breeder) of each species seen, heard, or captured at each MAPS station on each day of operation was recorded using techniques similar to those employed for breeding bird atlas projects.

The computer entry, proofing, and verification of all banding, effort, and breeding status data were completed by IBP biologists using specially designed data entry, verification, and editing programs. The critical data for each banding record (capture code, band number, species, age, sex, date, capture time, station, and net number) were proofed by hand against the raw data and any computer-entry errors were corrected. All banding data were then run through a series of verification programs as follows:

- (1) Clean-up programs to check the validity of all codes entered and the ranges of all numerical data;
- (2) Cross-check programs to compare station, date, and net fields from the banding data with those from the effort and breeding status data;
- (3) Cross-check programs to compare species, age, and sex determinations against degree of skull pneumaticization, breeding condition (extent of cloacal protuberance and brood patch), extent of juvenal plumage, extent of body and flight-feather molt, extent of primary-feather wear, and presence of molt limits and plumage characteristics;
- (4) Screening programs which allow identification of unusual or duplicate band numbers or unusual band sizes for each species; and
- (5) Verification programs to screen banding and recapture data from all years of operation for inconsistent species, age, or sex determinations for each band number.

Any discrepancies or suspicious data identified by any of these programs were examined manually and corrected if necessary. Wing chord, body mass, fat content, date and station of capture, and any pertinent notes were used as supplementary information for the correct determination of species, age, and sex in all of these verification processes. The proofed, verified, and corrected banding data from each year were then run through a series of analysis programs that calculated for each species and for all species pooled at each station and for all stations pooled on each forest:

- (1) the numbers of newly banded birds, recaptured birds, and birds released unbanded;
- (2) the numbers and capture rates (per 600 net-hours) of first captures (in each year) for individual adult and young birds; and
- (3) the proportion of young in the catch.

Following the procedures pioneered by the British Trust for Ornithology (BTO) in their CES Scheme (Peach et al. 1996), the number of adult birds captured was used as an index of adult population size. For our estimate of post-fledging productivity, we are now using “reproductive index” (number of young divided by number of adults) as opposed to “proportion of young in the catch” previously used. Reproductive index is a more intuitive value for productivity, and it is also more comparable to other calculated MAPS parameters such as recruitment indices.

Survival of target species was estimated using Modified Cormack-Jolly-Seber (CJS) mark-recapture analyses (Pollock et al. 1990, Lebreton et al. 1992) on 13 years (1995-2007) of capture histories of adult birds from the seven stations ever operated at this location. Target species were those for which, on average, at least 2.5 individual adults per year and at least two between-year returns were recorded from the seven stations pooled, at which the species was a breeder during more than half of the years the station was operated. Using the computer program TMSURVIV (White 1983, Hines et al. 2003), we calculated, for each target species, maximum-likelihood estimates and standard errors (*SEs*) for adult survival probability, adult recapture probability, and the proportion of residents among newly captured adults using a time-constant, between- and within-year transient model (Pradel et al. 1997, Nott and DeSante 2002, Hines et al. 2003). The use of the transient model accounts for the existence of transient adults (dispersing and floater individuals which are only captured once) in the sample of newly captured birds, and provides survival estimates that are unbiased with respect to these transient individuals (Pradel et al. 1997). Recapture probability is defined as the conditional probability of recapturing a bird in a subsequent year that was banded in a previous year, given that it survived and returned to the place it was originally banded.

Results and Discussion

We operated six MAPS stations on Fort Bragg during the summer of 2007 (Table 1). A total of 2029.3 net-hours were accumulated at all six stations pooled, representing 62.6% of the maximum possible effort (3400 net-hours) at the six stations. The reason for the lower than normal proportion of effort was because operation began late in 2007 (in period 5 rather than period 3) due to funding uncertainties. Of these net hours, 1869.5 overlap with those of 2006. The details of the operation of these six stations during 2007 are presented in Table 1.

For each individual species and for all species pooled, the numbers of birds newly banded, captured and released unbanded, and recaptured are presented for each station in Table 2 and, for all stations combined, in Table 4. A total of 344 captures of 37 species occurred at Fort Bragg during the summer of 2007 (Table 4). Newly banded birds comprised 66.8% of the total captures. The greatest number of total captures (69) was recorded at the Sandstone Hill station and the smallest number of total captures (26) was recorded at the S112 station. The highest species richness occurred at Stations I104 and I113 (20 species each) and the lowest species richness occurred at S112 (9 species).

The capture rates (per 600 net-hours) of individual adult and young birds and the proportion of young in the catch are presented for each species and for all species pooled at each station in Table 3 and, for all stations combined, in Table 4. We present capture rates (captures per 600

net-hours) of adults and young in these tables so that the data can be compared among stations which, because of the vagaries of weather and accidental net damage, can differ from one another in effort expended (Table 1). Adult population size (for all species pooled) was highest at Station S114 (52.7 adults/600 net hours; Table 3), followed by Sandstone Hill and Station I104 (51.7 each), Station I113 (51.7), Station S110 (45.9), and Station S112 (31.3). Reproductive index (number of young birds per adult) showed a different pattern, being highest at Sandstone Hill (1.14), followed by Station I104 (0.48), Station S110 (0.46), Station S113 (0.44), Station I114 (0.27), and Station S112 (0.22). The mean adult capture rate for the six stations combined was 47.0 per 600 net hours in 2007, compared with 41.3 adults per 600 net-hours for the same six stations in 2006, and the overall reproductive index was 0.52 in 2007, compared with 0.55 in 2006, indicating slightly larger breeding population sizes and slightly decreased productivity between the two years.

Among individual species, Carolina Wren was the most frequently captured species at the six stations in 2007, followed by Pine Warbler, Common Yellowthroat, Ruby-throated Hummingbird, Northern Cardinal, Prairie Warbler, and Tufted Titmouse (Table 4). The most abundant breeding species, having a capture rate of at least 2.0 adults per 600 net-hours, in decreasing order, were Carolina Wren, Northern Cardinal, Prairie Warbler, Common Yellowthroat, Tufted Titmouse, Hooded Warbler, Eastern Towhee, and Indigo Bunting (Table 4); the rate for Ruby-throated Hummingbird could not be calculated because individuals are not banded. The most abundant breeding species at each station, having a capture rate of at least 3.0 birds per 600 net-hours in 2007 are as follows (species of concern, as noted above, in italics):

Sandstone Hill

Prairie Warbler
Great Crested Flycatcher*
Pine Warbler
Bachman's Sparrow
Blue Grosbeak
Northern Flicker*
Chipping Sparrow

S110

Prairie Warbler
Common Yellowthroat
Red-eyed Vireo*
Carolina Wren*
Hooded Warbler*
Bachman's Sparrow*
Northern Cardinal*
Indigo Bunting
Great Crested Flycatcher[†]

I 104

Common Yellowthroat
Carolina Wren*
Brown Thrasher*
Eastern Wood-Pewee*
*Prairie Warbler**
American Goldfinch
Carolina Chickadee[†]
Eastern Towhee[†]
Chipping Sparrow[†]

S114

Carolina Wren
Hooded Warbler
Eastern Towhee*
Indigo Bunting
Tufted Titmouse*
Northern Cardinal
Blue-gray Gnatcatcher[†]

I113

Tufted Titmouse*
Carolina Wren
Eastern Towhee
Northern Cardinal
Common Yellowthroat
Hooded Warbler*
Chipping Sparrow*
Pine Warbler[†]
American Goldfinch[†]
Great Crested Flycatcher[†]
Summer Tanager[†]
Prairie Warbler[†]

S112

Northern Cardinal
Tufted Titmouse
Carolina Wren
American Goldfinch*

* At least 3.0 adults per 600 net hours in 2007 but not in 2006.

[†] At least 3.0 adults per 600 net hours in 2006 but not in 2007.

As can be seen from the number and position of marked species (* and †), there was considerable turnover between 2006 and 2007, despite a similar overall capture rate. Prairie Warbler, a species of concern, increased at one station but decreased at another, whereas Carolina Wren and Hooded Warbler each increased at two stations.

Using 13 years of data (1995-2007) from all seven stations ever operated on Fort Bragg combined, estimates of adult survival and recapture probabilities were obtained for 20 target species breeding at Fort Bragg. Maximum-likelihood estimates of annual adult survival probability, recapture probability, and proportion of residents among newly captured adults from the time-constant transient model are presented in Table 5 for these 20 species. Survival-rate estimates for all 20 species showed variable precision (CVs between 13% and 76%) with a mean CV of 33.8%. The mean CV for 17 species with estimates last year (Nott et al. 2006) was 29.4% compared with 31.4% after 12 years, indicating continued improvement in precision with the addition of an 13th year at Fort Bragg. Annual adult survival rates for these 20 species in 2007 ranged from a low of 0.279 for Great Crested Flycatcher to a high of 0.547 for Red-eyed Vireo and Ovenbird, with a mean survival rate of 0.424 for the 20 species.

More precise estimates (CV<20%) were revealed for seven species, which included a group of five resident species (Carolina Chickadee, Tufted Titmouse, Carolina Wren, Eastern Towhee, and Northern Cardinal) and two Neotropical migrants (Prairie Warbler, Common Yellowthroat).

Survival estimates are low at Fort Bragg compared to other locations, especially for resident species. In comparing survival values from Fort Bragg (1995-2007) with those of the [Southeast Region of the United States \(1992-2001\)](#), for example, survival at Fort Bragg was lower than that of the Southeast Region for 9 of the 14 target species, with the mean at Fort Bragg (0.424) being 12.5% lower than that of the Southeast Region (0.485). Importantly, survival for the five resident species were substantially lower at Fort Bragg (mean 0.415) than in the Southeast Region (mean 0.478), perhaps indicating problems with survival of landbirds on the Fort Bragg installation and/or vicinity. In the Breeding Bird Survey 1980-2007 trend data for North Carolina (Sauer et al. 2007) Carolina Chickadee significantly declined (P<0.05) and Northern Cardinal significantly (P<0.05) increased, while the other three species remained stable. This further supports the hypothesis that there is a local problem with the populations of resident birds.

The survival rates of the two Neotropical migrants, Prairie Warbler (0.381) and Common Yellowthroat (0.351), were also lower than those for Southeast region (0.421 and 0.420, respectively). Breeding Bird Survey trend data (1980-2007) showed stable populations of both species in North Carolina.

These results provide a strong suggestion that overwintering survival of individuals wintering on Fort Bragg may be poor (DeSante et al. 2004, 2005a). The Institute for Bird Populations has initiated the Monitoring Avian Wintering Survival (MAWS) Program to assess habitat-specific overwintering survival rates in the southern parts of the United States. Six of the initial 24 MAWS stations have been established on Fort Bragg through funding from the Legacy Resources Management Program. Four or five years of data from these MAWS stations should

be able to provide information as to the extent of any overwintering survival problem and relationships between survival and various habitat variables. However, no significant declines or increases were detected in MAWS data between the winters of 2003-2004 and 2006-2007. The MAWS Program if it can be re-established, in conjunction with MAPS, should lead to the formulation of management strategies and guidelines to enhance overwintering survival, especially for declining species of conservation concern that overwinter in the United States.

As mentioned earlier, analyses aimed at identifying and describing relationships between four demographic parameters (adult population size, population trends, numbers of young, and productivity) and landscape-level habitat characteristics have been completed for 13 military installations including Fort Bragg ([Nott et al. 2003](#), [Nott and Michel 2005](#)). These analyses were funded by Legacy Resource Management Program Project #103. At Fort Bragg, two species (Wood Thrush and Prairie Warbler) emerged as candidates for particular management concern. Since, the numbers of Wood Thrush have declined below acceptable levels and it is no longer considered a target species of management concern.

In 2003 the I102 station was replaced by the Sandstone Hill station in a mosaic of upland patchy forest, shrubland, and grasslands that are frequently managed to reduce risk of wildfire. This station was established to specifically monitor the effects of a prescribed fire regime upon Prairie Warbler populations. Prior to the 2004 season the area around Sandstone Hill was burned, and in 2004 a single Prairie Warbler was captured. In 2005 the capture rates Prairie Warblers increased abruptly to 15.5 adult and 4.8 young individuals per 600 net-hours. In 2006, these values were 14.3 and 0.0, respectively, and in 2007 these values were 12.5 and 1.8 (Table 3). This indicates that the management actions at Sandstone Hill successfully resulted in increased populations of Prairie Warblers, but that perhaps the habitat is beginning to succeed, resulting in a slight decreasing trend from 2005 to 2007 and fewer young. Since 2002, Wood Thrush captures and recaptures recorded at S112 have declined; they were not captured at all at Fort Bragg in 2007. However, Bachman's Sparrow, a USFWS Bird of Conservation Concern with IUCN Red List near threatened status, was captured at Sandstone Hill, I113, and S110 in 2007 (Table 3).

In summary, our data suggest that Prairie Warblers will occupy potential habitat immediately following a springtime burn but that breeding individuals will recruit into the habitat the second year after fire. As this fire-managed "disclimax" community succeeds towards forest, we predict that continued effectiveness monitoring of these populations will detect the onset of a decline in reproductive success or population size. Perhaps that has already becoming apparent, with a slight decline in breeding adults between 2005 and 2007. However, based on an understanding of these temporal dynamics, we can adjust the prescribed fire frequency to maximize mean annual productivity of Prairie Warbler populations (and perhaps Bachman's Sparrows and other species) while meeting the management goals associated with Readiness and Range Sustainment. Through such adaptive management cycles, we are confident that we can achieve the long-term goal of reversing declining populations and maintaining stable or increasing source populations of target landbird species at Fort Bragg and other military installations.

Acknowledgments

We thank IBP field biologist intern Emily Hockman for operating the station in 2007, and IBP biologist Danielle Kaschube for providing additional training and supervision. We also thank volunteer Lara Hartley for assistance with station operation. We especially thank Jessie Schillaci for her enthusiastic support of and kind assistance with all of the logistical and administrative aspects of this work. Financial support for this program was provided by U.S. Army Fort Bragg for which we are very grateful. This is Contribution Number 336 of The Institute for Bird Populations.

Literature Cited

- DeSante, D.F. (1992) Monitoring Avian Productivity and Survivorship (MAPS): a sharp, rather than blunt, tool for monitoring and assessing landbird populations. *In*: D. R. McCullough and R. H. Barrett (Eds.), Wildlife 2001: Populations, pp. 511-521. (London, U.K.: Elsevier Applied Science).
- DeSante, D.F. (1995) Suggestions for future directions for studies of marked migratory landbirds from the perspective of a practitioner in population management and conservation. Journal Applied Statistics 22, pp. 949-965.
- DeSante, D.F., Burton, K.M., Saracco, J.F., & Walker, B.L. (1995) Productivity indices and survival rate estimates from MAPS, a continent-wide programme of constant-effort mist netting in North America. Journal Applied Statistics, 22, pp. 935-947.
- DeSante, D.F., Burton, K.M., Velez, P., & Froehlich, D. (2007) MAPS Manual, Point Reyes Station, CA: The Institute for Bird Populations; 49 pp.
- [DeSante, D.F., Nott, M.P., & O'Grady, D.R. \(2001a\)](#) Identifying the proximate demographic cause(s) of population change by modeling spatial variation in productivity, survivorship, and population trends. Ardea 89 (special issue):185-207.
- DeSante, D.F. and D.R. O'Grady. (2000). The Monitoring Avian Productivity and Survivorship (MAPS) program 1997 and 1998 report. Bird Populations 5:49-101.
- DeSante, D.F., & O'Grady, D. (2002) The 2000-2002 report of the Monitoring Avian Productivity and Survivorship (MAPS) Program on Fort Bragg. Point Reyes Station, CA: The Institute for Bird Populations; 7 pp. plus 12 tables.
- DeSante, D.F., O'Grady, D.R. & Pyle, P. (1999) Measures of productivity and survival derived from standardized mist netting are consistent with observed population changes. Bird Study 46 (suppl.):S178-188.
- DeSante, D.F., Pyle, P., and O'Grady, D.R. (2001b) The 1992-1999 report of the Monitoring Avian Productivity and Survivorship (MAPS Program on military installations in eastern United States. The Institute for Bird Populations, Point Reyes Station, CA 113 pp.
- DeSante, D.F., Pyle, P., & Kaschube, D. (2004) The 2003 report of the Monitoring Avian Productivity and Survivorship (MAPS) Program at Fort Bragg. Point Reyes Station, CA: The Institute for Bird Populations; 10 pp.
- DeSante, D.F., Pyle, P., & Kaschube, D. (2005) The 2004 report of the Monitoring Avian Productivity and Survivorship (MAPS) Program at Fort Bragg. Point Reyes Station, CA: The Institute for Bird Populations; 10 pp.
- DeSante, D.F., & Rosenberg, D.K. (1998) What do we need to monitor in order to manage landbirds? *In*: J. Marzluff & R. Sallabanks (Eds.), Avian Conservation: Research Needs and

- Effective Implementation, pp. 93-106. Island Press, Washington, DC.
- Hines, J.E., Kendall, W.L., & Nichols, J.D. (2003) On the use of the robust design with transient capture-recapture models. Auk, 120, pp.1151-1158
- Lebreton, J.-D., Burnham, K.P., Clobert, J., & Anderson, D.R. (1992) Modeling survival and testing biological hypotheses using marked animals: a unified approach with case studies, Ecological Monographs, 62, pp. 67-118.
- Nott, M.P., & DeSante, D.F. (2002) Demographic monitoring and the identification of transients in mark-recapture models. Pp. 727-736 *in*: J.M. Scott & P. Heglund (eds.), Predicting Species Occurrences: Issues of Scale and Accuracy. Island Press, NY.
- [Nott, M.P., & N. Michel. \(2005\) Management strategies for reversing declines in landbirds of conservation concern on military installations: Predictive modeling of landbird populations on military installations.](#) The Institute for Bird Populations, Pt. Reyes Station, CA.
- [Nott, M.P., D.F. DeSante, and N. Michel. \(2003\) Management strategies for reversing declines in landbirds of conservation concern on military installations: A landscape-scale analysis of MAPS data.](#) The Institute for Bird Populations, Pt. Reyes Station, CA.
- Nott, P., Pyle, P., and Kaschube, D. (2007) The 2006 report of the Monitoring Avian Productivity and Survivorship (MAPS) Program at Fort Bragg. Point Reyes Station, CA: The Institute for Bird Populations; 11 pp.
- Peach, W.J., Buckland, S.T., & Baillie, S.R. (1996) The use of constant effort mist-netting to measure between-year changes in the abundance and productivity of common passerines. Bird Study, 43, pp. 142-156.
- Peterjohn, B.G., Sauer, J.R., & Robbins, C.S. (1995) Population trends from the North American Breeding Bird Survey. *In*: T.E. Martin and D.M. Finch, Ecology and Management of Neotropical Migratory Birds, New York: Oxford University Press; pp. 3-39.
- Pollock, K.H., Nichols, J.D., Brownie, C., & Hines, J.E. (1990) Statistical inference for capture-recapture experiments, Wildlife Monographs, No. 107.
- Pradel, R., Hines, J., Lebreton, J.-D., & Nichols, J.D. (1997) Estimating survival probabilities and proportions of 'transients' using capture-recapture data. Biometrics, 53, pp. 60-72.
- Pyle, P., Kaschube, D., Nott, P., and Desante, D.F. (2006) The 2005 report of the Monitoring Avian Productivity and Survivorship (MAPS) Program at Fort Bragg. Point Reyes Station, CA: The Institute for Bird Populations; 10 pp.
- Robbins, C.S., Sauer, J.R., Greenberg, R.S., & Droege, S. (1989) Population declines in North American birds that migrate to the Neotropics, Proceedings of the National Academy of Sciences (USA), 86, pp. 7658-7662.
- Sauer, J. R., J. E. Hines, and J. Fallon. 2007. The North American Breeding Bird Survey, Results and Analysis 1966 - 2006. Version 10.13.2007. USGS Patuxent Wildlife Research Center, Laurel, MD
- Terborgh, J. (1989) Where Have All the Birds Gone? Essays on the Biology and Conservation of Birds that Migrate to the American Tropics, Princeton, NJ: Princeton Univ. Press; 207 pp.
- White, G.C. (1983) Numerical estimation of survival rates from band-recovery and biotelemetry data. J. Wildlife Management, 47, pp. 716-728.

Table 1. Summary of the 2007 MAPS program on U.S. Army Fort Bragg.

Station		No.	Major Habitat Type	Latitude-longitude	Avg Elev. (m)	2007 operation		
Name	Code					Total number of net-hours ¹	No. of periods	Inclusive dates
Sandstone Hill	SAHI	16706	Controlled burn pine savanna, mixed scrub oak woodland	35°03'04"N,79°19'37"W	152	336.7 (314.8)	6	6/18 – 8/01
I 104	I104	16657	Controlled burn pine savanna, riparian fields and scrub	35°07'00"N,79°19'13"W	84	336.7 (305.2)	6	6/15 – 8/04
I 113	I113	16658	Controlled burn riparian, savanna and pine-oak woodland	35°05'37"N,79°19'28"W	110	329.7 (320.5)	6	6/20 – 8/05
S 110	S110	16659	Riparian woodland, pine savanna and pine-oak woodland	35°07'07"N,79°20'04"W	94	340.0 (286.7)	6	6/14 – 8/02
S 114	S114	16661	Pine-oak and riparian woodland bordering grain fields	35°02'58"N,79°16'11"W	70	341.3 (324.0)	6	6/19 – 7/31
S 112	S112	16660	Pine-oak mixed with riparian woodland	35°06'46"N,79°21'45"W	114	345.0 (318.3)	6	6/13 – 8/03
ALL STATIONS COMBINED						2029.3(1869.5)	6	6/13 – 8/05

¹ Total net-hours in 2007. Net-hours in 2007 that could be compared in a constant-effort manner to 2006 are shown in parentheses.

Table 2. (cont.) Capture summary for the six individual MAPS stations operated on U.S. Army Fort Bragg in 2007.
 N = Newly Banded, U = Unbanded, R = Recaptures of banded birds.

Species	Sandstone Hill			I 104			I 113			S 110			S 114			S 112		
	N	U	R	N	U	R	N	U	R	N	U	R	N	U	R	N	U	R
Kentucky Warbler													1					
Common Yellowthroat				7	1	8	3			4		4	1					
Hooded Warbler							2			2		1	4		4			
Summer Tanager	1			1														
Eastern Towhee				2			3		1	1			3					
Bachman's Sparrow	4		1				1			2	1							
Chipping Sparrow	3		1				2											
Northern Cardinal				2		2	6		3	2		1	2			4		3
Blue Grosbeak	3		1	1	1					1						1		
Indigo Bunting				1			1			2			3	1	1			
American Goldfinch				2			1									2		
ALL SPECIES POOLED	61	2	6	37	5	21	34	7	10	36	11	21	37	8	22	18	2	6
Total Number of Captures		69			63			51			68			67			26	
Number of Species	13	1	5	19	4	6	17	2	4	16	2	7	13	2	6	8	2	3
Total Number of Species		14			20			20			17			14			9	

Table 3. Numbers of adult and young individual birds captured per 600 net-hours and reproductive index (young/adult) at the six individual MAPS stations operated on U.S. Army Fort Bragg in 2007.

Species	Sandstone Hill			I 104			I 113			S 110			S 114			S 112		
	Ad.	Yg.	Repr. index	Ad.	Yg.	Repr. index	Ad.	Yg.	Repr. index	Ad.	Yg.	Repr. index	Ad.	Yg.	Repr. index	Ad.	Yg.	Repr. index
Red-bellied Woodpecker				1.8	0.0	0.00												
Downy Woodpecker	0.0	1.8	und. ¹	0.0	1.8	und. ¹												
Red-cockaded Woodpecker	1.8	0.0	0.00															
Northern Flicker	3.6	0.0	0.00															
Eastern Wood-Pewee				3.6	0.0	0.00							1.8	0.0	0.00			
Acadian Flycatcher																1.7	0.0	0.00
Great Crested Flycatcher	5.3	0.0	0.00	1.8	0.0	0.00	0.0	1.8	und. ¹	1.8	0.0	0.00				1.7	0.0	0.00
Eastern Kingbird				1.8	0.0	0.00												
Loggerhead Shrike	1.8	0.0	0.00															
White-eyed Vireo							0.0	1.8	und.				1.8	0.0	0.00			
Yellow-throated Vireo							1.8	0.0	0.00									
Red-eyed Vireo				1.8	0.0	0.00				3.5	1.8	0.50						
Blue Jay	1.8	0.0	0.00															
Carolina Chickadee	1.8	0.0	0.00	1.8	0.0	0.00	1.8	0.0	0.00	1.8	1.8	1.00	0.0	3.5	und. ¹			
Tufted Titmouse				1.8	1.8	1.00	5.5	5.5	1.00	0.0	1.8	und. ¹	3.5	1.8	0.50	5.2	1.7	0.33
White-breasted Nuthatch	1.8	0.0	0.00				1.8	0.0	0.00									
Brown-headed Nuthatch							1.8	0.0	0.00									
Carolina Wren				5.3	7.1	1.33	5.5	1.8	0.33	3.5	12.4	3.50	17.6	8.8	0.50	5.2	5.2	1.00
Blue-gray Gnatcatcher				0.0	1.8	und.				1.8	0.0	0.00						
Gray Catbird				5.3	0.0	0.00				1.8	0.0	0.00	1.8	0.0	0.00			
Brown Thrasher										1.8	0.0	0.00						

Table 3. (cont.) Numbers of adult and young individual birds captured per 600 net-hours and reproductive index (young/adult) at the six individual MAPS stations operated on U.S. Army Fort Bragg in 2007.

Species	Sandstone Hill			I 104			I 113			S 110			S 114			S 112		
	Ad.	Yg.	Repr. index	Ad.	Yg.	Repr. index	Ad.	Yg.	Repr. index	Ad.	Yg.	Repr. index	Ad.	Yg.	Repr. index	Ad.	Yg.	Repr. index
Pine Warbler	5.3	51.7	9.67				1.8	0.0	0.00				1.8	0.0	0.00			
Prairie Warbler	12.5	1.8	0.14	3.6	1.8	0.50	0.0	1.8	und.	7.1	1.8	0.25						
American Redstart							1.8	0.0	0.00									
Louisiana Waterthrush																1.7	0.0	0.00
Kentucky Warbler													1.8	0.0	0.00			
Common Yellowthroat				10.7	7.1	0.67	3.6	1.8	0.50	5.3	1.8	0.33	1.8	0.0	0.00			
Hooded Warbler							3.6	0.0	0.00	3.5	0.0	0.00	7.0	0.0	0.00			
Summer Tanager	1.8	0.0	0.00	1.8	0.0	0.00												
Eastern Towhee				1.8	1.8	1.00	5.5	0.0	0.00	1.8	0.0	0.00	5.3	0.0	0.00			
Bachman's Sparrow	5.3	1.8	0.33				1.8	0.0	0.00	3.5	0.0	0.00						
Chipping Sparrow	3.6	1.8	0.50				3.6	0.0	0.00									
Northern Cardinal				1.8	1.8	1.00	5.5	7.3	1.33	3.5	0.0	0.00	3.5	0.0	0.00	10.4	0.0	0.00
Blue Grosbeak	5.3	0.0	0.00	1.8	0.0	0.00				1.8	0.0	0.00				1.7	0.0	0.00
Indigo Bunting				1.8	0.0	0.00	1.8	0.0	0.00	3.5	0.0	0.00	5.3	0.0	0.00			
American Goldfinch				3.6	0.0	0.00	1.8	0.0	0.00							3.5	0.0	0.00
ALL SPECIES POOLED	51.7	58.8	1.14	51.7	25.0	0.48	49.1	21.8	0.44	45.9	21.2	0.46	52.7	14.1	0.27	31.3	7.0	0.22
Number of Species	13	5		17	8		16	7		15	6		12	3		8	2	
Total Number of Species		14			19			19			16			13			8	

¹ Reproductive index (young/adult) is undefined because no adults of this species were captured at this station in this year.

Table 4. Summary of results for all six U.S. Army Fort Bragg MAPS stations combined in 2007.

Species	Birds captured			Birds/600 nethours		Reprod. Index
	Newly banded	Un-banded	Recap-tured	Adults	Young	
Ruby-throated Hummingbird		25				
Red-bellied Woodpecker	1			0.3	0.0	0.00
Downy Woodpecker	2			0.0	0.6	und. ¹
Red-cockaded Woodpecker			1	0.3	0.0	0.00
Northern Flicker	2			0.6	0.0	0.00
Eastern Wood-Pewee	3			0.9	0.0	0.00
Acadian Flycatcher	1			0.3	0.0	0.00
Great Crested Flycatcher	7			1.8	0.3	0.17
Eastern Kingbird	1			0.3	0.0	0.00
Loggerhead Shrike	1			0.3	0.0	0.00
White-eyed Vireo	2			0.3	0.3	1.00
Yellow-throated Vireo	1			0.3	0.0	0.00
Red-eyed Vireo	4		1	0.9	0.3	0.33
Blue Jay	1			0.3	0.0	0.00
Carolina Chickadee	6		3	1.2	0.9	0.75
Tufted Titmouse	14		5	2.7	2.1	0.78
White-breasted Nuthatch	1	1		0.6	0.0	0.00
Brown-headed Nuthatch	1			0.3	0.0	0.00
Carolina Wren	35	1	35	6.2	5.9	0.95
Blue-gray Gnatcatcher	2	2		0.3	0.3	1.00
Gray Catbird	5		3	1.5	0.0	0.00
Brown Thrasher	1			0.3	0.0	0.00
Pine Warbler	34	2	2	1.5	8.6	5.80
Prairie Warbler	15		5	3.8	1.2	0.31
American Redstart	1			0.3	0.0	0.00
Louisiana Waterthrush	1			0.3	0.0	0.00
Kentucky Warbler	1			0.3	0.0	0.00
Common Yellowthroat	15	1	12	3.5	1.8	0.50
Hooded Warbler	8		5	2.4	0.0	0.00
Summer Tanager	2			0.6	0.0	0.00
Eastern Towhee	9		1	2.4	0.3	0.13
Bachman's Sparrow	7	1	1	1.8	0.3	0.17
Chipping Sparrow	5		1	1.2	0.3	0.25

Table 4. Summary of results for all six U.S. Army Fort Bragg MAPS stations combined in 2007.

Species	Birds captured			Birds/600 nethours		Reprod. Index
	Newly banded	Un-banded	Recap-tured	Adults	Young	
Northern Cardinal	16		9	4.1	1.5	0.36
Blue Grosbeak	6	1	1	1.8	0.0	0.00
Indigo Bunting	7	1	1	2.1	0.0	0.00
American Goldfinch	5			1.5	0.0	0.00
ALL SPECIES POOLED	223	35	86	47.0	24.5	0.52
Total Number of Captures		344				
Number of Species	35	9	16	35	15	
Total Number of Species		37			36	

¹ Reproductive index (young/adult) is undefined because no adults of this species were captured at this location in this year.

Table 5. Estimates of adult annual survival and recapture probabilities and proportion of residents among newly captured adults using a time-constant model for 20 species breeding at MAPS stations on U.S. Army Fort Bragg obtained from 13 years (1995-2007) of mark-recapture data.

Species	Num. sta2. ¹	Num. ind. ²	Num. caps. ³	Num. ret. ⁴	Survival probability ⁵	Surv. C.V. ⁶	Recapture probability ⁷	Proportion of residents ⁸
Red-bellied Woodpecker ‡†	5	40	45	2	0.380 (0.288)	75.6	0.087 (0.223)	1.000 (2.614)
Great Crested Flycatcher ‡	6	124	139	8	0.279 (0.141)	50.6	0.218 (0.213)	0.815 (0.838)
White-eyed Vireo	2	44	90	6	0.315 (0.139)	44.0	0.543 (0.298)	0.455 (0.344)
Red-eyed Vireo ‡	4	62	70	4	0.547 (0.188)	34.3	0.166 (0.168)	0.339 (0.364)
Carolina Chickadee	6	102	139	14	0.535 (0.101)	18.8	0.319 (0.124)	0.340 (0.165)
Tufted Titmouse	6	130	221	28	0.370 (0.069)	18.8	0.724 (0.129)	0.465 (0.158)
Carolina Wren	5	167	375	37	0.294 (0.054)	18.4	0.726 (0.125)	0.794 (0.235)
Blue-gray Gnatcatcher ‡	6	67	78	3	0.416 (0.214)	51.4	0.357 (0.308)	0.089 (0.104)
Wood Thrush	1	58	85	6	0.492 (0.157)	31.9	0.141 (0.114)	0.727 (0.595)
Gray Catbird	2	114	173	9	0.451 (0.122)	27.1	0.189 (0.112)	0.437 (0.278)
Brown Thrasher	2	49	65	5	0.494 (0.153)	30.9	0.411 (0.229)	0.090 (0.096)
Pine Warbler	6	112	122	5	0.410 (0.195)	47.5	0.198 (0.198)	0.338 (0.357)
Prairie Warbler	4	231	348	34	0.381 (0.068)	17.9	0.381 (0.104)	0.623 (0.199)
Ovenbird †	2	40	55	8	0.547 (0.132)	24.2	0.184 (0.125)	1.000 (0.721)
Common Yellowthroat	5	365	836	51	0.351 (0.047)	13.5	0.627 (0.099)	0.235 (0.069)
Summer Tanager	6	82	98	7	0.387 (0.158)	40.8	0.155 (0.154)	0.944 (0.974)
Eastern Towhee	6	113	200	32	0.403 (0.067)	16.7	0.771 (0.111)	0.504 (0.158)
Northern Cardinal	5	176	409	53	0.473 (0.050)	10.6	0.663 (0.084)	0.320 (0.099)
Indigo Bunting	4	59	68	6	0.539 (0.164)	30.4	0.343 (0.213)	0.284 (0.218)
American Goldfinch ‡†	4	49	53	2	0.414 (0.303)	73.2	0.071 (0.198)	1.000 (2.853)

¹ Analysis of all stations pooled include data from 1995-2007 from the I 104, I 113, S 110, and S112, from 1995-2002 from the S 102 station and 2003-2007 from the Sandstone Hill station which replaced the S 102 station. Only data from 1995-2001 is included from the S114 station.

² Number of stations where the species was a regular or usual breeder and at which adults of the species were captured. Stations within one km of each other were combined into a single super-station to prevent individuals whose home ranges included portions of two or more stations from being counted as multiple individuals.

³ Number of adult individuals captured at stations where the species was a regular or usual breeder (i.e., number of capture histories).

Table 5. (cont.) Estimates of adult annual survival and recapture probabilities and proportion of residents among newly captured adults using a time-constant model for 20 species breeding at MAPS stations on U.S. Army Fort Bragg obtained from 13 years (1995-2007) of mark-recapture data.

⁴ Total number of captures of adult birds of the species at stations where the species was a regular or usual breeder.

⁵ Total number of returns. A return is the first recapture in a given year of a bird originally banded at the same station in a previous year.

⁶ Survival probability (ϕ) presented as the maximum likelihood estimate (standard error of the estimate).

⁷ The coefficient of variation for survival probability, $CV(\phi)$.

⁸ Recapture probability (p) presented as the maximum likelihood estimate (standard error of the estimate).

⁹ The proportion of residents among newly captured adults (τ) presented as the maximum likelihood estimate (standard error of the estimate).

‡ The estimate for survival probability should be viewed with caution because it is based on fewer than five between-year recaptures, or the estimate is very imprecise ($SE(\phi) > 0.200$ or $CV(\phi) > 50.0\%$), or the proportion of residents is equal to zero.

† The estimate for recapture probability (and possibly survival probability as well) may be biased low because the estimate for τ was 1.000.

Appendix I. Numerical listing (in AOU checklist order) of all the species sequence numbers, species alpha codes, and species names for all species banded or encountered during the 13 years, 1995-2007, of the MAPS Program on the seven stations ever operated on **U.S. Army Fort Bragg**.

Cumulative breeding status for all years in which each station was operated are also included (B = Regular Breeder (all years); U = Usual Breeder (>½, not all, years); O = Occasional Breeder (≤½ years); T = Transient; M = Migrant; A= Altitudinal Disperser; ? = Uncertain Species ID

NUMB	SPEC	SPECIES NAME	Sandstone Hill (SAHI)	I 104 (I104)	I 113 (I113)	S 110 (S110)	S 114 (S114)	S 112 (S112)	I 102 (I102)
00860	DCCO	Double-crested Cormorant		T					
00950	AMBI	American Bittern		T					
01010	GBHE	Great Blue Heron	T	T	T	T		T	T
01040	GREG	Great Egret						T	
01130	GRHE	Green Heron		T					
01290	BLVU	Black Vulture		T		T			
01300	TUVU	Turkey Vulture	T	T	T	T	T	O	T
01460	CANG	Canada Goose		T		T		T	
01570	WODU	Wood Duck		U	T	T	T	T	T
01630	MALL	Mallard		T					
02200	SSHA	Sharp-shinned Hawk		M		M			
02210	COHA	Cooper's Hawk				T			T
02380	RSHA	Red-shouldered Hawk		T	T	T	O	O	
02400	BWHA	Broad-winged Hawk		T	T		T	T	
02460	RTHA	Red-tailed Hawk		T	T	T	T	O	T
02630	AMKE	American Kestrel	O	T	T	T	T	T	U
03040	WITU	Wild Turkey			T		T		
03160	NOBO	Northern Bobwhite	B	B	U	U	U	U	B
04490	AMWO	American Woodcock		T	T	T			
05570	MODO	Mourning Dove	B	U	B	B	U	B	B
06410	YBCU	Yellow-billed Cuckoo	O	U	U	U	U	U	O
06680	EASO	Eastern Screech-Owl		T	O	T	T	T	T
06800	GHOW	Great Horned Owl		T	T				
07080	CONI	Common Nighthawk	O	U	U	O	O	O	U
07170	CWWI	Chuck-will's-widow	T	O		O	T	O	
07230	WPWI	Whip-poor-will		T	O	O		T	
07400	CHSW	Chimney Swift	T	O	T	T	T	O	T
08630	RTHU	Ruby-throated Hummingbird	O	O	U	U	O	U	O
09110	BEKI	Belted Kingfisher	T	O				T	T
09420	RHWO	Red-headed Woodpecker	O	U	U	O	O	O	U
09550	RBWO	Red-bellied Woodpecker	O	B	U	U	U	U	B
09650	DOWO	Downy Woodpecker	O	U	U	U	U	U	O
09660	HAWO	Hairy Woodpecker		T	T	T	O	U	O

Appendix I. Continued.

NUMB	SPEC	SPECIES NAME	SAHI	I104	I113	S110	S114	S112	I102
09680	RCWO	Red-cockaded Woodpecker	U	O	U	O	T	T	B
09800	YSFL	Yellow-shafted Flicker	B	U	U	U	U	U	B
09860	PIWO	Pileated Woodpecker	O	O	O	O	U	U	U
11390	EAWP	Eastern Wood-Pewee	O	B	B	U	O	O	B
11460	ACFL	Acadian Flycatcher		T	T	T	O	O	T
11595	UEFL	Unidentified Empidonax Flycatcher				?			
11610	EAPH	Eastern Phoebe		T					T
11760	GCFL	Great Crested Flycatcher	B	B	B	B	U	B	B
12030	EAKI	Eastern Kingbird	T	O	T	O	T	T	
12520	LOSH	Loggerhead Shrike	T						
12550	WEVI	White-eyed Vireo	T	U	O	U	O	O	O
12690	YTVI	Yellow-throated Vireo	T	O	O	O	O	O	
12720	BHVI	Blue-headed Vireo						T	T
12790	REVI	Red-eyed Vireo	T	O	U	U	U	B	T
12930	BLJA	Blue Jay	B	U	U	U	U	B	B
13190	AMCR	American Crow	U	O	O	O	O	U	O
13270	FICR	Fish Crow	B	O	O	O	O	T	O
13340	PUMA	Purple Martin	T	T	T	T	T	T	T
13410	TRES	Tree Swallow	M					M	M
13540	BARS	Barn Swallow			T	T	T	T	T
13560	CACH	Carolina Chickadee	B	B	B	B	B	B	B
13660	TUTI	Tufted Titmouse	B	B	B	B	B	B	B
13700	WBNU	White-breasted Nuthatch	U	U	U	O	U	U	U
13720	BHNU	Brown-headed Nuthatch	B	U	U	U	T	O	U
14000	CARW	Carolina Wren	O	B	B	B	B	B	B
14350	BGGN	Blue-gray Gnatcatcher	U	B	U	B	U	B	U
14560	EABL	Eastern Bluebird	U	O	O	O	T	O	U
14780	VEER	Veery				M		M	M
14790	GCTH	Gray-cheeked Thrush		M		M			
14810	SWTH	Swainson's Thrush		M		M		M	M
14830	WOTH	Wood Thrush		T	T	T	O	B	
15000	AMRO	American Robin	T	O	O	O	O	O	U
15130	GRCA	Gray Catbird		U	O	O	O	O	U
15150	NOMO	Northern Mockingbird		T	T				
15200	BRTH	Brown Thrasher	O	U	O	U	O	O	U
15550	CEDW	Cedar Waxwing	M	M		M			
15630	BWWA	Blue-winged Warbler		M					
15730	NOPA	Northern Parula		T	T	T	T	T	T
15750	YWAR	Yellow Warbler			M			M	
15770	MAWA	Magnolia Warbler		M		M	M	M	M

Appendix I. Continued.

NUMB	SPEC	SPECIES NAME	SAHI	I104	I113	S110	S114	S112	I102
15790	BTBW	Black-throated Blue Warbler		M	M	M	M	M	M
15830	BTNW	Black-throated Green Warbler		M				M	
15870	YTWA	Yellow-throated Warbler	T	O	T	O	O	O	T
15910	PIWA	Pine Warbler	B	B	B	B	U	U	B
15930	PRAW	Prairie Warbler	B	B	U	B	O	O	B
15970	BLPW	Blackpoll Warbler		M				M	M
16030	BAWW	Black-and-white Warbler		T	O	O	O	U	
16040	AMRE	American Redstart		O	T	T	T	T	T
16050	PROW	Prothonotary Warbler		T	T		O	T	T
16060	WEWA	Worm-eating Warbler			T			T	
16080	OVEN	Ovenbird	T	O	O	O	U	U	T
16090	NOWA	Northern Waterthrush		M	M		M	M	
16100	LOWA	Louisiana Waterthrush		T		O	T	T	
16110	KEWA	Kentucky Warbler				T	U	O	T
16150	COYE	Common Yellowthroat	T	B	B	B	U	U	B
16280	HOWA	Hooded Warbler		O	O	U	B	U	U
16290	WIWA	Wilson's Warbler		M					
16300	CAWA	Canada Warbler					M		
16460	YBCH	Yellow-breasted Chat		O	O	T	T	T	T
16820	SUTA	Summer Tanager	B	B	B	B	U	B	U
17820	EATO	Eastern Towhee	B	B	B	B	U	B	B
17930	BACS	Bachman's Sparrow	B	U	U	U		O	U
18020	CHSP	Chipping Sparrow	U	U	U	U	O	T	U
18050	FISP	Field Sparrow	O		T	O			T
18080	VESP	Vesper Sparrow		T					
18270	WTSP	White-throated Sparrow		M					M
18560	NOCA	Northern Cardinal	O	B	B	B	B	B	B
18640	BLGR	Blue Grosbeak	U	O	O	O	T	O	T
18670	INBU	Indigo Bunting	O	U	U	B	U	B	U
18730	RWBL	Red-winged Blackbird		O					
18800	EAME	Eastern Meadowlark			T				
18870	COGR	Common Grackle	T	O	T	T		T	T
18960	BHCO	Brown-headed Cowbird	O	O	O	U	O	U	U
19040	OROR	Orchard Oriole							T
19160	BAOR	Baltimore Oriole						M	
19370	HOFI	House Finch	T			T		O	
19510	AMGO	American Goldfinch	U	B	U	U	U	B	U
19920	HOSP	House Sparrow				T	T		