



THE INSTITUTE FOR  
BIRD POPULATIONS




## The 2008 Annual Report of the Monitoring Avian Productivity and Survivorship (MAPS) Program in Yosemite National Park

Rodney B. Siegel<sup>1</sup>, Peter Pyle<sup>1</sup>, Danielle R. Kaschube<sup>1</sup>, and Sarah Stock<sup>2</sup>

<sup>1</sup>The Institute for Bird Populations  
P.O. Box 1346  
Point Reyes Station, CA 94956-1346

<sup>2</sup>Division of Resources Management and Science  
Yosemite National Park

February 26, 2008





## TABLE OF CONTENTS

<b>INTRODUCTION</b> .....	2
<b>METHODS</b> .....	4
<b>Establishment and Operation of Stations</b> .....	4
<b>Data Collection</b> .....	5
<b>Computer Data Entry and Verification</b> .....	6
<b>Data Analysis</b> .....	6
Adult population index and productivity analyses.....	7
Analyses of trends in adult population size and productivity .....	7
<b>RESULTS</b> .....	8
<b>2008 Indices of Adult Population Size and Post-fledging Productivity</b> .....	8
<b>Multi-year Trends in Adult Population Size and Productivity</b> .....	10
<b>DISCUSSION</b> .....	11
<b>ACKNOWLEDGMENTS</b> .....	11
<b>LITERATURE CITED</b> .....	12
 <b>TABLES AND FIGURES</b>	
<b>Table 1. Summary of the 2008 operation of the five MAPS stations in Yosemite National Park</b> .....	14
<b>Table 2. Capture summary for the five individual MAPS stations operated in Yosemite National Park in 2008</b> .....	15
<b>Table 3. Numbers of aged individual birds captured per 600 net-hours and proportion of young in the catch at the five individual MAPS stations operated in Yosemite National Park in 2008</b> .....	18
<b>Figure 1. Locations of ongoing Monitoring Avian Productivity and Survivorship (MAPS) bird banding stations at Yosemite National Park</b> .....	21
<b>Figure 2. Population trends for 25 species and all species pooled at the five currently operating MAPS stations in Yosemite National Park over the 16 years 1993-2008</b> .....	22
<b>Figure 3. Trend in productivity for 25 species and all species pooled at the five currently operating MAPS stations in Yosemite National Park over the 16 years 1993-2008</b> .....	25
 <b>APPENDIX 1. Breeding status classifications for all species banded or encountered during the 19 years, 1990-2008, of the MAPS Program on the six stations ever operated at Yosemite National Park</b> .....	28

## INTRODUCTION

National parks can fulfill vital roles as both refuges for bird species dependent on late successional forest conditions, and to provide reference sites for assessing the effects of land use and land cover changes on bird populations throughout the larger geographic area (Silsbee and Peterson 1991). These changes may result from regional activities such as land conversion and forest management, or from broader-scale processes such as global climate change. Indeed, monitoring vital rates and population trends at 'control' sites in national parks is especially important because the parks are among the few sites in the United States where population trends due to large-scale regional or global change patterns are relatively unconfounded with local changes in land-use practices (Simons et al. 1999).

Landbirds are excellent indicators of environmental change in terrestrial ecosystems, because of their high body temperature, rapid metabolism, and high ecological position on most food webs. Furthermore, their abundance and diversity in virtually all terrestrial habitats, diurnal nature, discrete reproductive seasonality, and intermediate longevity facilitate the monitoring of their population and demographic parameters. An added benefit is that landbird monitoring is often particularly efficient, in the sense that many species can be monitored simultaneously with the same survey protocol, and costs are relatively low. Finally, landbirds hold high and growing public interest (Cordell et al. 1999; Cordell and Herbert 2002) and are perhaps the most visible faunal component of park ecosystems.

Population-trend data on Neotropical migrant birds, while suggesting alarming declines in some species, provide no information on primary demographic parameters (productivity and survivorship). Without demographic information, population-trend data alone provide no means for determining at what point(s) in the life cycles problems are occurring, or to what extent population trends are driven by causal factors that affect birth rates, death rates, or both (DeSante 1995). The lack of such information for migratory birds in particular is an obstacle to effective conservation actions, as it leaves unresolved whether critical problems that drive population declines are occurring primarily on temperate breeding grounds, during migration, or on distant tropical wintering grounds. Lack of data on productivity and survivorship thus impedes the formulation of effective management and conservation strategies to reverse population declines (DeSante 1992).

Environmental factors and management actions affect primary demographic parameters directly and these effects can be observed over a short time period (Temple and Wiens 1989). Because of the buffering effects of floater individuals and density-dependent responses of populations, there may be substantial time lags between changes in primary parameters and resulting changes in population size or density as measured by census or survey methods (DeSante and George 1994). Thus, a population could be in trouble long before this becomes evident from population trend data alone. Perhaps even more importantly, because of the vagility of many bird species, local variation in secondary parameters (e.g., population size or density) may be masked by recruitment from a wider region (George et al. 1992) or accentuated by lack of recruitment from a wider area (DeSante 1990). Local abundance can sometimes be a poor indicator of reproductive success, particularly in habitats that have been modified substantially by humans (Bock and Jones 2004).

In 1989 The Institute for Bird Populations ([IBP](#)) established the Monitoring Avian Productivity and Survivorship ([MAPS](#)) program, a cooperative effort among public agencies, private organizations, and individual bird banders in North America. MAPS has since grown into a continent-wide network of over 500 constant-effort mist-netting and banding stations that provide long-term demographic data on landbirds (DeSante et al. 1995). The design of the MAPS program was patterned after the very successful British Constant Effort Sites (CES) Scheme that has been operated by the British Trust for Ornithology since 1981 (Peach et al. 1996). The MAPS program was endorsed in 1991 by both the Monitoring Working Group of PIF and the USDI Bird Banding Laboratory, and has subsequently has attracted participation from numerous federal agencies, including the National Park Service, Department of Defense, Department of the Navy, Department of the Army, Texas Army National Guard, USDA Forest Service, and US Fish and Wildlife Service.

The MAPS Program is organized to fulfill three sets of goals and objectives: monitoring, research, and management:

*Monitoring goals.* For over 100 target species, including Neotropical-wintering migrants, temperate-wintering migrants, and permanent residents, MAPS provides: (a) annual indices of adult population size and post-fledging productivity from data on the numbers and proportions of young and adult birds captured; and (b) annual estimates of adult population size, adult survival rates, proportions of residents, and recruitment into the adult population from modified Cormack- Jolly-Seber analyses of mark-recapture data on adult birds.

*Research goals.* MAPS identifies and describes: (a) temporal and spatial patterns in these demographic indices and estimates at a variety of spatial scales ranging from the local landscape to the entire continent; and (b) relationships between these patterns and ecological characteristics of the target species, population trends of the target species, station-specific and landscape-level habitat characteristics, and spatially-explicit weather variables.

*Management goals.* MAPS uses these patterns and relationships to: (a) identify thresholds and trigger points to notify appropriate agencies and organizations of the need for further research and/or management actions; (b) determine the proximate demographic cause(s) of population change; (c) suggest management actions and conservation strategies to reverse population declines and maintain stable or increasing populations; and (d) evaluate the effectiveness of management actions and conservation strategies that are implemented.

The MAPS program was established in Yosemite National Park in 1990, and Yosemite now hosts some of the longest-running MAPS stations in the country.

In this report we briefly summarize results of the MAPS program at five stations in Yosemite National Park from 1993 (1998 at the Gin Flat East Meadow station) through 2008. [Siegel et al. \(2007\)](#) present a complete report summarizing data and results through 2006, and additional data from the Hodgdon Meadow station from 1990-1992 are presented in previous reports (e.g., [Pyle et al. 2006](#)). Here we present indices of adult population size and productivity for each station

and for all stations combined through 2008. For selected target species and all species pooled, we present temporal trends in adult population size and productivity. More in-depth analyses, particularly analyses assessing effects of annual weather variation and climate change, are slated to be conducted after the 2010 field season.

## METHODS

### Establishment and Operation of Stations

Five MAPS stations were re-established and operated in Yosemite National Park in 2008, at the same locations they were operated in previous years (Fig 1). The five stations, located along an elevation gradient from highest to lowest, were:

- White Wolf Meadow (WHWO), set in a wet montane meadow surrounded by mixed red fir and lodgepole pine forest at 2,402 m elevation.
- Gin Flat East Meadow (GFEM), located in a wet montane meadow surrounded by mixed red fir and lodgepole pine forest at 2,073 m elevation.
- Crane Flat Meadow (CRFL), located in a wet montane meadow with willow and aspen thickets, surrounded by mixed conifer forest at 1,875 m elevation.
- Hodgdon Meadow (HODG), located in a wet montane meadow with willow and dogwood thickets, surrounded by mixed conifer forest and a patch of California Black Oak woodland at 1,408 m elevation.
- Big Meadow (BIME), located in riparian willows and mixed conifer forest (largely consumed by a stand-replacing fire in 1990) in an open, dry meadow at 1,311 m elevation.

The Hodgdon Meadow station was established and first operated in 1990, followed by White Wolf, Crane Flat, and Big Meadow in 1993, and Gin Flat East Meadow in 1998. See Table 1 for details of habitats and operation of each station in 2008.

Through the efforts of three IBP field biologist interns (Jeff Moker, Bucky Ballou, and Craig Ballou), trained and supervised by crew leader Marika Bourdeaux, these five MAPS banding stations were operated during 2008 in accordance with the standardized banding protocols developed for the MAPS Program throughout North America ([DeSante et al. 2006](#)).

Ten net sites (14 sites at the Hodgdon Meadow station) were re-established at each of the stations in 2008, at the exact same locations where they were established and operated in each of the preceding years. One 12-m-long, 30-mm-mesh, nylon mist net was erected at each of the ten net sites at four of the stations on each day of operation. At Hodgdon Meadow, seven of the 14 net sites were operated on one day with the remaining seven net sites operated on a second day. Each of the stations was operated for six morning hours per day (beginning at about local sunrise) during one day (two days for Hodgdon Meadow) in each of eight consecutive 10-day

periods between May 21 and August 4 or, for the two higher-elevation stations (White Wolf Meadow and Gin Flat East Meadow), for one day in each of seven periods between June 5 and August 5 (Table 1). The operation of all stations occurred on schedule in 2008 during each of the ten-day periods.

## Data Collection

With few exceptions, all birds captured at MAPS stations were identified to species, age, and sex. If unbanded, the birds were banded with USGS/BRD numbered aluminum bands. Birds were released immediately upon capture and before being banded or processed if situations arose where bird safety was compromised. Such situations could involve exceptionally large numbers of birds being captured at once, or the sudden onset of adverse weather conditions such as high winds or rainfall. The following data were collected from all birds captured, including recaptures:

- capture code (newly banded, recaptured, band changed, unbanded);
- band number
- species
- age and how aged
- sex (if possible) and how sexed (if applicable)
- extent of skull pneumaticization
- breeding condition of adults (i.e., extent of cloacal protuberance or brood patch)
- extent of juvenal plumage in young birds
- extent of body and flight-feather molt
- extent of primary-feather wear
- presence of molt limits and plumage characteristics
- wing chord
- fat class and body mass
- date and time of capture (net-run time)
- station and net site where captured
- any pertinent notes

Effort data (i.e., the number and timing of net-hours on each day of operation) were also collected in a standardized manner. In order to allow constant-effort comparisons of data, the times of opening and closing the array of mist nets and of beginning each net check were recorded to the nearest ten minutes. The breeding (summer residency) status (confirmed breeder, likely breeder, non-breeder) of each species seen, heard, or captured at each MAPS station on each day of operation was recorded using techniques similar to those employed for breeding bird atlas projects.

For each of the five stations, simple habitat maps prepared in previous years (indicating extent and location of major habitats, as well as structures, roads, trails, and streams) were checked and updated where necessary. The pattern and extent of cover of each of four major vertical layers of vegetation (upperstory, midstory, understory, and ground cover), in each major habitat type, were

classified into one of twelve pattern types and eleven cover categories according to guidelines in the MAPS Habitat Structure Assessment Protocol ([Nott et al. 2003](#)).

### **Computer Data Entry and Verification**

The computer entry of all banding data was completed by John W. Shipman of Zoological Data Processing, Socorro, NM. The critical data for each banding record (capture code, band number, species, age, sex, date, capture time, station, and net number) were proofed by hand against the raw data and any computer-entry errors were corrected. Computer entry of effort and vegetation data was completed by IBP biologists using custom data entry programs. All banding data were then run through a series of verification programs as follows:

- Clean-up programs to check the validity of all codes entered and the ranges of all numerical data.
- Cross-check programs to compare station, date, and net fields from the banding data with those from the summary of mist netting effort data.
- Cross-check programs to compare species, age, and sex determinations against degree of skull pneumaticization, breeding condition (extent of cloacal protuberance and brood patch), and extent of body and flight-feather molt, primary-feather wear, and juvenal plumage.
- Screening programs which allow identification of unusual or duplicate band numbers or unusual band sizes for each species.
- Verification programs to screen banding and recapture data from all years of operation for inconsistent species, age, or sex determinations for each band number.

Any discrepancies or suspicious data identified by any of these programs were examined manually and corrected if necessary. Wing chord, weight, station of capture, date, and any pertinent notes were used as supplementary information for the correct determination of species, age, and sex in all of these verification processes.

### **Data Analysis**

We classified the landbird species captured in mist nets into six groups based upon their breeding or summer residency status. Each species was classified as one of the following:

- a regular breeder (B) if we had positive or probable evidence of breeding or summer residency within the boundaries of the MAPS station *during all years* that the station was operated.
- a usual breeder (U) if we had positive or probable evidence of breeding or summer residency within the boundaries of the MAPS station *during more than half but not all of the years* that the station was operated.



- an occasional breeder (O) if we had positive or probable evidence of breeding or summer residency within the boundaries of the MAPS station *during half or fewer of the years* that the station was operated.
- a transient (T) if the species was *never* a breeder or summer resident at the station, but the station was within the overall breeding range of the species.
- an altitudinal disperser (A) if the species breeds only at lower elevation than that of the station but disperses to higher elevations after breeding.
- a migrant (M) if the station was not located within the overall breeding range of the species.

Data for a given species from a given station were included in productivity analyses if the station was within the breeding range of the species; that is, data were included from stations where the species was a breeder (B, U, or O), or transient (T), but not where the species was an altitudinal disperser (A) or a migrant (M). Data for a given species from a given station were included in trend analyses only if the species was classified as a regular (B) or usual (U) breeder at the station. Throughout this report we define “target species” for trend and survivorship analyses as those for which an average of 2.5 individual adult birds were captured per year at all stations combined or at each station for station-specific analysis.

#### Adult population index and productivity analyses

The proofed, verified, and corrected banding data from all sixteen years were run through a series of analysis programs that calculated for each species:

- the numbers of newly banded birds, recaptured birds, and birds released unbanded.
- the numbers and capture rates (per 600 net-hours) of first captures (in a given year) of individual adult and young birds.
- the reproductive index. Following the procedures pioneered by the British Trust for Ornithology (BTO) in their CES Scheme (Peach et al. 1996), we used the number of adult birds captured as an index of adult population size. For each species each year, we calculated a yearly reproductive index as the number of young divided by the number of adults.

#### Analyses of trends in adult population size and productivity

For each target species and for all species pooled we examined 16-year trends in adult population size and productivity (reproductive index) using data from all five stations combined. Year-to-year comparisons were made in a “constant-effort” manner by means of an analysis program that used actual net-run (capture) times and net-opening and -closing times on a net-by-net and period-by-period basis. We excluded captures that occurred in a given net in a

given period in one year during the time when that net was not operated in that period in the other year. For trends in population size, we first calculated adult population indices for each species for each of the 16 years based on an arbitrary starting index of 1.0 in the first year of station operation or analysis (1993). The constant-effort changes were used to calculate these “chain” indices in each subsequent year by multiplying the proportional change (percent change divided by 100) between the two years times the index of the previous year and adding that figure to the index of the previous year:

$$PSI_{i+1} = PSI_i + PSI_i * (d_i/100),$$

where  $PSI_i$  is the population size index for year  $i$  and  $d_i$  is the percentage change in constant-effort numbers from year  $i$  to year  $i+1$ . A regression analysis was then run to determine the slope ( $PT$ ) of these indices. Because the indices for adult population size are based on percentage changes, we further calculated the annual percent change ( $APC$ ), defined as the average change per year, to provide an estimate of the population trend for the species;  $APC$  was calculated as:

$$(\text{slope of the regression line} / \text{predicted value for the first year}) * 100$$

We present the  $APC$ , the standard error of the slope ( $SE$ ), the correlation coefficient ( $r$ ), and the significance of the correlation ( $P$ ) to describe each trend. For 16-year trends, species for which  $r \geq 0.30$  are considered to have a substantially increasing trend, those for which  $r \leq -0.30$  are considered to have a substantially decreasing trend, those for which absolute  $r < 0.3$  and  $SE \leq 0.015$  are considered to have a non-substantial and non-fluctuating trend, and those for which absolute  $r \leq 0.3$  and  $SE > 0.015$  are considered to have non-substantial, widely fluctuating trends.

Trends in productivity,  $PrT$ , for all stations combined were calculated in an analogous manner by starting with actual productivity values in 1993 and calculating each successive year's value based on the actual constant-effort changes in productivity between each pair of consecutive years. For trends in productivity, the slope ( $PrT$ ) and its standard error ( $SE$ ) are presented, along with the correlation coefficient ( $r$ ), and the significance of the correlation ( $P$ ). Productivity trends are characterized in a manner analogous to that for population trends, except that, for non-substantial trends, we do not attempt to distinguish between those that are widely fluctuating and those that are non-fluctuating.

## RESULTS

A total of 2,147.3 net-hours was accumulated at the five MAPS stations operated in Yosemite National Park in 2008 (Table 1). Data from 1,987.7 of these net-hours could be compared directly to the previous year's data in a constant-effort manner.

### 2008 Indices of Adult Population Size and Post-fledging Productivity

We present the 2008 numbers of newly-banded, unbanded, and recaptured birds for each species at each of the five stations individually and for all stations combined in Table 2. A total of 2,663 captures of 66 species was recorded during the summer of 2008. Newly banded birds comprised

75.3% of the total captures. The greatest number of total captures (827) was recorded at the Crane Flat station and the smallest number of total captures (242) was recorded at the Big Meadow station. The highest species richness occurred at Hodgdon Meadow (43 species) and the lowest species richness occurred at White Wolf Meadow (29 species).

The 2008 capture rates (per 600 net-hours) of individual adult and young birds and the 2008 reproductive index (number of young birds per adult) are presented for each species and for all species pooled at each station and all stations combined in Table 3. We present capture rates (captures per 600 net-hours) rather than absolute numbers of birds in this table so that the data can be compared among stations which, because of the vagaries of weather and accidental net damage, can differ from one another in effort expended (see Table 1). These capture indices suggest that the total adult population size in 2008 was greatest at Crane Flat (293.7 adults/600 net-hours), followed in descending order by Hodgdon Meadow, Gin Flat East Meadow, Big Meadow, and White Wolf Meadow (Table 3). The capture rate of young of all species pooled at each station in 2008 was highest at Crane Flat (388.8 young/600 net-hours), followed by Gin Flat East Meadow, White Wolf Meadow, Hodgdon Meadow, and Big Meadow (Table 3).

Reproductive index (the number of young per adult) at the five stations in 2008 was greatest at Gin Flat East Meadow (1.73), followed by White Wolf Meadow (1.64), Crane Flat (1.32), Big Meadow (0.98), and Hodgdon Meadow (0.61). The mean adult capture rate for the five stations combined was 214.3 per 600 net hours and the overall reproductive index was 1.14 in 2008.

In 2008 Orange-crowned Warbler was the most frequently captured, followed by Dark-eyed Junco, MacGillivray's Warbler, Lincoln's Sparrow, Yellow-rumped Warbler, Song Sparrow, Nashville Warbler, Golden-crowned Kinglet, Anna's Hummingbird, Hermit Warbler, and Mountain Chickadee (Table 2). Overall, the most abundant species in 2008 (as determined by the number of adults captured per 600 net-hours; Table 3) for which the five Yosemite National Park MAPS stations are within the breeding range of the species (not including Orange-crowned Warbler because it doesn't breed in Yosemite and Anna's Hummingbird because birds were not banded to determine individual adults), in decreasing order, were MacGillivray's Warbler, Dark-eyed Junco, Lincoln's Sparrow, Yellow-rumped Warbler, Song Sparrow, Mountain Chickadee, Western Tanager, and American Robin. The following is a list of such species (captured at a rate of at least 8.0 adults per 600 net-hours), in decreasing order, at each station in 2008 (Table 3):

**White Wolf Meadow**

Dark-eyed Junco  
Yellow-rumped Warbler  
American Robin  
Lincoln's Sparrow  
Mountain Chickadee  
Dusky Flycatcher

**Crane Flat**

Yellow-rumped Warbler  
Lincoln's Sparrow  
Dark-eyed Junco  
MacGillivray's Warbler  
Warbling Vireo  
Dusky Flycatcher  
Mountain Chickadee  
American Robin  
Hermit Warbler  
Red-breasted Nuthatch  
Brown Creeper

**Big Meadow**

Bushtit  
Lazuli Bunting  
Purple Finch  
Wrentit  
MacGillivray's Warbler  
Spotted Towhee  
Black-headed Grosbeak  
Hammond's Flycatcher  
Nashville Warbler

**Gin Flat East Meadow**

Lincoln's Sparrow  
Dark-eyed Junco  
Dusky Flycatcher  
Yellow-rumped Warbler  
MacGillivray's Warbler  
Dusky Flycatcher  
Mountain Chickadee  
Red-breasted Sapsucker

**Hodgdon Meadow**

MacGillivray's Warbler  
Song Sparrow  
Warbling Vireo  
Western Tanager  
Black-headed Grosbeak  
Lincoln's Sparrow  
Dark-eyed Junco  
Red-breasted Nuthatch  
American Robin

**Multi-year Trends in Adult Population Size and Productivity**

“Chain” indices of adult population size for the 16-year period 1993-2008 are presented for 25 target species and for all species pooled in Figure 2. We used annual percent change (*APC*) for each species as an estimate of the mean annual population trend for that species. These estimates of *APC*, along with the standard error of the slope (in parentheses), the correlation coefficient (*r*), and the significance of the correlation (*P*), are included for each target species and for all species pooled on each graph.

Populations of 12 species as well as all species pooled showed substantial declining trends ( $r \leq -0.3$  for a 16-year trend). The declines for Golden-crowned Kinglet, Hermit Warbler, Chipping Sparrow, and Lazuli Bunting were highly significant; those for Western Wood-Pewee, Dusky Flycatcher, Yellow Warbler, Dark-eyed Junco, and Purple Finch were significant; that of Warbling Vireo was nearly significant, and those of Hermit Thrush, Black-headed Grosbeak, and all species pooled were not significant. In contrast, populations of only five species showed substantial increasing trends ( $r \geq 0.3$ ), which were highly significant for Mountain Chickadee, MacGillivray's Warbler, and Western Tanager, and nearly significant for Yellow-rumped Warbler and Song Sparrow. Populations of the remaining eight target species (Red-breasted Sapsucker, Hammond's Flycatcher, Cassin's Vireo, Brown Creeper, American Robin, Lincoln's Sparrow, Pine Siskin, and Lesser Goldfinch) showed non-substantial (absolute  $r < 0.3$ ) trends. Seven of these eight species showed substantially fluctuating (SE of the slope  $\geq 0.015$ ) population trends, whereas only one species (Lincoln's Sparrow) showed a non-fluctuating trend. Overall, 16 of the 25 species showed negative trends, one species (American Robin) showed a flat trend, and eight species showed positive trends. The substantial negative trend for all species pooled had been highly significant ( $P = 0.009$ ) after 14 years of data had been collected (Siegel et al. 2007) but based on increasing populations in 2006-2008 was no longer significant after 16 years of data had been collected. Nevertheless, the decline continues to be substantial at a decreasing rate of -0.9% per year, suggesting that total populations of landbirds in Yosemite have declined by 14.5% over the 16-year period (1993-2008).

“Chain” indices of productivity for each of the 16 years (1993-2008) are shown in Figure 3 for the same 25 target species and all species pooled, at all five stations combined. Five species showed substantially declining productivity trends ( $r \leq -0.30$ ), which were highly significant for Lesser Goldfinch, significant for Hermit Thrush, marginally significant for Hammond's

Flycatcher and MacGillivray's Warbler, and not significant for Western Wood-Pewee. In contrast, 11 species as well as all species pooled showed substantially increasing productivity trends ( $r \geq 0.30$ ); these were highly significant for Red-breasted Sapsucker, American Robin, Yellow Warbler, Black-headed Grosbeak, and Lazuli Bunting; significant for Mountain Chickadee, Lincoln's Sparrow, and Purple Finch; nearly significant for Golden-crowned Kinglet; and not significant for Warbling Vireo, Hermit Warbler, and all species pooled. The remaining nine species (Dusky Flycatcher, Cassin's Vireo, Brown Creeper, Yellow-rumped Warbler, Western Tanager, Chipping Sparrow, Song Sparrow, Dark-eyed Junco, and Pine Siskin) showed non-substantial productivity trends. Overall, 15 of the 25 target species had positive productivity trends and ten had negative productivity trends. The productivity trend for all species pooled indicated an average annual increase of 0.027 per year.

## DISCUSSION

The MAPS Program in Yosemite continues to yield station-specific indices of adult population size and post-fledging productivity, park-wide estimates of annual survival rates of adults, and important information on annual changes and longer-term trends in these indices and estimates, for over 25 target species. The results in this and previous reports underscore the complexity of the population dynamics of Yosemite's breeding birds, complexity which can only be unraveled through long-term data collection.

The Yosemite MAPS Program also continues to yield both new findings and new hypotheses about landbird population dynamics in the park. The generation of preliminary findings and new hypotheses that can then be followed up with targeted research is one of the hallmarks of an effective ecological monitoring program. A recent example of this monitoring-research cycle is that MAPS results over the past two decades suggested disturbing declines in the park's Willow Flycatcher population, prompting an intensive research project on Willow Flycatcher in the park (Siegel et al. 2008). We believe that the MAPS program at Yosemite is particularly well-suited to studying the effects of annual weather variation and climate change on birds, and this will be an important focus of our next major analytical report for the Yosemite MAPS program, which is slated to be completed after the 2010 field season.

## ACKNOWLEDGMENTS

All data presented in this report were collected by field biologist interns of The Institute for Bird Populations (IBP). In 2008, they were Jeff Moker, Bucky Ballou, and Craig Ballou, who were trained and supervised by crew leader Marika Bourdeaux. We thank the field crew for their excellent work. We thank Steve Thompson of Yosemite National Park for his support of and assistance with much of the logistical and administrative aspects of this work. Financial support for the MAPS Program in Yosemite National Park during 2008 was provided by the National Park Service through a Cooperative Agreement between Yosemite National Park and The Institute for Bird Populations, with funding provided by the Yosemite Fund. This is Contribution Number 362 of The Institute for Bird Populations.

## LITERATURE CITED

- Bock, C. E. and Z. F. Jones. 2004. Avian habitat evaluation: should counting birds count? *Frontiers in Ecology and Environment* 2:403-410.
- Cordell, H. K. and N. G. Herbert. 2002. The popularity of birding is still growing. *Birding* Feb.:54-61.
- Cordell, H. K., N. G. Herbert, and F. Pandolfi. 1999. The growing popularity of birding in the United States. *Birding* 3:168-176.
- DeSante, D. F. 1990. The role of recruitment in the dynamics of a Sierran subalpine bird community. *American Naturalist* 136:429-455.
- DeSante, D. F. 1992. Monitoring Avian Productivity and Survivorship (MAPS): a sharp, rather than blunt, tool for monitoring and assessing landbird populations. *In*: D. R. McCullough and R. H. Barrett (Eds.), *Wildlife 2001: Populations*, pp. 511-521. (London, U.K.: Elsevier Applied Science).
- DeSante, D. F. 1995. Suggestions for future directions for studies of marked migratory Landbirds from the perspective of a practitioner in population management and conservation. *Journal of Applied Statistics* 22:949-965.
- DeSante, D. F., K. M. Burton, J. F. Saracco, and B. L. Walker. 1995. Productivity indices and survival rate estimates from MAPS, a continent-wide programme of constant-effort mist netting in North America. *Journal of Applied Statistics* 22:935-947.
- [DeSante, D. F., K. M. Burton, P. Velez, D. Froehlich, and D. R. Kaschube. 2006. MAPS manual. The Institute for Bird Populations, Point Reyes Station, CA.](#)
- DeSante, D. F., and T. L. George. 1994. Population trends in the landbirds of western North America, *In*: J. R. Jehl, Jr. and N. K. Johnson (Eds.), *A Century of Avifaunal Change in Western North America*, *Studies in Avian Biology*, No. 15, pp. 173-190 (Cooper Ornithological Society).
- George, T. L., A. C. Fowler, R. L. Knight, and L.C. McEwen. 1992. Impacts of a severe drought on grassland birds in western North America. *Ecological Applications* 2:275-284.
- Nott, M. P., D. F. DeSante, and N. Michel. 2003. Monitoring Avian Productivity and Survivorship (MAPS) Habitat Structure Assessment Protocol. The Institute for Bird Populations, Pt. Reyes Station, CA.
- Peach, W. J., S. T. Buckland, and S. R. Baillie. 1996. The use of constant effort mist-netting to measure between-year changes in the abundance and productivity of common passerines. *Bird Study* 43:142-156.

[Pyle, P., D. R. Kaschube, R. B. Siegel, and D. F. DeSante. 2006.](#) The 2005 annual report of the Monitoring Avian Productivity and Survivorship (MAPS) Program in Yosemite National Park. The Institute for Bird Populations, Point Reyes Station, CA.

Siegel, R. B., R. L. Wilkerson, and D. F. DeSante. 2008. Extirpation of the Willow Flycatcher from Yosemite National Park. *Western Birds* 39:8-21.

[Siegel, R. S., P. Pyle, D. Kaschube, and D. F. Desante. 2007.](#) The 2006 Annual Report of the Monitoring Avian Productivity and Survivorship (MAPS) Program in Yosemite National Park. The Institute for Bird Populations, Point Reyes Station, CA 94956.

Silsbee, G. G. and D. L. Peterson. 1991. Designing and implementing comprehensive long-term inventory and monitoring programs for National Park System lands. Natural Resources Report NPS/NRUW/NRR-91/04, Denver, CO.

Simons, T. R., K. N. Rabenold, D. A. Buehler, J. A. Collazo, and K. E. Fransreb. 1999. The role of indicator species: neotropical migratory song birds. Pages 187-208 *in* J. D. Peine, (ed.), *Ecosystem management for sustainability: principles and practices illustrated by a regional biosphere reserve cooperative*. Lewis Publishers. New York.

Temple, S. A., and J. A. Wiens. 1989. Bird populations and environmental changes: can birds be bio-indicators? *American Birds* 43:260-270.

Table 1. Summary of the 2008 operation of the five MAPS stations in Yosemite National Park.

Station		No.	Major Habitat Type	Latitude-longitude	Avg Elev. (m)	2008 operation		
Name	Code					Total number of net-hours <sup>1</sup>	No. of periods	Inclusive dates
White Wolf Meadow	WHWO	11904	Wet montane meadow, red fir/ lodgepole pine forest	37°52'10"N,-119°39'08"W	2402	367.0 (333.6)	7	6/14 - 8/05
Gin Flat East Meadow	GFEM	11980	Wet montane meadow, mixed fir forest	37°45'59"N,-119°45'37"W	2073	382.3 (357.3)	7	6/05 - 7/31
Crane Flat	CRFL	11907	Wet montane meadow, willow/ aspen thickets, mixed coniferous forest	37°45'20"N,-119°48'13"W	1875	429.0 (403.2)	8	5/27 - 8/01
Hodgdon Meadow	HODG	11107	Wet montane meadow, willow/ dogwood thickets, mixed oak and coniferous forest	37°47'41"N,-119°51'50"W	1408	595.3 (561.8)	8	5/21 - 8/03
Big Meadow	BIME	11905	Riparian willows, mixed coniferous forest (largely consumed by a stand- replacing fire in 1990), open dry meadow	37°42'16"N,-119°45'07"W	1311	373.7 (331.7)	8	5/23 - 8/04
ALL STATIONS COMBINED						2147.3 (1987.7)	8	5/21 - 8/05

<sup>1</sup> Total net-hours in 2008. Net-hours in 2008 that could be compared in a constant-effort manner to 2007 are shown in parentheses.



Table 2. Capture summary for the five individual MAPS stations operated in Yosemite National Park in 2008. N = Newly Banded, U = Unbanded, R = Recaptures of banded birds.

Species	White Wolf			Gin Flat East Meadow			Crane Flat			Hodgdon Meadow			Big Meadow			All five stations combined			
	N	U	R	N	U	R	N	U	R	N	U	R	N	U	R	N	U	R	
Mountain Quail														1			1		
California Quail														2			2		
Anna's Hummingbird					3			2			55			16			76		
Calliope Hummingbird		1						4			4						9		
Rufous Hummingbird		4			5			6			6			2			23		
Unident. Selasphorus Hum.		1			1												2		
Williamson's Sapsucker	3					1										3		1	
Red-breasted Sapsucker				7		1	4		4	10	1	3	3			24	1	8	
Nuttall's Woodpecker													1			1			
Downy Woodpecker										4						4			
Hairy Woodpecker				1			1			1						3			
White-headed Woodpecker				1			2		2	2						5		2	
Northern Flicker	2			1			1									4			
Olive-sided Flycatcher										2						2			
Western Wood-Pewee				1						8			6		1	15		1	
Willow Flycatcher											1						1		
Hammond's Flycatcher	3			5			9		2	2						19		2	
Dusky Flycatcher	6		1	13			9		4	8			2			38		5	
Pacific-slope Flycatcher				5	1		1			13		1	1			20	1	1	
Unidentified Empidonax Flyc.								1									1		
Black Phoebe							1						11		1	12		1	
Ash-throated Flycatcher													1			1			
Cassin's Vireo							7		1	8		1	3		2	18		4	
Warbling Vireo							15		5	23		4				38		9	
Steller's Jay	1			2						3			1			7			
Western Scrub-Jay													1		1	1		1	
Mountain Chickadee	13		7	18		1	19		1	7		5				57		14	

Table 2. (cont.) Capture summary for the five individual MAPS stations operated in Yosemite National Park in 2008. N = Newly Banded, U = Unbanded, R = Recaptures of banded birds.

Species	White Wolf			Gin Flat East Meadow			Crane Flat			Hodgdon Meadow			Big Meadow			All five stations combined		
	N	U	R	N	U	R	N	U	R	N	U	R	N	U	R	N	U	R
Oak Titmouse													2			2		
Bushtit													26	1	5	26	1	5
Red-breasted Nuthatch				7			38		1	11		2				56		3
White-breasted Nuthatch							2									2		
Brown Creeper	12	1	1	1		2	14	1	2	4		2	2			33	2	7
Bewick's Wren															1			1
House Wren	2		1	2			2			7		3	5	1		18	1	4
Winter Wren										1						1		
Golden-crowned Kinglet	2			37			37			4						80		
Hermit Thrush	2		1				1			2		1				5		2
American Robin	7		2	2			7		1	7	1	2	1			24	1	5
Wrentit													20		12	20		12
Orange-crowned Warbler	56	1		56		1	137	1	11	166		18	6			421	2	30
Nashville Warbler	34	1	2	9			32		2	28	1	7	5		2	108	2	13
Yellow Warbler	2									4			2		2	8		2
Yellow-rumped Warbler	38	1	4	57	1	2	49		10	9						153	2	16
Black-throated Gray Warbler				1			3			3			5			12		
Townsend's Warbler	1															1		
Hermit Warbler	24			8			28		1	13		1				73		2
MacGillivray's Warbler	1			12	1		24		26	62	2	77	11		1	110	3	104
Wilson's Warbler	7						2			6		3				15		3
Western Tanager	3			2			8			16		4	1			30		4
Green-tailed Towhee				2												2		
Spotted Towhee							1			6		6	5	1	6	12	1	12
California Towhee													1			1		
Chipping Sparrow	3						4		3	1		1	1			9		4
Fox Sparrow				2		1										2		1

Table 2. (cont.) Capture summary for the five individual MAPS stations operated in Yosemite National Park in 2008. N = Newly Banded, U = Unbanded, R = Recaptures of banded birds.

Species	White Wolf			Gin Flat East Meadow			Crane Flat			Hodgdon Meadow			Big Meadow			All five stations combined		
	N	U	R	N	U	R	N	U	R	N	U	R	N	U	R	N	U	R
Song Sparrow	1			12		2	28		24	36	1	22	1		1	78	1	49
Lincoln's Sparrow	7		4	22	4	28	25	3	58	12	2	12	1			67	9	102
Dark-eyed Junco	88	3	23	42	1	7	98	4	27	26		5				254	8	62
Black-headed Grosbeak										14		4	13			27		4
Lazuli Bunting				1			5	1		1			14		3	21	1	3
Indigo Bunting							1									1		
Red-winged Blackbird										5		2				5		2
Brewer's Blackbird				5			1									6		
Brown-headed Cowbird													2			2		
Pine Grosbeak	1		1													1		1
Purple Finch				1			1			1			17		3	20		3
Cassin's Finch	4			1			2						1			8		
Pine Siskin	1			6						1						8		
Lesser Goldfinch				7		1							5			12		1
<b>ALL SPECIES POOLED</b>	<b>324</b>	<b>13</b>	<b>47</b>	<b>349</b>	<b>17</b>	<b>47</b>	<b>619</b>	<b>23</b>	<b>185</b>	<b>537</b>	<b>74</b>	<b>186</b>	<b>177</b>	<b>24</b>	<b>41</b>	<b>2006</b>	<b>151</b>	<b>506</b>
Total Number of Captures		384			413			827			797			242			2663	
Number of Species	27	7	11	33	7	11	36	8	19	39	10	23	33	7	14	59	21	40
Total Number of Species		29			36			39			43			38			66	

Table 3. Numbers of aged individual birds captured per 600 net-hours and proportion of young in the catch at the five individual MAPS stations operated in Yosemite National Park in 2008.

Species	White Wolf			Gin Flat East Meadow			Crane Flat			Hodgdon Meadow			Big Meadow			All five stations combined		
	Ad.	Yg.	Prop. Yg.	Ad.	Yg.	Prop. Yg.	Ad.	Yg.	Prop. Yg.	Ad.	Yg.	Prop. Yg.	Ad.	Yg.	Prop. Yg.	Ad.	Yg.	Prop. Yg.
Williamson's Sapsucker	4.9	0.0	0.00	1.6	0.0	0.00										1.1	0.0	0.00
Red-breasted Sapsucker				9.4	3.1	0.33	2.8	4.2	1.50	7.1	6.0	0.86	3.2	1.6	0.50	4.8	3.4	0.71
Nuttall's Woodpecker													0.0	1.6	und.	0.0	0.3	und.
Downy Woodpecker										2.0	2.0	1.00				0.6	0.6	1.00
Hairy Woodpecker				1.6	0.0	0.00	0.0	1.4	und.	1.0	0.0	0.00				0.6	0.3	0.50
White-headed Woodpecker				0.0	1.6	und.	4.2	1.4	0.33	1.0	1.0	1.00				1.1	0.8	0.75
Northern Flicker	3.3	0.0	0.00	1.6	0.0	0.00	1.4	0.0	0.00							1.1	0.0	0.00
Olive-sided Flycatcher										2.0	0.0	0.00				0.6	0.0	0.00
Western Wood-Pewee				0.0	1.6	und.				7.1	1.0	0.14	8.0	3.2	0.40	3.4	1.1	0.33
Hammond's Flycatcher	0.0	4.9	und.	1.6	6.3	4.00	8.4	5.6	0.67	0.0	2.0	und.				2.0	3.6	1.86
Dusky Flycatcher	8.2	1.6	0.20	15.7	3.1	0.20	12.6	1.4	0.11	5.0	3.0	0.60	1.6	1.6	1.00	8.4	2.2	0.27
Pacific-slope Flycatcher				3.1	3.1	1.00	0.0	1.4	und.	7.1	6.0	0.86	1.6	0.0	0.00	2.8	2.5	0.90
Black Phoebe							1.4	0.0	0.00				3.2	11.2	3.50	0.8	2.0	2.33
Ash-throated Flycatcher													1.6	0.0	0.00	0.3	0.0	0.00
Cassin's Vireo							5.6	4.2	0.75	6.0	2.0	0.33	6.4	0.0	0.00	3.9	1.4	0.36
Warbling Vireo							16.8	5.6	0.33	17.1	8.1	0.47				8.1	3.4	0.41
Steller's Jay	1.6	0.0	0.00	3.1	0.0	0.00				3.0	0.0	0.00	1.6	0.0	0.00	2.0	0.0	0.00
Western Scrub-Jay													1.6	1.6	1.00	0.3	0.3	1.00
Mountain Chickadee	9.8	13.1	1.33	11.0	17.3	1.57	11.2	16.8	1.50	7.1	3.0	0.43				7.8	9.5	1.21
Oak Titmouse													1.6	1.6	1.00	0.3	0.3	1.00
Bushtit													20.9	19.3	0.92	3.6	3.4	0.92
Red-breasted Nuthatch				1.6	9.4	6.00	9.8	43.4	4.43	8.1	4.0	0.50				4.5	11.5	2.56
White-breasted Nuthatch							0.0	2.8	und.							0.0	0.6	und.
Brown Creeper	8.2	13.1	1.60	1.6	0.0	0.00	9.8	11.2	1.14	2.0	3.0	1.50	0.0	3.2	und.	4.2	5.9	1.40
Bewick's Wren													1.6	0.0	0.00	0.3	0.0	0.00
House Wren													1.6	6.4	4.00	0.3	1.1	4.00

Table 3. (cont.) Numbers of aged individual birds captured per 600 net-hours and proportion of young in the catch at the five individual MAPS stations operated in Yosemite National Park in 2008.

Species	White Wolf			Gin Flat East Meadow			Crane Flat			Hodgdon Meadow			Big Meadow			All five stations combined		
	Ad.	Yg.	Prop. Yg.	Ad.	Yg.	Prop. Yg.	Ad.	Yg.	Prop. Yg.	Ad.	Yg.	Prop. Yg.	Ad.	Yg.	Prop. Yg.	Ad.	Yg.	Prop. Yg.
Winter Wren										1.0	0.0	0.00				0.3	0.0	0.00
Golden-crowned Kinglet	0.0	3.3	und.	1.6	56.5	36.00	7.0	43.4	6.20	1.0	3.0	3.00				2.0	20.1	10.29
Hermit Thrush	4.9	0.0	0.00				1.4	0.0	0.00	2.0	1.0	0.50				1.7	0.3	0.17
American Robin	11.4	1.6	0.14	3.1	0.0	0.00	11.2	0.0	0.00	8.1	1.0	0.13	0.0	1.6	und.	7.0	0.8	0.12
Wrentit													12.8	25.7	2.00	2.2	4.5	2.00
Nashville Warbler										4.0	24.2	6.00	8.0	1.6	0.20	2.5	7.0	2.78
Yellow Warbler	0.0	3.3	und.							2.0	2.0	1.00	3.2	3.2	1.00	1.1	1.7	1.50
Yellow-rumped Warbler	24.5	44.1	1.80	15.7	75.3	4.80	43.4	28.0	0.65	7.1	2.0	0.29				17.6	27.1	1.54
Black-throated Gray Warbler				0.0	1.6	und.	0.0	4.2	und.	0.0	3.0	und.	0.0	8.0	und.	0.0	3.4	und.
Hermit Warbler	1.6	37.6	23.00	3.1	9.4	3.00	11.2	28.0	2.50	6.0	8.1	1.33				4.8	15.9	3.35
MacGillivray's Warbler	0.0	1.6	und.	14.1	4.7	0.33	28.0	18.2	0.65	57.4	22.2	0.39	9.6	8.0	0.83	25.7	12.3	0.48
Wilson's Warbler	0.0	11.4	und.				1.4	1.4	1.00	4.0	2.0	0.50				1.4	2.8	2.00
Western Tanager	4.9	0.0	0.00	3.1	0.0	0.00	7.0	4.2	0.60	15.1	4.0	0.27	1.6	0.0	0.00	7.3	2.0	0.27
Green-tailed Towhee				3.1	0.0	0.00										0.6	0.0	0.00
Spotted Towhee							1.4	0.0	0.00	4.0	4.0	1.00	9.6	3.2	0.33	3.1	1.7	0.54
California Towhee													0.0	1.6	und.	0.0	0.3	und.
Chipping Sparrow	0.0	4.9	und.				5.6	1.4	0.25	1.0	0.0	0.00	0.0	1.6	und.	1.4	1.4	1.00
Fox Sparrow				4.7	0.0	0.00										0.8	0.0	0.00
Song Sparrow	0.0	1.6	und.	0.0	17.3	und.	5.6	33.6	6.00	24.2	18.1	0.75	1.6	0.0	0.00	8.1	15.1	1.86
Lincoln's Sparrow	11.4	0.0	0.00	34.5	12.6	0.36	40.6	14.0	0.35	12.1	4.0	0.33	0.0	1.6	und.	19.6	6.4	0.33
Dark-eyed Junco	49.0	112.8	2.30	22.0	51.8	2.36	37.8	109.1	2.89	12.1	16.1	1.33				23.2	54.8	2.36
Black-headed Grosbeak										14.1	2.0	0.14	9.6	11.2	1.17	5.6	2.5	0.45
Lazuli Bunting				0.0	1.6	und.	4.2	2.8	0.67	1.0	0.0	0.00	16.1	9.6	0.60	3.9	2.5	0.64
Red-winged Blackbird										7.1	0.0	0.00				2.0	0.0	0.00
Brewer's Blackbird				6.3	1.6	0.25	1.4	0.0	0.00							1.4	0.3	0.20
Brown-headed Cowbird													0.0	3.2	und.	0.0	0.6	und.

Table 3. (cont.) Numbers of aged individual birds captured per 600 net-hours and proportion of young in the catch at the five individual MAPS stations operated in Yosemite National Park in 2008.

Species	White Wolf			Gin Flat East Meadow			Crane Flat			Hodgdon Meadow			Big Meadow			All five stations combined		
	Ad.	Yg.	Prop. Yg.	Ad.	Yg.	Prop. Yg.	Ad.	Yg.	Prop. Yg.	Ad.	Yg.	Prop. Yg.	Ad.	Yg.	Prop. Yg.	Ad.	Yg.	Prop. Yg.
Pine Grosbeak	3.3	0.0	0.00													0.6	0.0	0.00
Purple Finch				0.0	1.6	und.	0.0	1.4	und.	1.0	0.0	0.00	16.1	14.5	0.90	3.1	3.1	1.00
Cassin's Finch	6.5	0.0	0.00	1.6	0.0	0.00	2.8	0.0	0.00				1.6	0.0	0.00	2.2	0.0	0.00
Pine Siskin	1.6	0.0	0.00	4.7	4.7	1.00				1.0	0.0	0.00				1.4	0.8	0.60
Lesser Goldfinch				1.6	11.0	7.00							6.4	1.6	0.25	1.4	2.2	1.60
<b>ALL SPECIES POOLED</b>	<b>155.3</b>	<b>255.0</b>	<b>1.64</b>	<b>171.1</b>	<b>295.0</b>	<b>1.73</b>	<b>293.7</b>	<b>388.8</b>	<b>1.32</b>	<b>261.0</b>	<b>158.2</b>	<b>0.61</b>	<b>150.9</b>	<b>147.7</b>	<b>0.98</b>	<b>214.3</b>	<b>243.4</b>	<b>1.14</b>
Number of Species	16	14		25	21		27	25		35	28		25	25		52	45	
Total Number of Species		23			31			32			37			33			57	




Figure 1. Locations of ongoing Monitoring Avian Productivity and Survivorship (MAPS) bird banding stations at Yosemite National Park.

22 - The MAPS Program in Yosemite National Park, 2008




Figure 2. Population trends for 25 species and all species pooled at the **five currently operating MAPS stations** in Yosemite National Park over the 16 years 1993-2008. The index of population size was arbitrarily defined as 1.0 in 1993. Indices for subsequent years were determined from constant-effort between-year changes in the number of adult birds captured from stations where the species was a regular or usual breeder and summer resident. The annual percentage change in the index of adult population size was used as the measure of the population trend (APC), and it and the standard error of the slope (in parentheses) are presented on each graph. The correlation coefficient (r) and significance of the correlation coefficient (P) are also shown on each graph.



The MAPS Program in Yosemite National Park, 2008 - 23




Figure 2 (cont.). Population trends for 25 species and all species pooled at the **five currently operating MAPS stations** in Yosemite National Park over the 16 years 1993-2008. The index of population size was arbitrarily defined as 1.0 in 1993. Indices for subsequent years were determined from constant-effort between-year changes in the number of adult birds captured from stations where the species was a regular or usual breeder and summer resident. The annual percentage change in the index of adult population size was used as the measure of the population trend (APC), and it and the standard error of the slope (in parentheses) are presented on each graph. The correlation coefficient (r) and significance of the correlation coefficient (P) are also shown on each graph.




Figure 2 (cont.). Population trends for 25 species and all species pooled at the **five currently operating MAPS stations** in Yosemite National Park over the 16 years 1993-2008. The index of population size was arbitrarily defined as 1.0 in 1993. Indices for subsequent years were determined from constant-effort between-year changes in the number of adult birds captured from stations where the species was a regular or usual breeder and summer resident. The annual percentage change in the index of adult population size was used as the measure of the population trend (APC), and it and the standard error of the slope (in parentheses) are presented on each graph. The correlation coefficient (r) and significance of the correlation coefficient (P) are also shown on each graph.

The MAPS Program in Yosemite National Park, 2008 - 25




Figure 3. Trend in productivity for 25 species and all species pooled at the **five currently operating MAPS stations** in Yosemite National Park over the 16 years 1993-2008. The 1993 reproductive index was defined as the actual reproductive index; indices for subsequent years were determined from constant-effort between-year changes in the reproductive index. The slope of the regression line for annual change in the index of productivity was used as the measure of the productivity trend (PrT), and it and the standard error of the slope (in parentheses) are presented on each graph. The correlation coefficient (r) and significance of the correlation coefficient (P) are also shown on each graph.

26 - The MAPS Program in Yosemite National Park, 2008




Figure 3 (cont.). Trend in productivity for 25 species and all species pooled at the **five currently operating MAPS stations** in Yosemite National Park over the 16 years 1993-2008. The 1993 reproductive index was defined as the actual reproductive index; indices for subsequent years were determined from constant-effort between-year changes in the reproductive index. The slope of the regression line for annual change in the index of productivity was used as the measure of the productivity trend (PrT), and it and the standard error of the slope (in parentheses) are presented on each graph. The correlation coefficient (r) and significance of the correlation coefficient (P) are also shown on each graph.




Figure 3 (cont.). Trend in productivity for 25 species and all species pooled at the **five currently operating MAPS stations** in Yosemite National Park over the 16 years 1993-2008. The reproductive index was defined as the actual reproductive index value in 1993. Indices for subsequent years were determined from constant-effort between-year changes in reproductive index from stations where the species was a regular or usual breeder and summer resident. The slope of the regression line for annual change in the index of productivity was used as the measure of the productivity trend (PrT), and it and the standard error of the slope (in parentheses) are presented on each graph. The correlation coefficient (r) and significance of the correlation coefficient (P) are also shown on each graph.

28 - The MAPS Program in Yosemite National Park, 2008

Appendix I. Numerical listing (in AOU checklist order) of all the species sequence numbers, species alpha codes, and species names for all species banded or encountered during the 19 years, 1990-2008, of the MAPS Program on the six stations ever operated on **Yosemite National Park**.

Cumulative breeding status for all years in which each station was operated are also included (**B** = Regular Breeder (all years); **U** = Usual Breeder (>½, not all, years); **O** = Occasional Breeder (<½ years); **T** = Transient; **M** = Migrant; **A**= Altitudinal Disperser; **?** = Uncertain Species ID

NUMB	SPEC	SPECIES NAME	White Wolf (WHWO)	Gin Flat East Meadow (GFEM)	Crane Flat (CRFL)	Hodgdon Meadow (HODG)	Big Meadow (BIME)	Tamarak Meadow (TAME)
01010	GBHE	Great Blue Heron					T	
01300	TUVU	Turkey Vulture	T	T	T	T	T	
01630	MALL	Mallard		O		O	O	
01980	COME	Common Merganser					T	
02020	OSPR	Osprey					T	
02170	NOHA	Northern Harrier					T	
02200	SSHA	Sharp-shinned Hawk		T		T		
02210	COHA	Cooper's Hawk	T		T	O	T	
02240	NOGO	Northern Goshawk	T	T		T		
02380	RSHA	Red-shouldered Hawk	T		M	T		
02460	RTHA	Red-tailed Hawk	T	T	T	U	O	
02510	GOEA	Golden Eagle					T	
02545	UNHA	Unidentified Hawk				?	?	
02630	AMKE	American Kestrel					O	
03000	DUGR	Dusky Grouse	T	T	O	O		
03040	WITU	Wild Turkey				T	T	
03100	MOUQ	Mountain Quail	O	U	O	U	B	
03130	CAQU	California Quail				T	O	
03370	VIRA	Virginia Rail				T		T
05440	BTPI	Band-tailed Pigeon	T	T	T	O	T	
05570	MODO	Mourning Dove		T	T	O	O	
06670	WESO	Western Screech-Owl				T		
06800	GHOW	Great Horned Owl	T		T	O	T	
06830	NOPO	Northern Pygmy-Owl				T	T	
06940	SPOW	Spotted Owl				O		
06970	GGOW	Great Gray Owl	T	U	O	O		
07040	NSWO	Northern Saw-whet Owl				T		
07330	BLSW	Black Swift					T	
07410	VASW	Vaux's Swift				T	T	
07530	WTSW	White-throated Swift		T		T		
08640	BCHU	Black-chinned Hummingbird			T	T	T	
08670	ANHU	Anna's Hummingbird	T	O	O	U	U	T

## Appendix I, continued.

NUMB	SPEC	SPECIES NAME	WHWO	GHEM	CRFL	HODG	BIME	TAME
08690	CAHU	Calliope Hummingbird	T	O	O	O	O	T
08730	RUHU	Rufous Hummingbird	M	M	M	M	M	M
08740	ALHU	Allen's Hummingbird	M	M	M	M	M	
08774	USHU	Unidentified Selasphorus Hummingbird	?	?	?	?	?	
08775	UNHU	Unidentified Hummingbird	?	?	?	?	?	
09110	BEKI	Belted Kingfisher			T	T	U	
09390	LEWO	Lewis's Woodpecker					M	
09430	ACWO	Acorn Woodpecker	T		T		U	
09570	WISA	Williamson's Sapsucker	U	O	T	T		
09600	RBSA	Red-breasted Sapsucker	O	B	B	B	O	O
09640	NUWO	Nuttall's Woodpecker				T	T	
09650	DOWO	Downy Woodpecker	T		T	O	U	T
09660	HAWO	Hairy Woodpecker	U	U	U	U	U	B
09690	WHWO	White-headed Woodpecker	O	B	B	B	O	B
09710	BBWO	Black-backed Woodpecker						U
09800	RSFL	Red-shafted Flicker	U	B	U	B	B	U
09860	PIWO	Pileated Woodpecker	T	U	U	U	T	O
09915	UNWO	Unidentified Woodpecker	?					
11340	OSFL	Olive-sided Flycatcher	T	U	O	B	O	B
11380	WEWP	Western Wood-Pewee	U	U	O	B	B	B
11475	WIFL	Willow Flycatcher		T	T	U	O	T
11510	HAFL	Hammond's Flycatcher	O	U	U	U	T	O
11515	HDFL	Hammond's/Dusky Flycatcher		?	?	?		
11520	GRFL	Gray Flycatcher	M		M	M	M	
11530	DUFL	Dusky Flycatcher	B	B	B	U	T	B
11555	PSFL	Pacific-slope Flycatcher	T	O	O	U	O	T
11595	UEFL	Unidentified Empidonax Flycatcher	?	?	?	?	?	
11600	BLPH	Black Phoebe	T	T	T	O	B	
11740	ATFL	Ash-throated Flycatcher					O	T
12020	WEKI	Western Kingbird	T			T	T	
12085	UNFL	Unidentified Flycatcher	?	?	?	?	?	
12710	CAVI	Cassin's Vireo	T	O	B	B	U	U
12740	HUVI	Hutton's Vireo		T	O	O		
12760	WAVI	Warbling Vireo	U	U	B	B	B	B
12790	REVI	Red-eyed Vireo			M	M		
12920	STJA	Steller's Jay	B	B	B	B	U	B
13110	WESJ	Western Scrub-Jay	T			T	O	
13150	CLNU	Clark's Nutcracker	T	T		T		
13190	AMCR	American Crow		M		M		
13300	CORA	Common Raven	U	U	U	B	U	O

30 - The MAPS Program in Yosemite National Park, 2008  
 Appendix I, continued.

NUMB	SPEC	SPECIES NAME	WHWO	GHEM	CRFL	HODG	BIME	TAME
13410	TRES	Tree Swallow		T		T	T	T
13440	VGSW	Violet-green Swallow		T		T	O	T
13490	NRWS	Northern Rough-winged Swallow				T	O	
13540	BARS	Barn Swallow					O	
13555	UNSW	Unidentified Swallow					?	
13580	MOCH	Mountain Chickadee	B	B	B	U	O	B
13600	CBCH	Chestnut-backed Chickadee	T	T	T	O		T
13640	OATI	Oak Titmouse					O	
13680	BUSH	Bushtit			T	O	U	T
13690	RBNU	Red-breasted Nuthatch	B	B	B	B	O	B
13700	WBNU	White-breasted Nuthatch	T	O	O	O	O	O
13710	PYNU	Pygmy Nuthatch		T				
13730	BRCR	Brown Creeper	B	B	B	B	B	B
14040	BEWR	Bewick's Wren	T	T		T	O	
14070	HOWR	House Wren	A	A	A	A	U	A
14110	WIWR	Winter Wren	T	T	O	O	O	T
14205	UNWR	Unidentified Wren				?	?	
14210	AMDI	American Dipper					O	
14240	GCKI	Golden-crowned Kinglet	B	B	B	B	T	U
14250	RCKI	Ruby-crowned Kinglet	O			T		
14570	WEBL	Western Bluebird		T		O	U	
14590	TOSO	Townsend's Solitaire	T	O	O	T	T	
14810	SWTH	Swainson's Thrush	T			O		
14820	HETH	Hermit Thrush	B	O	U	U	T	T
15000	AMRO	American Robin	B	B	B	B	B	B
15110	WREN	Wrentit					O	
15370	EUST	European Starling				O	O	
15550	CEDW	Cedar Waxwing				M	M	
15660	OCWA	Orange-crowned Warbler	A	A	A	A	A	A
15670	NAWA	Nashville Warbler	A	A	A	B	U	A
15750	YWAR	Yellow Warbler	O	T	O	U	B	T
15800	AUWA	Audubon's Warbler	B	B	B	B	O	B
15800	YRWA	Yellow-rumped Warbler		T				
15810	BTYW	Black-throated Gray Warbler	T	T	T	O	O	T
15840	TOWA	Townsend's Warbler	M	M	M	M		M
15850	HEWA	Hermit Warbler	U	B	B	B	T	U
16040	AMRE	American Redstart				M		
16090	NOWA	Northern Waterthrush					M	
16140	MGWA	MacGillivray's Warbler	T	B	B	B	U	B
16150	COYE	Common Yellowthroat				M		
16280	HOWA	Hooded Warbler				M		
16290	WIWA	Wilson's Warbler	T	T	O	U	T	B



## Appendix I, continued.

NUMB	SPEC	SPECIES NAME	WHWO	GHEM	CRFL	HODG	BIME	TAME
16460	YBCH	Yellow-breasted Chat				T	T	
16495	UNWA	Unidentified Warbler			?	?	?	
16840	WETA	Western Tanager	O	B	B	B	U	B
17790	GTTO	Green-tailed Towhee		O	T	T	T	
17810	SPTO	Spotted Towhee			T	O	U	
17850	CALT	California Towhee					T	
18020	CHSP	Chipping Sparrow	U	T	U	U	U	B
18110	SAGS	Sage Sparrow					T	
18130	SAVS	Savannah Sparrow					M	
18140	GRSP	Grasshopper Sparrow					M	
18220	FOSP	Fox Sparrow	T	O	T	T	T	O
18230	SOSP	Song Sparrow	O	O	U	B	B	O
18240	LISP	Lincoln's Sparrow	B	B	B	B	O	B
18290	MWCS	Mountain White-crowned Sparrow	T			T		
18320	ORJU	Oregon Junco	B	B	B	B	U	B
18335	UNSP	Unidentified Sparrow		?	?	?	?	
18600	RBGR	Rose-breasted Grosbeak				M		
18610	BHGR	Black-headed Grosbeak	O	O	U	B	B	O
18660	LAZB	Lazuli Bunting	T	T	U	O	B	T
18670	INBU	Indigo Bunting			M	M		
18730	RWBL	Red-winged Blackbird	T	T	T	B	O	O
18810	WEME	Western Meadowlark					O	
18820	YHBL	Yellow-headed Blackbird					M	
18860	BRBL	Brewer's Blackbird	U	O	O	B	B	
18960	BHCO	Brown-headed Cowbird	O	T	O	U	U	
19105	BUOR	Bullock's Oriole		T		T	U	T
19330	PIGR	Pine Grosbeak	U	T	T			
19350	PUFI	Purple Finch	O	O	U	U	U	O
19360	CAFI	Cassin's Finch	U	O	O	O	O	O
19370	HOFI	House Finch			T	T	T	
19375	UCFI	Unidentified Carpodacus Finch			?	?	?	
19380	RECR	Red Crossbill	O	T	T	T	O	
19430	PISI	Pine Siskin	B	B	U	U	O	U
19490	LEGO	Lesser Goldfinch	T	O	T	O	B	T
19500	LAGO	Lawrence's Goldfinch		T	T	T	O	T
19510	AMGO	American Goldfinch				M	M	M
19580	EVGR	Evening Grosbeak	O	T	T	T	O	T
19920	HOSP	House Sparrow					T	
20085	UNBI	Unidentified Bird	?		?	?	?	?